FOOD HANDLING EQUIPMENT, TABLES, CONVEYORS, FLOORS, ETC., APPLY A SOLUTION OF 100 PPM TO 200 PPM AVAILABLE CHLORINE TO ALL SURFACES. (NEVER ALLOW AVAILABLE CHLORINE TO DROP BELOW 50 PPM). THIS MAY BE ACCOMPLISHED EITHER BY SPRAYING OR WIPING ON WITH A CLOTH OR BRUSH. LET STAND 15 TO 20 MINUTES, THEN RINSE THOROUGHLY WITH WATER. STORE THE SOLUTION ONLY IN GLASS, CROCKERY OR PLASTIC CONTAINERS. CLEAN SPRAY EQUIPMENT WITH WATER AFTER EACH USE.

DEGRADES WITH AGE. USE A TEST KIT AND INCREASE DOSAGE AS NECESSARY TO OBTAIN REQUIRED LEVEL OF AVAILABLE CHLORINE.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD.

#### CAUTION

HARMFUL IF SWALLOWED. DO NOT BREATHE VAPORS OR FUMES. DO NOT GET IN EYES, ON SKIN OR CLOTHING. IF SWALLOWED CALL PHYSICIAN IMMEDIATELY.

ANTIDOTE: INTERNAL — DRINK A TEASPOONFUL OR MORE OF MAGNESIA, CHALK OR SMALL PIECES OF SOAP, SOFTENED IN WATER, IN MILK OR RAW EGG WHITE. EXTERNAL — WIPE OFF GENTLY. IMMEDIATELY FLOOD THE SURFACE WITH WATER, USING SOAP FREELY, THEN COVER WITH MOIST MAGNESIA OR BAKING SODA.

CALL PHYSICIAN

KEEP OUT OF CHILDREN'S REACH

KEEP IN A COOL PLACE

• DAIRIES • POOLS

## D-CHLOF

POULTRY EQUIPMENT

KENNEL

### CAUTION: KEEP OUT OF REACH OF CHILDREN

EPA

USDA REG. NO. 8963-3 -/ 12/1

Manufactured by INTERMOUNTAIN SOAP AND CHEMICAL CO. Nampa, Idaho 83651

BULL O 1972

BULLEN DENTICIDE ACT
FI. NGICIUE AGU L'ODENTICIDE ACT
FI. RECINOME POLION REGISTER.
FI. UNDER NO. 2014 - 2. SUBJECT
TO ATTACHEU COMMENTS.

### T GAL.

#### DIRECTIONS

DAIRY EQUIPMENT: TREAT EQUIPMENT WITH 50 PPM TO 200 PPM (4 TABLESPOONS DISINFECTANT TO 8 GALLONS OF WATER PROVIDE 200 PPM) AFTER THOROUGHLY CLEANING WITH WATER AND CLEANER. CAUTION: TEST THE SOLUTION IN USE TO MAKE CERTAIN THAT THE CONCENTRATION DOES NOT FALL BELOW 50 PPM AT ANY TIME. IF A CONCENTRATION GREATER THAN 200 PPM IS USED, THEN THE EQUIPMENT MUST BE RINSED AFTER TREATMENT WITH POTABLE WATER.

POULTRY AND POULTRY EQUIPMENT: WASH INCUBATORS, BREEDERS, FEEDING UTENSILS, ETC., THEN APPLY A SOLUTION 2 TABLESPOONS DISINFECTANT TO EACH GALLON OF WATER. TO TREAT DRINKING WATER FOR FOWL, ADD 25 PPM DIS NFECTANT (1 TEASPOON DISINFECTANT TO 5 GALLONS WATER.)

KENNELS, DOGS, CATS: TO DEODORIZE AND CLEANSE DOGS AND CATS, FIRST CLEAN WITH WARM WATER THEN WASH WITH A SOLUTION OF 25 PPM TO

**DAIRIES** 

· POOLS

# D-CHLOR

POULTRY EQUIPMENT

KENNELS

**DAIRIES** 

• POOLS

# D-CHLOR

POULTRY EQUIPMENT

KENNELS

### AUTION: KEEP OUT OF REACH OF CHILDREN

EPA TUSBA REG. NO. 8963-3 -/39/3

Manufactured by
INTERMOUNTAIN SOAP AND CHEMICAL CO.
Nampa, Idaho 83651

TO 200 PPM (4 ABLESPOONS DISINFECTANT TO 8 GALLONS OF WATER PROVIDE 200 PPM) AFTER THOROUGHLY CLEANING WITH WATER AND CLEANER. CAUTION: TEST THE SOLUTION IN USE TO MAKE CERTAIN THAT THE CONCENTRATION DOES NOT FALL BELOW 50 PPM AT ANY TIME. IF A CONCENTRATION GREATER THAN 200 PPM IS USED, THEN THE EQUIPMENT MUST BE RINSED AFTER TREATMENT WITH POTABLE WATER.

POULTRY AND POULTRY EQUIPMENT: WASH INCUBATORS, BREEDERS, FEEDING UTENSILS, ETC., THEN APPLY A SOLUTION 2 TABLESPOONS DISINFECTANT TO EACH GALLON OF WATER. TO TREAT DRINKING WATER FOR FOWL, ADD 25 PPM DISINFECTANT (1 TEASPOON DISINFECTANT TO 5 GALLONS WATER.)

KENNELS, DOGS, CATS: TO DEODORIZE AND CLEANSE DOGS AND CATS, FIRST CLEAN WITH WARM WATER THEN WASH WITH A SOLUTION OF 25 PPM TO EACH GALLON OF WARM SUDSY WATER (1 TEASPOON OF DISINFECTANT TO 5 GALLONS OF WATER). RINSE WELL WITH CLEAR WATER, THEN DRY. FOR GENERAL SANITATION AROUND THE KENNEL, USE 2 TABLESPOONS PER GALLON OF WATER (800 PPM) TO DISINFECT AND DEODORIZE EQUIPMENT AND FLOORS, AFTER CLEANING WITH WARM, SUDSY WATER.

**HORSE TROUGH:** USE 25 PPM IN THE DRINKING WATER.

**SWIMMING POOLS:** USE 1½ PINTS OF DISINFECTANT TO 12,500 GALLONS OF WATER. ADD MORE AS NEEDED TO MAINTAIN 0.6 TO 1.0 PPM OF AVAILABLE CHLORINE IN THE WATER AT ALL TIMES.

KEEP OUT OF CHILDREN'S REACH

KEEP IN A COOL PLACE