UNDER THE ELLIPSE INSERTICIDE FUNGICIDE AND ROLENSE DE ACT FOR ECONOMIC POISON REGISTERED UNDER NORTH - 4 - 4

Cross-infection is of major housekeeping concern hot only in hospitals, but in schools, institutions, and industry. MALCO DISINFECTANT CLEANER DC-100 is formulated for this problem area. It both cleans and disinfects effectively when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection.

Two ounces of MALCO DISINFECTANT CLEANER DC-100 per gallon of water will deodorize bathrooms, garbage storage areas, and other areas where bacterial growth can cause mal odors.

WARNING

Keep out of reach of children. Causes severe eye and skin irritation. Do not get in eyes, on skin, or Harmful if swallowed. Avoid on clothing. contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink promptly a large quantity of water. Avoid alcohol. Call a physician immediately.



Sta Sáli

Active Ingredients

Did Tet Soc

Soc

Inert Ingredients

9019-4 **EPA Registration No.**

Net Contents:

KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR ADDITIONAL PRECAU-**TIONARY STATEMENTS.**

MALCO DISINFECTANT CLEANER DC·100

Cleaner-Disinfectant-Deodorizer-Fungicide for Hospital and Institutional Use

AOAC Phenol Coefficients

ph. aureus (ATCC No. 6538)	100
monella typhosa (ATCC No. 6539)	50

decyl dimethyl ammonium chloride trasodium ethylenediamine tetraacetate dium carbonate dium metasilicate, anhydrous	4.5% 2.0% 1.0% 0.5%
dium metasilicate, annyarous	0.5%

92.0% 100.00%

MALCO PRODUCTS, INC. BARBERTON, OHIO, U.S.A. 44203

WARNING

USE DIRECTIOI

Apply MALCO walls, floors and and bed frames areas, a prelimina

For Hospitals 8 gallon of water.

At two ounces ANT CLEANE aeruginosa, Stap Escherichia coli formance agains firmed by the performance wa AOAC Fungicid

For Schools, Ir ounce per gallor

At one ounce ANT CLEANE germicidal. It Salmonella chol interdigitale, th tests used abc hospitals were

Rinse empty (

+--HOE. ACT RED

ern not only in MALCO Justry. ed for this probtively when used n will reduce the

LEANER DC-100 :, garbage storage n cause mal odors.

ses severe eye es, on skin, or Avoid wed.

ih eyes or skin minutes. For and wash all If swallowed, water. Avoid γ.

MALCO DISINFECTANT CLEANER DC-100

Cleaner-Disinfectant-Deodorizer-Fungicide for Hospital and Institutional Use

AOAC Pheno' Coefficients

Stanh. aureus (ATCC No. 6538) Salmonella typhosa (ATCC No. 6539)	100
	50

Active Ingredients

Didecyl dimethyl ammonium chloride Tetrasodium ethylenediamine tetraacetate Sodium carbonate Sodium metasilicate, anhydrous	4.5% 2.0% 1.0% 0.5%
Inert Ingredients	92.0%
	00.00%
EPA Registration No. 9019-4	

EPA Registration No.

Net Contents:

MALCO PRODUCTS, INC. BARBERTON, OHIO, U.S.A. 44203

WARNING

KEEP OUT OF REACH OF CHILDREN SEE LEFT PANEL FOR ADDITIONAL PRECAU-TIONARY STATEMENTS.

USE DIRECTIONS:

Apply MALCO DISINFECTANT CLEANER DC-100 to walls, floors and other hard surfaces such as tables, chairs and bed frames with a cloth or mop. For heavily soiled areas, a preliminary cleaning may be required.

For Hospitals & Nursing Homes: Add two ounces per gallon of water.

At two ounces per gallon use-level, MALCO DISINFECT-ANT CLEANER DC-100 is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, and pathogenic fungi. Germicidal performance against the first four organisms has been confirmed by the AOAC Use-Dilution Test. Fungicidal performance was determined against T. interdigitale by the AOAC Fungicidal Test.

For Schools, Institutional and Industrial Uses: Add one ounce per gallon of water.

At one ounce per gallon use-level, MALCO DISINFECT-ANT CLEANER DC-100 delivers excellent cleaning and is germicidal. It is effective against Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, and Trichophyton interdigitale, the athlete's foot fungus. The same AOAC tests used above to confirm suitable performance for hospitals were used.

Rinse empty container with water before discarding.