

# CO-OP CATTLE MINERAL CONTAINS RABON® ORAL LARVICIDE

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	0.93% *
INERT INGREDIENTS	99.07% **
TOTAL	100.00%

\*Each pound contains 4.22 gm. Rabon

\*\*Refers only to ingredients which are not Larvicided

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

## ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

## GUARANTEED ANALYSIS

Calcium (Ca)	(max.)	16.00%
	(min.)	14.00%
Phosphorus (P)	(min.)	12.00%
Salt (NaCl)	(max.)	18.00%
	(min.)	16.00%
Chlorine (Cl)	(min.)	0.01%
Vitamin A	(min.)	250,000 units/lb.

## FEED INGREDIENTS

1. Dicalcium Phosphate, Salt, Calcium Formate, Calcium Hydroxide, Vitamin A Acetate, D-Calcium Pantothenate, Vitamin D3, Sulfur, Potassium Iron Sulfate, Magnesium Oxide, Magnesium Sulfate, Iron Carbonate, Iron Sulfate, Iron Oxide, Copper Oxide, Copper Carbonate, Ethylene Diamine Dihydrochloride and Zinc Oxide.

ACCEPTED

Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	0.93% *
INERT INGREDIENTS	99.07% **
TOTAL	100.00%

\*Each pound contains 4.22 gm. Rabon

\*\*Refers only to ingredients which are not Larvicided

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel for additional precautionary statements.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
CAUTION**

Harmful if swallowed. Avoid contact with skin or eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.


**GUARANTEED ANALYSIS**

Calcium (Ca)	(max) .....	16.00%
	(min) .....	14.00%
Phosphorus (P)	(min) .....	12.00%
Salt (NaCl)	(max) .....	18.00%
	(min) .....	16.00%
Iron (I)	(min) .....	.01%
vitamin A	(min) .....	280,000 units/lb

**FEED INGREDIENTS**

Includes Ethylene Salt, Concentrated Meal, Corn Meal, Hay, Vitamin A Acetate, D Active Vitamin Sterol (source of Vitamin D3), Sulfur, Potassium Iron Sulfate, Magnesium Oxide, Manganese Oxide, Iron Carbonate, Zinc Sulfate, Iron Oxide, Copper Oxide, Dental Carbonate, Ethylene Diamine Dihydrochloride and Zinc Oxide)

EPA Reg. No. 9078-6  
EPA Est. No. 9078-TN-4  
9078-TN-5  
Net Wt. 50 lbs.



Manufactured by  
**TENNESSEE FARMERS COOPERATIVE**  
LaVergne, Tennessee 37086

**(back of tag)**  
**DIRECTIONS FOR USE**

**GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

Feed Co-op Cattle Mineral at the rate of 2 ounces per 750 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

Start feeding Co-op Cattle Mineral early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

Co-op Cattle Mineral may be fed up to slaughter and to lactating dairy cow.

Co-op Cattle Mineral prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Co-op Cattle Mineral should be used in conjunction with other good management and sanitation practices.

**STORAGE AND DISPOSAL**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.