

## CO-OP PASTURE BLOCK WITH RABON<sup>®</sup>

To prevent the development of horn flies, face flies and stable flies in the manure of treated cattle

### ACTIVE INGREDIENT

2-Chloro-1 (2,4,5-Trichlorophenyl)  
Vinyl Dimethyl Phosphate

0.310

### INERT INGREDIENTS \*\*

99.690

\* Each pound contains 1.4 gms. Rabon

100.000

\*\* Refers only to ingredients which are not insecticidal.

### GUARANTEED ANALYSIS

Crude Protein	5.5% Min.
Crude Fat	0.5% Min.
Crude Fiber	10.0% Max.
Salt (NaCl)	19.0% Min. 24.0% Max.
Calcium	5.0% Max.
Phosphorus	2.5% Min.
Vitamin A, USP Units/lb	40,000 Min.
Vitamin D, USP Units/lb	5,000 Min.

### FEED INGREDIENTS

Cane Molasses, Roughage Products, Salt, Animal Fat, preserved with BHA (Butylated Hydroxyanisole), Calcium Carbonate, DiCalcium Phosphate, traces of Manganese Oxide, Iron Carbonate, Copper Oxide, Cobalt Carbonate, Calcium Iodate, Zinc Oxide.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE OTHER PRECAUTIONS ON REVERSE SIDE.

NET WT. 33 1/3 LBS. — 2 Blocks, weight 100 lbs.

EPA Reg. No. 9078-12

EPA Est. No. 9078-TN-4

9078-TN-5

Manufactured By  
TENNESSEE FARMERS COOPERATIVE  
LAVERGNE, TENNESSEE  
GENERAL OFFICE

## CO-OP PAS

### PRECAUTIONS HAZARDS

### CAL

Keep out of reach of children. Harmful to eyes. Avoid breathing dust. Wash throat and before eating or smoking. If in eyes, irritation persists, see a physician.

### ENVIRONMENT

This product is toxic to fish. Keep from contaminating water by cleaning of equipment.

### DIRECTIONS GENERAL CLASS

It is a violation of federal law to use this product in violation of the labeling.

1. Feed (approximately 1/2 lb.) of block of Pasture Block may be fed up to slaughter withholding the milk from market during treatment.
2. Place one block per five head of cattle.
3. Provide blocks where animals congregated.
4. Feed blocks continuously, never allow an empty block when each block has been one-half consumed.
5. Animals should not be starved for minor fly problems. PASTURE BLOCK program, but omit salt if a salt program is started. Do not feed free choice containing salt in any form.
6. When starting a feeding program during an infestation, use other measures to reduce the population. This product is not effective against existing populations. Additional measures may be needed in and around the feedlot to control adult flies and stable flies, which can breed in other decaying vegetable matter or sludge.
7. In order to achieve optimum fly control, this product should be used in conjunction with good management practices.

### STORAGE AND

Do not contaminate water or food by spillage. Do not reuse empty containers. Containers cannot be used, or chemically reprocessed, for approved pesticides. Dispose of in approved pesticide containers, or bury in a safe place.

## CO-OP PASTURE BLOCK

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE GENERAL CLASSIFICATIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Feed (approximately 1/2 lb.) of block per 1000 pound animal daily. Co-op Pasture Block may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.
2. Place one block per five head of cattle.
3. Provide blocks where animals congregate.
4. Feed blocks continuously, never allow animals to run out. Add an additional block when each block has been one-half consumed.
5. Animals should not be starved for minerals before initiation of the CO-OP PASTURE BLOCK program, but omit all salt from other feed after the program is started. Do not feed free choice sources of salt or minerals containing salt in any form.
6. When starting a feeding program during the fly season, it may be necessary to use other measures to reduce the population of existing adult flies. This product is not effective against existing adult flies. Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult flies and stable flies, which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.
7. In order to achieve optimum fly control, CO-OP PASTURE BLOCK should be used in conjunction with good management and sanitation practices.

### STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

9078-12