

10031

9078-12

CO-OP PASTURE BLOCK WITH RABON®

Control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

ACTIVE INGREDIENTS: Tetrachlorvinphos, 2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate 0.31%
INERT INGREDIENTS: 99.69%**
Total 100.00%

*Each pound contains 1.4 gm. Rabon
**Refers only to ingredients which are not Larvicidal Rabon is a registered trademark of Fermenta Animal Health Co

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash water.

GUARANTEED ANALYSIS

Crude Protein (min.) 5.50%
Crude Fat (min.) 0.50%
Crude Fiber (max) 10.00%
Calcium (min) 4.50% (max) 5.50%
Phosphorus (min) 2.50%
Salt (min) 19.00% (max) 22.50%
Vitamin A 40,000 IU/lb
Vitamin D-3 5,000 IU/lb

INGREDIENTS

Processed Grain By-Products, Salt, Dicalcium Phosphate, Cane Molasses, Roughage Products, Lignin Sulfonate, Calcium Carbonate, Plant Protein Products, Hydrolyzed Animal Fat and Hydrolyzed Plant Oil with Ethoxyquin (a preservative), Iron Oxide, Magnesium Oxide, Vitamin D-3 Supplement, Vitamin A Acetate, Vitamin E Supplement, Manganese Oxide, Copper Oxide, Cobalt Carbonate, Zinc Oxide, Ferrous Sulfate, Sodium Selenite, Calcium Iodate, and Ethylenediamine Dihydrochloride.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
1. Feed approximately 1/2 lb. of block per 1000 pound animal daily. Co-op Pasture Block may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.
 2. Place one block per five head of cattle.
 3. Provide blocks where animals congregate.
 4. Feed blocks continuously, never allow animals to run out. Add an additional block when each block has been one-half consumed.
 5. Animals should not be starved for minerals before initiation of the Co-op Pasture Block program, but omit all salt from other feed after the program is started. Do not feed free choice sources of salt or minerals containing salt in any form.
 6. When starting a feeding program during the fly season, it may be necessary to use other measures to reduce the population of existing adult flies. This product is not effective against existing adult flies. Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult flies and stable flies, which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.
 7. In order to achieve optimum fly control, Co-op Pasture Block should be used in conjunction with good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Storage: Store in a dry place in original container.
Disposal: Completely empty container into application (mixer/feeder) equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed, by State and local authorities by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

EPA Reg. No. 9078-12
EPA Est. No. 9078-TN-4 9078-TN-5
Net Wt. 33 1/3 lb.

Manufactured by
TENNESSEE FARMERS COOPERATIVE
LaVergne, Tennessee 37086
General Office

ITEM NO. 625
TAG CODE NO. 011290-625

