

Key # - 9086-5

PM - 10

193



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
9086-5

Date of Issuance:  
**23 JUN 1992**

NOTICE OF PESTICIDE:  
  X   Registration  
       Reregistration

(under FIFRA, as amended)

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Revenge Farm And Home  
Fly Bomb Insect  
Fogger

Name and Address of Registrant (include ZIP Code):

Roxide International, Inc.  
Box 249  
New Rochelle, NY 10802

Note: Changes in labeling differing from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 9086-5".
- b. The following Environmental Hazards statement must appear on the label: ENVIRONMENTAL HAZARDS-This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Apply this product only as specified on this label.
- c. In the food Handling caution note that the word, application, appears twice.

Signature of Approving Official:

Date: **23 JUN 1992**

page 2  
EPA Reg. No.

3. Submit one (1) copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

RFM

Richard F. Mountfort  
Product Manager 10  
Insecticide & Rodenticide Branch  
Registration Division (H7505C)



BEST AVAILABLE

# REVENGE

## FARM AND HOME FLY BOMB INSECT FOGGER

**Kills & Repels Flying Insects Crawling Insects (See Full List on Back)  
For Dairy Farms, Milk Houses, Stables, Hog Operations, Homes, Commercial Plants, on Livestock and more as listed on back.**

Active Ingredient	
Pyrethrins	0.50%
Piperonyl butoxide Tech*	1.00%
Bicycloheptene Dicarboximide	1.67%
Inert Ingredients	96.83%
Equivalent to 0.8% ( Butyl- carbityl 6 Propylpiperonyl) Ether and 0.2% related compounds	

ACCEPTED FOR COMMERCIAL SALE BY EPA LISTED 23 JUN 1992 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registration under EPA Reg. 9006-5

EPA Reg 9086-00 - EPA Est 4391-OH-2

Keep Out of Reach of Children

### CAUTION

See Additional Precautions On Back Panel

- IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration pneumonia hazard.
- IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
- IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

Net Weight \_\_\_\_\_ Ounces

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**POULTRY OPERATIONS:** Use only in empty poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. Treat daily or as necessary.

**HOG OPERATIONS:** In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after application. Ventilate the area after treatment is completed. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**KENNELS AND CATTERIES:** Remove animals before spraying. Ventilate after treatment is completed.

**HOME USES:** Follow Commercial Directions below

#### COMMERCIAL USES:

**OUTDOOR CONTROL:** Flies, Gnats and Mosquitoes in open areas near building and campgrounds. Best results attained when wind speed is five (5) MPH or less. Apply at a rate of 60 - 80 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

**INDOOR CONTROL:** Flying Insects - Cluster Flies, Flies, Fruit Flies, Gnats, Mosquitoes, Small Flying Moths. **GENERAL VOLUMETRIC TREATMENT:** Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. Fill room with mist. Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat treatment as necessary.

**Crawling Insects:** - Roaches, Ants, Spiders, Crickets, Boxelder Beetles, Silverfish. Spray into cracks and crevices, behind sinks, cabinet bottoms, under appliances, etc. **GENERAL VOLUMETRIC TREATMENT:** Open cabinets and drawers, close off air conditioners, fans and close doors or windows before spraying. Spray at rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary. For Use in and around bakeries, beef cattle operations, bottling plants, dairies, farms (including milk house, milk parlor, loafing shed and hog operations, homes, horse barns and stable, kennels, meat packing plants, poultry operations other industrial building and warehouses, and swine.

**Kills and Repels** ants, centipedes, cluster flies, crickets, deer flies, horn flies, hornets, house flies, lice, mosquitoes, roaches, sow bugs, etc.

**CALL US TOLL FREE FOR EXTRA TIPS AT 1-800-431-5500**

**DAIRY FARM USE:** Milking Parlor and Milk Room - Close all windows and doors. Spray upward and in all directions at a rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after treatment. Open and ventilate before reoccupying to prevent contamination from spray and falling dead insects.

**ANIMAL USE:** (Do not spray sick animals) - From approximately 10 feet, spray entire animal as it is being released to pasture. Do not spray directly into the face or eyes. Spot treat withers, shoulders and back where saliva accumulates.

**STANCHION BARN USE:** Walk behind animals and direct spray at the head. For most effective results, apply each morning.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**HORSE BARN AND STABLES:** Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per horse and pony. Do not spray directly toward the face or eyes. Repeat application daily or as necessary.

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

#### CAUTION:

Harmful if swallowed or absorbed through skin. Do not remain in treated area after treatment is completed. Avoid breathing of vapors. Wash thoroughly after using and before smoking or eating food, or foodstuffs. Remove all pets and birds. Cover fish tanks. **Food Areas of Food Handling Establishments:** Foods should be covered during treatment. All food processing surfaces should be covered during treatment with an effective cleaning compound followed by a potable water rinse to prevent contamination. When using this product as a space spray, operation. Except in Federally inspected meat and poultry processing plants, application for inspection purposes is permissible. **PHYSICAL OR CHEMICAL HAZARDS:** Contents under pressure. Do not use or store near heat or open flame. Exposure to heat may cause bursting. Do not spray on plastic, painted or varnished surfaces. Do not use on electronic equipment such as radios, televisions, computers, etc.

**STORAGE AND DISPOSAL:** **Storage:** Store in a cool area away from heat or open flame. **Disposal:** Replace cap and discard empty container in trash. **ABSOLUTE SATISFACTION:** Anything less, and you get your money back. For tips or questions call Toll Free 1-800-431-5500. Weekdays, 9 - 5 P.M. Eastern Time. Consumer Relations. ROXIDE INTERNATIONAL, INC., P.O. Box 10802, USA Mfg. for Roxide International, Inc. STOCK #30025