



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (M7505C)
401 M St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

9086-8

Date of Issuance:

11 FEB 1993

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Revenge Farm & Home
Fly Bomb Insect
Fogger

Name and Address of Registrant (include ZIP Code):

Roxide International, Inc.
Box 249
New Rochelle, NY 10802

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar product to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 9086-8".
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Joseph M. Parsons

Date:

February 11, 1993

2 for 2

REVENGE FARM AND HOME FLY BOMB INSECT FOGGER

Kills & Repels Flying Insects Crawling Insects (See Full List on Back)
For Dairy Farms, Milk Houses, Stables, Hog Operations, Homes, Commercial Plants, on Livestock and more as listed on back.

Active Ingredient:
Pyrethrins 0.50%
Piperonyl butoxide Tech* 5.00%
Inert Ingredients 94.50%
* Equivalent to 0.4% (Butyl- carbityl 6 Propylpiperonyl) Ether and 1% related compounds.
Contains Petroleum Distillate

EPA Reg 9086- EPA Est. 9551-OH-2

Keep Out of Reach of Children

CAUTION

See Additional Precautions On Back

IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration pneumonia hazard.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated

Net Weight _____ Ounces

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

POULTRY OPERATIONS: Use only in empty poultry houses (when birds are not present.) Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 Minutes after application. Ventilate the area after treatment is completed. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

KENNELS AND CATTERIES: Use on animals use - spray against natural lay of hair. Do not spray toward face or eyes. As a space spray - Remove animals before spraying. Ventilate before re-occupying. Not recommended for use on kittens. or puppies less than four weeks old.

HOME USES: Follow Commercial Directions below.

COMMERCIAL USES:

OUTDOOR CONTROL: Flies, Gnats and Mosquitoes in open areas near building and campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60 - 80 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

INDOOR CONTROL: Flying Insects - Cluster Flies, Flies, Fruit Flies, Gnats, Mosquitoes, Small Flying Moths. **GENERAL VOLUMETRIC TREATMENT:** Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. Fill room with mist. Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat treatment as necessary.

Crawling Insects: - Roaches, Ants, Spiders, Crickets, Boxelder Bugs, Earwigs, Grain mites, Silverfish. Spray into cracks and crevices, behind sinks, cabinets, baseboards and window frames,

GENERAL VOLUMETRIC TREATMENT: Open cabinets and air conditioners, fans and close doors or windows before spray. Keep area closed for 15 minutes. Open and up and destroy fallen insects. Repeat treatment as necessary. For Use in and around bakeries, beef cattle operations, bottling plants, dairies, farms (including milk house, milk parlor, loafing shed and operations, homes, horse barns and stable, kennels, meat markets, poultry operations other industrial building and warehouses, swine, cats and dogs.

Kills and Repels ants, centipedes, cluster flies, crickets, deer flies, horn flies, house flies, lice, mosquitoes, roaches, scorpions, ticks, wasps.

CALL US TOLL FREE FOR EXTRA TIPS AT 1-800-431-5500

DAIRY FARM USE: Milking Parlor and Milk Room - Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after treatment. Open and ventilate before reoccupying. Repeat application daily or as necessary to prevent contamination from spray and falling dead insects.

ANIMAL USE: (Do not spray sick animals) - From approximately 10 feet. Do not spray directly at animal. Do not spray entire animal as it is being released to pasture. Do not spray treat withers, shoulders and back where saliva accumulates.

STANCHION BARN USE: Walk behind animals and direct spray at the back of the animal. For most effective results, apply each morning.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Repeat application daily or as necessary. Repeat application daily or as necessary.

HORSE BARN AND STABLES: Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Repeat application daily or as necessary. For the backs of horses and ponies from a distance of at least 10 feet. Repeat application daily or as necessary.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION:

Harmful if swallowed or absorbed through skin. Do not remain in treated area until after treatment is completed. Avoid breathing of vapor. Avoid contact with eyes. Wash thoroughly after using and before smoking or eating foodstuffs. Remove all pets and birds. Cover fish tanks or reefs. Do not use in Food Areas of Food Handling Establishments: Foods should not be prepared or served in treated areas. All food processing surfaces should be covered and cleaned with an effective cleaning compound followed by a potable water rinse. When using this product as a space spray, application in homes, schools, day care centers, and other public places for pest control purposes is permissible at all times.

PHYSICAL OR CHEMICAL HAZARDS: - Extremely flammable liquid and vapor. Do not use or store near open flames or incinerate container. Do not use or store near temperatures above 130°F may cause bursting. Do not spray on or directly into any electronic equipment such as radios, televisions, etc.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds, and bees. Do not apply directly to water. Apply this product only as specified on this label.

STORAGE AND DISPOSAL: Storage: Store in a cool, dry place. Disposal: Replace cap and discard empty container in trash.

ABSOLUTE SATISFACTION: Anything less, and you get your money back. For tips or questions call Toll Free 1-800-431-5500. Weekdays, 9 AM to 5 PM. Consumer Relations. ROXIDE INTERNATIONAL, INC., P.O. Box 10602, USA Mfg. for Roxide International, Inc.

Approved with comments in EPA Letter dated 11 FEB 1993 Under the Federal Insecticide, Fungicide, and Rodenticide Act 9086-8