

PM 03

9086-P

5/15/97

P9 172

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 05-31-98



United States Environmental Protection Agency  
Washington, DC 20460

Registration  
Amendment  
Other

OPP Identifier Number

251655

Application for Pesticide - Section I

1. Company/Product Number 9086-8	2. EPA Product Manager MARION JOHNSON	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) REVENGE FARM AND HOME FLY BOMB INSECT FOGGER	PM# 10	
5. Name and Address of Applicant (Include ZIP Code) ROXIDE INTERNATIONAL, INC. P.O. BOX 249 NEW ROCHELLE, NY 10802  <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

ADDITIONAL STATEMENT OF THE PRECAUTIONARY STATEMENT SECTION: "PROLONGED OR FREQUENTLY REPEATED SKIN CONTACT MAY CAUSE ALLERGIC REACTION IN SOME INDIVIDUALS."

Notification  
MAY 15 1997

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
Certification must be submitted		If "Yes" Unit Packaging wgt.    No. per container If "Yes" Package wgt        No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name JAMES B. COWEN	Title PRESIDENT	Telephone No. (Include Area Code) 914-235-5300
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>James B. Cowen</i>	3. Title PRESIDENT	
4. Typed Name JAMES B. COWEN	5. Date APRIL 28, 1997	

# REVENGE FARM AND HOME FLY BOMB INSECT FOGGER

Kills & Repels Flying Insects Crawling Insects (See Full List on Back)  
For Dairy Farms, Milk Houses, Stables, Hog Operations, Homes, Commercial Plants, on Livestock and more as listed on back.

### Active Ingredient:

Pyrethrins ..... 0.50%  
Piperonyl butoxide Tech\* ..... 5.00%  
Inert Ingredients ..... 94.50%  
\* Equivalent to 0.4% ( Butyl- carbityl 6 Propylpiperonyl) Ether and 1% related compounds,  
Contains Petroleum Distillate

EPA Reg 9086- EPA Est. 9591-OH-2

Keep Out of Reach of Children

## CAUTION

See Additional Precautions On Back Panel

IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration pneumonia hazard.  
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.  
IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.  
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

Net Weight \_\_\_\_\_ Ounces

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**POULTRY OPERATIONS:** Use only in empty poultry houses (when birds are not present.) Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. Treat daily or as necessary.

**HOG OPERATIONS:** In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after application. Ventilate the area after treatment is completed. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**KENNELS AND CATTERIES:** On animals use - spray against natural lay of hair. Do not spray toward face or eyes. As a space spray - Remove animals before spraying. Ventilate before re-occupying. Not recommended for use on kittens, or puppies less than four weeks old.

**HOME USES:** Follow Commercial Directions below.

### COMMERCIAL USES:

**OUTDOOR CONTROL:** Flies, Gnats and Mosquitoes in open areas near building and campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

**INDOOR CONTROL:** Flying Insects - Cluster Flies, Flies, Fruit Flies, Gnats, Mosquitoes, Small Flying Moths. **GENERAL VOLUMETRIC TREATMENT:** Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. Fill room with mist. Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat treatment as necessary.

**Crawling Insects:** - Roaches, Ants, Spiders, Crickets, Boxelder Bugs, Earwigs, Grain mites, Silverfish. Spray into cracks and crevices, behind sinks, cabinets, baseboards and window frames.

**GENERAL VOLUMETRIC TREATMENT:** Open cabinets and doors in area to be treated. Shut off air conditioners, fans and close doors or windows before spraying. Apply at a rate of 20 seconds per 1000 cu. ft. Keep area closed for 15 minutes. Open and ventilate before re-occupying. Sweep up and destroy fallen insects. Repeat treatment as necessary.

For Use in and around bakeries, beef cattle operations, bottling plants, breweries, catteries, dairy farms (including milk house, milk parlor, loafing shed and holding lot), food storage areas, hog operations, homes, horse barns and stable, kennels, meat packing and food processing plants, motels, poultry operations other industrial building and warehouses. For direct use on cattle, horses, swine, cats and dogs.

Kills and Repels ants, centipedes, cluster flies, crickets, deer flies, face flies, fleas, fruit flies, gnats, horn flies, hornets, house flies, lice, mosquitoes, roaches, sowbugs and spiders.

CALL US TOLL FREE FOR EXTRA TIPS AT 1-800-431-5500 (WEEKDAYS, 9-5 EST).

**DAIRY FARM USE:** Milking Parlor and Milk Room - Close all windows and doors. Direct fog upward and in all directions at a rate of 1 to 2 seconds per 1000 cu. ft. Keep room closed for 15 minutes after treatment. Open and ventilate before reoccupying. Cover milking utensils and milk to prevent contamination from spray and falling dead insects.

**ANIMAL USE:** (Do not spray sick animals) - From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

**STANCHION BARN USE:** Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double dosage. For animal use thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**HORSE BARN AND STABLES:** Close all windows and doors. Spray at a rate of 1 to 2 seconds per 1000 cu ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least two feet. Do not spray directly towards the face or eyes. Repeat application daily or as necessary.

## PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

### CAUTION:

Harmful if swallowed or absorbed through skin. Do not remain in treated area and ventilate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Do not contaminate water, food, or food-stuffs. Remove all pets and birds. Cover fish tanks or remove before spraying.

**Food Areas of Food Handling Establishments:** Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. When using this product as a space spray, apply only when plant is not in operation. Except in Federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all times.

**PHYSICAL OR CHEMICAL HAZARDS -** Extremely flammable. Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, etc.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish, birds and other wild life. Do not apply directly to water. Apply this product only as specified on this label.

**STORAGE AND DISPOSAL:** Storage: Store in a cool area away from heat or open flame. Disposal: Replace cap and discard empty container in trash.

**ABSOLUTE SATISFACTION:** Comments or questions? Call 1-914-235-5300, Weekdays, 9-5 Eastern Time. Or write Jim Barry, Consumer Relations, ROXIDE INTERNATIONAL, INC., P.O. Box 249, New Rochelle, New York 10802. USA Mfg. For Roxide International, Inc.

0.50%  
5.00%  
94.50%  
EPA LETTER DATED  
11 FEB 1993  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under 9086-8

NOTIFICATION  
1997

2/2