03/30/2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENC WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAR 3-0 2009

- 6 H.

Carrie Crawford Product Registration Carrie Crawford, Regulatory Consultant Agent for Roxide International, Inc. 5927 Paint Bank Rd. New Castle, VA 24127

SUBJECT:

Application for Pesticide Notification (PRN 98-10) Request Alternate Brand Name "Revenge Fly Bomb" EPA Reg. No. 9086-8 Application Dated December 17, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 12/17/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" received but not reviewed and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

 \sim

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before completing form. United States Environmental Protection A Washington, DC 20460			Agency		A. OMB No. 2070 Registratio Amendmei Other	
	A	oplication	for Pesticide -	Section	I (
1. Company/Product Number 9086-8			2. EPA Product Manager / Richard Gebken			3. Proposed Classification
4. Company/Product (Name) Revenge Farm and Home Fly Bomb Insect Fogger			PM# []			
 5. Name and Address of Applicant (Include ZIP Code) Roxide International, Inc. P. O. Box 239 Larchmont, New York 10538 Check if this is a new address 			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. NOTIFICATION Product Name MAR 3 0 2009			
			Section - II			
Amendment - Explain below Resubmission in response to Notification - Explain below. Explanation: Use additional page	o Agency letter da		Age *Me	printed labe ncy letter da Too" Applic r - Explain b	ation.	· · · · · · · · · · · · · · · · · · ·
Notification of Alternate Brand Name This notification is consistent with the labeling or the confidential statement EPA. I further understand that if this FIFRA and I may be subject to enfor	e provisions of PR I t of formula of this p notification is not o	Notice 98-10 a product. I und consistent with	nd EPA regulations at 4 erstand that it is a violati the terms of PR Notice er sections 12 and 14 of	0 CFR 152.4 on of 18 U.S 98-10 and 40	6, and no other cha .C. Sec. 1001 to wi	lifully make any false statement to
1. Material This Product Will Be Pac			Section - III			
Child-Resistant Packaging Yes No Ves No Certification must De submitted Ves No If "Yes" Unit Packaging Yes No No No No No No No No No No			Yes No If "Yes" No. per Package wgt container		GI Pa	etal astic ass per her (Specify)
3. Location of Net Contents Information 4. Size(s) H			stall Container 5. L			
			ngraph Other r glued ciled			
			Section - IV		, 	
1. Contact Point <i>(Complete items directly below for identificati</i> Name Carrie Crawford			ion of individual to be contacted, if necessary, to p Title Regulatory Agent			s this application.) phone No. (Include Area Code))-864-8490
I certify that the statements I I acknowledge that any know both under applicable law.			attachments thereto		imprisonment or .	C (Stamped)
2. Signature Chassie Gaugerd			Regulatory Agent			
4. Typed Name			5. Date			

. . .

READ ENTIRE LABEL BEFORE EACH USE Commercial and home use USE ONLY ON BEEF CATLE, HORSES, PONIES, SWINE, CATS AND DOGS

See one of the former, non-less rolling, rolling

Sensitivities may occur after using any pesticide product for pets. It signs of sensitivity occur, battle your pet with mild seap and rince with darge anomatis of water. Consult a veterinariant before using this product on depikitated, incucated, aged, pregnant or nursing animals. se with

llowed, initialed, in eyes, or on skin of animals, call a veterinarian. for treatment advice. tor incommittion on this product including health concerns, medical emergencies or pesticida incidents with animals, call ASPCA Animal Poison Control Center Hotline at 1-888-426-4435

Poison Control Center Hotline at 1-883-425 -455. ENVIRONMENTAL BAZABOS To protect the environment, do not allow possible to enter or run of this storm drams, drainage diches, guithers or surface waters. Applying this product in calm wrather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or watch pesticide off the treatment area. See Directions for Use for additional processions and reindraments reguirer

PHYSICAL OR CHENICAL HAZARDS: Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or individent container. Expositive to temperatures above 1907 may cause bursting. On not apply this product around electrical equipment due to the possibility of a shock hazard.

FIRST AID

 Immediately call a poison control center or doctor.
 Do not induce vomiting unless told to do so by a poison control. center or doctor

center or doctor. = Do not give any liquid to the person. = Do not give any liquid by mouth to an unconscious person. **If on Skin or Chothlag:** = Take et foontaminated clothing. = Paines exin immediately with plenty of water for 15-20 minutes. = Daila poison control center or doctor for tréatmeni advice.

f Inholect

n maxeu Move person to fresh air. «If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

1 in Eves: How yes.
 Hold eye open and rinse slowly and gently with water for 15-20 ming
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note to Physician: Contains petroleum distillate - vomiting may cause aspetelon oneumonia. For information of this product (infeuding health concerns, medical emergencies or pestickle incidents) call the informal residue information Center at 1-800-589-7378.

STORAGE AND DISPOSAL

STORABE AND DISPOSAL: Storage: Store in a cool area away from heat or open flame: Disposal: Do not puncture or incineratal if entpy: Norrelitable container. Do not reuse or refill this container. Offer for resystemic if available. If party filled: Call your local solid waste agency or toil free 1-800-CLEANUP and/or website www.earth91 long for disposal.

instr

QUESTIONS OR COMMENTS? Call 1-914-235-5300 Weekdays, 9-5 Eastern Time or write to Jim Barry Consumer Relations, Rodde International, Inc., P. O. Box 249, New Rochelle, NY 10802 USA

75

STOCK# 31725 Manufactured by: Roxide International, Inc.

76807"31725 P. O. Box 249 New Rochelle, NY 10802 EPA REG. #9086-8 EPA EST. #5693-MA-1

 $q(A, \gamma)$

1 100 200

Y'R(

RIDOOR OR FOR

FARM & HOME **INSECT FOGGER**

KILLS FLYING AND CRAWLING INSECTS

Including ants, centipedes, cluster flies, crickets, deer flies, face flies, fruit flies, gnats, horn flies, hornets, house flies, lice, mosquitoes, cockroaches, sow bugs, spiders

LITE'N DRY OIL BASED FORMULA

For Use in or around Bakeries, Bottling Plants, Brewer Meat Packing and Food Processing Plants, Homes, Hotels, Motels, Dairy Farms including Milk House, Milk Parlor, Loofing Shed and Holding Lot, Food Storage Areas; Hog Operations, Commercial Plants, Industrial Buildings Warehouses, Poultry Operations, Kennels and Catteries Stanchion Barns, Beef Cattle Operations, Horse Barns and Stables, On Livestock Including: Cattle, Horses, Swine, Cats and Dogs

KEEP OUT OF

Equivalent to 4.0% (butkarbity) propylpiperany() ether and 1.0% related compounds. Contains Petroleum Distillate

(See Side/Back for Additional Precautionary Statements) NET WT. 1 LB. 1 OZ ...

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling

product in a manner inconsistent virus is tooening. General Precoursons and Restructions: Do not armain in treated areas, Except when applying cirectly to dogs and cats, remove puts. No not allow adults, children or pats to enter the treated area unit sprays have dried. Do not apply his product in a vary that will contact adults, children or pats offer drivently or through drift. For indoor use, vehicle area through with treatments commercial. For indoor use, vehicle area through the treatment is commercial. In one contaminate water, food or toodstuffs. Remove pets and birds and community value, you or buckstars remove person and buck and over or remove fails tanks buckle spraying. To not apply bucketly to or near water, storm drains, or drainage dutches. Do not apply when windy. To prevent product run-off, do not over water the treated area or apply prior to teary randial.

OUTDOOR CONTROL: Flies and Gnats in open areas near buildings OUTDODE CONTROL: Fues and Grass in open areas near buildings and campgrounds and Mosquides in the garden. Hest insults are obtained when wind speciel is five (5) MPH or tess. Apply at a rate of 1;2 seconds per 1,000 sp. n. Spray in white events across area to be treated. Allow spiry diff to penetrate dense foliage. Repeat treatment as allocessary. BLOODS CONTROL: Frying insects - Cluster Flies, Flies, Fruit Flies, Grats, Mosquitues in the home, Sneil Frying Motes. CELERAL VOLUMETRIC TREATMENT: Corea & windows and doors . and direct fog upward at a min of 1 to 3 seconds per 1,000 cu. R². Fill room with mist. Disperse In all locations. Keep area dosed for 15 min. Open and worklade before neccopying. Sweep up and dispose of fallen insects. Repet treatment as necessary.

Or hald rescal. Kepes treatment is necessary
 FOOD AREAS OF FOOD MAINLING ESTABLISHMENTS
 FOOD AREAS OF FOOD MAINLING ESTABLISHMENTS
 FOOds must be manwed or covered during treatment or thoroughly
 cleaned with an effective cleaning compound tobived by a potable
 what rise aprices must be covered during treatment. All hood
 processing surfaces aprix, apply only when plants is not in operation.
 Event in forderupt in precide meet and pourty plants careful crack
 and crivice application is permissible at all times.
 Diality FABM USE- MILING Perfor and MILIN Room - Cose all windows
 of doors. Cover milling uterials and in this prevent contamination
 from spray and tailing deal insects. (Direct fog upwerd and in all
 directions at a rate of 1 to 2 seconds per 1000 ou. h. Keep room
 cload of 1 is a second spraitment. Open and versitable before
 recoccypring.
 POR USE OI BEEF CATTLE, MINISES, PONIES, SWINE, CATS & DOGS
 from approximation 2.1. distance, thoroughly spray end semination

The provingely 2 it. distance, throughly prime animal as it, and provingely 2 it. distance, throughly prime animal as it, eing reases to pasture. Do not prive directly toward animals s. Soch treat withen, shoulders and back where safiva unulates from head tossing." Do not spray sick animals.

eccimitates from head tossing: Do not spray eick animals; STANCHION EARN USE: Wait betrind animals and direct spray over backs allowing accord per annual. Reapply acche morting. REEF CATTLE OPERATIONS: In barris, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Ventilatió aréa after treatment is completa; if area is not closed, cuble the dosage. Neap area closed for 15 minutes tollowing application. For animal use, horoughly spray entire annual all is it being relased to passime. Do not spray directly loward animals face or eyes. Repeat every day.

Non starty carbody carbody analysis had to reveal, respect very con-NORSE RANIS AND STARLESS. Coose all windows and doon." Spring at a rate of 1 to 2 seconds per 1000 cir. In. Kesp area closed for 15 minutes following application. Ventilate area after treatment is complete. If area carbod be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least two feet. To one spring friendly towards the face or eyes. Reapply every morning.

office. Incappy and present loss only in empty poultry houses (when birds are not present). Close all windows and doors. Surray at a rate of 1 to 2 seconds per 1000 cci. It. Keep area closed for 15 minutes following application. Venultae area after treatment is complete. If area cannot be closed, doubte the dosage. Treat daily.

Also sciently to detect, double the bodger that lead, HOG OPERATIONS. In high houses, close all windows & doors. Spray at a rate of 1 to 2 seconds per 1000 cu. ft. Keep and closed for 15 mins following application. Ventilate area after treatment is completes, if area cannot be closed, double the dosage. For animal application; direct spray over backs and spray for 1 to 2 seconds per tog. Do not spray directly toward animals lace or eyes. Reapply every day. KENNELS AND CATTERIES: As a space spray - Except when applying directly to dogs and cats, remove animals before spraying. Ventilate

bislore moccupying. USE QII DOUS AND CATRS. Spray against natural key of hair. Do noi: spray lowerd hour or yers. Use only on dogs and catss. Repeat every day, Do not use on kittems or pupples less than histive weeks old.

NOTIFICATION

MAR 3 0 2009



ACTIVE INGREDIENTS:

0.50% . . . Total 100.00%