MORE FUN / LESS WORK FOR POOL OWNERS



**DANGER:** KEEP OUT OF REACH OF CHILDREN. See other precautions and anti-dote on side panel.

Active Ingredient Trichtoro s-Triazinetrione Incrt Ingredient

(Available chlorine 86, 5%)

95%

50

ACCEPTED 9087-/ JUN 1 1983

Under the Federal Invanticule, Funginide, and Railand in Act, as an inced, for the pessence reputer 1 under Phi Reg. No.

## DIRECTIONS FOR USE:

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT'S LABELING

TABEX. Slow Dissolving Chlorination Tablets are an extremely concentrated frim of swim ining pool sanitizing agent. TABEX. Other nation Tablets are designed to dissolve slowly and at a uniform rate thereby eliminating the risidance of frequent hand feeding of granutar or figure chlorination. We suggest the use of the TABEX. Aquarian Feeder for the oitimate convenience of continuous, automatically dispensed chlorination.

## FOR CONTINUOUS DAYLY CHLORINATION

Before the first use of TABEX\_SLOW\_DISSOLVING\_CHLORINATION\_TABLETS\_Le sure to take the following steps.

- A Adjust the poor water pH to between 7.2 and 7.8 as determined by the TABEX Swimming Pool Test Kit. Further adjust your Total Alkalinity to a minimum of 100 ppm (parts per million).
- B. Stabilize pool water to at least 50 ppm with TABEX Conditioner as directed on container label. Thereafter regular use of TABEX Slow Dissolving Chlorination Tablets for daily chlorination will provide any needed makeup Conditioner.
- C. Superchlorinate (SHOCK\_TREAT) your pool water to a 4 ppm residual chlorine level using TABEX Super Shock Tablets as directed on the container tabel.
- D. Once your pool has been made "ready" by following A, B & C above, use these Slow Dissolving Chlorination Tablets to maintain the free residual chlorine level at a safe and efficient 1.0 to 1.5 ppm of FREE available chlorine at all times, as determined by the DAILY or more frequent use of the TABEX 4.1 Swimming Pool Test Kit.
- E. By means of the TABEX Aguarian Feeder, TABEX Floater or TABEX Skimmer Dispenser, immerse one TABEX Slow Dissolving Chlorination Tablet (or as many of them as may be neces Sary to maintain the safe FRCE available chlorine lever) for each 5 000 to 10 000 gallons (18,920 to 37,840 liters) of pool water.
- NEVER ADD THESE SLOW DISSOLVING CHLORINATION TABLETS DIRECTLY INTO THE WATER AS DISCOLORATION OF THE POOL LINER OR FINISH WILL OCCUR.
- F. Superchlorinate (SHOCK TREAT) your pool water at least twice monthly (more often in Sunbelt regions) during hot weather and peak bather loads. Failure to Superchlorinate regularly could result in the growth of Algae in your pool. To control algae, use TABEX Algae Out as directed on the container label.





EPA Reg. No. 9087-1 EPA Est. No. 3087-NY-1-2

**DANGER:** Highly corrosive. Causes skin and eye damage. May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Irritating to nose and throat. Avoid breathing dust. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not discharge into lakes, streams, ponds, or public waters unless in accordance with an NPDES Permit. For guidance contact the regional office of EPA.

PHYSICAL AND CHEMICAL HAZARDS: Strong Oxidizing Agent. Mix only with water. Use clean dry utensils. Do not add this product to any dispensing device containing remnants of any other product. Such use may cause a violent reaction leading to lire or explosion. Contamination with moisture, organic matter, or other chemicals may start a chemical reaction, with generation of heat, liberation of hazardous gases, and possible generation of fire and explosion. In case of contamination or decomposition, do not reseal container. If possible isolate container in open air or well ventilated area. Flood with large volumes of water, if necessary.

PRACTICAL TREATMENT: (First Aid) If swallowed feed bread soaked in milk followed by olive oil or cooking oil. Call a physician immediately.

If on skin. Brush off excess chemical and flush skin with cold water for at least 15 minutes. If irritation persists, get medical attention. If in Eyes. Flush with cold water for at least 15 minutes. Get medical attention

**STORAGE AND DISPOSAL:** Keep product dry in tightly closed container when not in use. Store in a cool dry, well ventilated area away from heat or open flame. In case of decomposition isolate container if possible, and flood with large amounts of water to dissolve all material before discarding. Place in trash collection or dispose in approved landfill area, or bury in a safe place.

Product No 102 Net Weight 2 lbs 12 oz (1 24 kg) Contents: 6, 7.33 oz. Tablets

(1182)



BEST AVAILABLE COPY

M