


TABEX FLOAT



Jan 3, 1973
 9087-4
 Comment
 # 9087-4
 FEDERAL REGISTER
 TRADE ACT
 POLYMER REGISTRATION NO.

MORE FUN / LESS WORK FOR POOL OWNERS

DIRECTIONS :

1. Adjust pool water pH to 7.2-7.8 range as determined by suitable test kit.
2. Stabilize pool water with 30 ppm conditioning agent (100% cyanuric acid).
3. Before first use of the feeder canister, superchlorinate (shock treat) pool water to 4 ppm chlorine residual, as determined by use of a chlorine test kit. Avoid entering pool until the chlorine residual drops to 2.0 ppm.
4. Before using the feeder canister, remove bottom cap (to permit water circulation) and remove tab over vent hole in top lid (to avoid air lock).
5. Maintain a chlorine residual of 1.0 to 1.5 ppm at all times (as determined by daily or more frequent use of a chlorine test kit) by immersing one Tabex chlorine tablet (or more until residual is obtained) in the Tabex Aquarian Feeder for up to 10,000 gallons of pool water per tablet. By the positioning of the canister in the feeder and/or adjustment of the water level in the feeder chamber, up to three tablets in a canister may be immersed. Two canisters may be inserted into the feeder; thus up to 6 tablets may be immersed at one time. This canister of tablets is designed for use in the Tabex Aquarian Feeder only. Tablets in canister should not be emptied directly in the pool as discoloration of the pool liner or paint could occur.
6. During the summer and peak bather loads, superchlorinate (shock treat) at least twice monthly as directed in "3" above. Failure to superchlorinate could result in growth of pool water algae. For such a condition, use Tabex Algae Out as directed on container.
7. Replace canister when last tablet drops into feeder.

Aspon
 INDUSTRIES, INC.
 TULLY, NEW YORK 13159



DANGER: KEEP OUT OF REACH OF CHILDREN. See other precautions and antidote on side panel.

Active Ingredients
 Trichloro-s-triazinetriene 95%
 Inert Ingredient 5%
 (Available Chlorine 85.5%)

CONVENIENT, ECONOMICAL POOL CHLORINATION

This canister containing 6 patented (U.S. Patent 3,325,411) slow dissolving, completely soluble Tabex Chlorine Tablets has been expressly designed for use with the Tabex Aquarian Feeder to provide a continuous, economical chlorine supply for control of bacteria and algae when used as directed.

DANGER: May cause eye damage. May be fatal or harmful if swallowed. May cause burns on wet skin. Rinse hands after handling. Strong oxidizer: Do not mix with any other chemical—mix only with water. Do not get into eyes. If in eyes, rinse with cold water, call physician. Avoid inhalation of dust and vapors. Keep away from organic matter (may cause fire), heat or open flame. Keep dry—chlorine and equally toxic gases liberated when wet. Do not reuse container; rinse thoroughly with water and discard.

Antidote: Drink milk of magnesia, vegetable oil or whites of eggs and call physician.

Treated pool water should not be discharged directly into lakes, streams or ponds.

Contents: 6 7.33-oz. tablets
 Net Weight: 2 lbs., 12 oz.
 EPA Reg. No. 9087-4
 Product No. 180



ACCEPTED WITH COMMENTS

MORE FUN / LESS WORK FOR POOL OWNERS

DIRECTIONS:

1. Adjust pool water pH to 7.2-7.8 range as determined by suitable test kit.
2. Stabilize pool water with 30 ppm conditioning agent (100% cyanuric acid).
3. Before first use of the float canister, superchlorinate (shock treat) pool water to 4 ppm chlorine residual as determined by use of a chlorine test kit. Avoid entering pool until the chlorine residual drops to 2.0 ppm.
4. Before using the canister, remove and discard plastic covering over holes around bottom of canister to admit water for tablet dissolution and remove tab over vent hole in top lid to avoid air lock.
5. Maintain a chlorine residual of 1.0 to 1.5 ppm at all times (as determined by daily or more frequent use of a chlorine test kit) by immersing 1 Tabex Chlorine Tablet (or more until residual is obtained) for up to 10,000 gallons of pool water per tablet. Immerse tablets by inserting canister through the Tabex Float and placing the unit in pool. To immerse 3 tablets, insert canister through wide mouth opening in the Tabex Float so that canister shoulders rest across narrow sides of opening. For 4 tablets, turn and depress canister shoulders into opening. If more than 4 tablets are required to obtain the above residual, an additional float canister unit will be needed. As bottom tablets dissolve, remaining tablets will drop down in canister. Do not use this canister in the Tabex Aquarion Feeder. Do not empty tablets directly into the pool as discoloration of the pool liner or paint could occur.
6. During the summer and peak bather loads, superchlorinate (shock treat) at least twice monthly as directed in 3 above. Failure to superchlorinate could result in growth of pool water algae. For such a condition, use Tabex Algae Out as directed on the container.
7. Replace canister when empty.

Apcon



Float Canister

DANGER: KEEP OUT OF REACH OF CHILDREN. See other precautions and antidote on side panel.

Active Ingredient: Trichloro-s-triazinetone 95%
 Inert Ingredient 5%
 (Available Chlorine 85.5%)

CONVENIENT, ECONOMICAL POOL CHLORINATION

This canister containing 6 patented U.S. Patent 2,854,111, slow dissolving, completely soluble Tabex Chlorine Tablets has been expressly designed for use with the Tabex Float to provide a continuous economical chlorine supply for control of bacteria and algae when used as directed.

DANGER: May cause eye damage. May be fatal or harmful if swallowed. May cause burns on wet skin. Rinse hands after handling. Strong oxidizer. Do not mix with any other chemical—mix only with water. Do not get into eyes. If in eyes, rinse with cold water, call physician. Avoid inhalation of dust and vapors. Keep away from organic matter (may cause fire), heat or open flame. Keep dry—chlorine and equally toxic gases liberated when wet. Do not reuse container. Destroy when empty.

Antidote: Drink milk of magnesia, vegetable oil or whites of eggs and call physician. Treated pool water should not be discharged directly into lakes, streams or ponds.

ACCEPTED

Net Wt. 2.12 lbs. (964 g)
 Net Vol. 1.00 gal. (3.78 l)
 EPA Reg. No. 908-3
 Product No. 181

ED UNDER N. 9087-3

DIRECTIONS:

1. Adjust pool water pH to 7.2-7.8 range as determined by suitable test kit
2. Stabilize pool water with 30 ppm conditioning agent (100% cyanuric acid).
3. Before first use of the float canister, superchlorinate (shock treat) pool water to 4 ppm chlorine residual as determined by use of a chlorine test kit. Avoid entering pool until the chlorine residual drops to 2.0 ppm.
4. Before using the canister, remove and discard plastic covering over holes around bottom of canister to admit water for tablet dissolution and remove tab over vent hole in top lid to avoid air lock.
5. Maintain a chlorine residual of 1.0 to 1.5 ppm at all times (as determined by daily or more frequent use of a chlorine test kit) by immersing 1 Tabex Chlorine Tablet (or more until residual is obtained) for up to 10,000 gallons of pool water per tablet. Immerse tablets by inserting canister through the Tabex Float and placing the unit in pool. To immerse 3 tablets, insert canister through wide mouth opening in the Tabex Float so that canister shoulders rest across narrow sides of opening. For 4 tablets, turn and depress canister shoulders into opening. If more than 4 tablets are required to obtain the above residual, an additional float canister unit will be needed. As bottom tablets dissolve, remaining tablets will drop down in canister. Do not use this canister in the Tabex Aquarian Feeder. Do not empty tablets directly into the pool as discoloration of the pool liner or paint could occur.
6. During the summer and peak bather loads, superchlorinate (shock treat) at least twice monthly as directed in 3 above. Failure to superchlorinate could result in growth of pool water algae. For such a condition, use Tabex Algae Out as directed on the container.
7. Replace canister when empty.

Aspon

MORE FUN / LESS WORK



Tabex
Chlorine Tablets

Float Canister

DANGER: KEEP OUT OF REACH OF CHILDREN. See other precautions and antidote on side panel.

Active Ingredient
Trichloro-s-triazinetrione 95%
Inert Ingredient 5%
(Available Chlorine 85.5%)

CONVENIENT ECONOMICAL POOL CHLORINATION

This canister containing 6 patented (Patent 3,475,411), slow dissolving, completely soluble Tabex Chlorine Tablets has been expressly designed for use with the Tabex Float to provide a continuous economical chlorine supply for control of bacteria and algae when used as directed.

DANGER: May cause eye damage. May be fatal or harmful if swallowed. May cause burns on wet skin. Rinse hands after handling. Strong oxidizer: Do not mix with any other chemical—mix only with water. Do not get into eyes. If in eyes, rinse with cold water, call physician. Avoid inhalation of dust and vapors. Keep away from organic matter (may cause fire), heat or open flame. Keep dry—chlorine and equally toxic gases liberated when wet. Do not reuse container. Destroy when empty.

Antidote: Drink milk of magnesia, vegetable oil or whites of eggs and call physician.

Treated pool water should not be discharged directly into lakes, streams or ponds.

ACCEPTED

1-8-73

Net Weight 2.25 lbs. 12 oz.
EPA Reg. No. 9087-3
Product No. 181

FOR LISTING UNDER NO. 9087-3

BEST AVAILABLE COPY



REPT. APPROVED 1973

ACCEPTED
JAN 8 1973
UNDE... FICIDE
FUNG... THE CASE ACT
FOR ECON... REGISTERED
ED UNDER NO. 9087-3

21