

RECEIVED
 Jan 3 1973
 9087-4
 W. M. W. W.

DIRECTIONS:

1. Adjust pool water pH to 7.2-7.8 range as determined by suitable test kit.
2. Stabilize pool water with 30 ppm conditioning agent (100% cyanuric acid)
3. Before first use of the feeder canister, superchlorinate (shock treat) pool water to 4 ppm chlorine residual, as determined by use of a chlorine test kit. Avoid entering pool until the chlorine residual drops to 2.0 ppm
4. Before using the feeder canister, remove bottom cap (to permit water circulation) and remove tab over vent hole in top lid (to avoid air lock).
5. Maintain a chlorine residual of 1.0 to 1.5 ppm at all times (as determined by daily or more frequent use of a chlorine test kit) by inserting one Tabex chlorine tablet (or more until residual is obtained) in the Tabex Aquarian Feeder for up to 10,000 gallons of pool water per tablet. By the positioning of the canister in the feeder and/or adjustment of the water level in the feeder chamber, up to three tablets in a canister may be immersed. Two canisters may be inserted into the feeder, thus up to 6 tablets may be immersed at one time. This canister of tablets is designed for use in the Tabex Aquarian Feeder only. Tablets in canister should not be emptied directly in the pool as discoloration of the pool liner or paint could occur
6. During the summer and peak bather loads, superchlorinate (shock treat) at least twice monthly as directed in "3" above. Failure to superchlorinate could result in growth of pool water algae. For such a condition, use Tabex Algae Out as directed on container.
7. Replace canister when last tablet drops into feeder.

Aspon

BEST DOCUMENT AVAILABLE

MORE FUN / LESS WORK FOR POOL OWNERS



Tabex
 Chlorine Tablets

Feeder Canister

DANGER: KEEP OUT OF REACH OF CHILDREN. See other precautions and antidote on side panel.

Active Ingredients
 Trichloro-s-triazinetrione 95%
 Stabilizer 5%
 Available Chlorine 85.5%

CONVENIENT, ECONOMICAL POOL CHLORINATION

This canister containing 6 patented (U.S. Patent 3,325,411) slow dissolving, completely soluble Tabex Chlorine Tablets has been expressly designed for use with the Tabex Aquarian Feeder to provide a continuous, economical chlorine supply for control of bacteria and algae when used as directed.

DANGER: May cause eye damage. May be fatal or harmful if swallowed. May cause burns on wet skin. Rinse hands after handling. Strong oxidizer. Do not mix with any other chemical—mix only with water. Do not get into eyes. If in eyes, rinse with cold water, call physician. Avoid inhalation of dust and vapors. Keep away from organic matter (may cause fire), heat or open flame. Keep dry—chlorine and equally toxic gases liberated when wet. Do not reuse container; rinse thoroughly with water and discard.
Antidote: Drink milk of magnesia, vegetable oil or whites of eggs and call physician.

Treated pool water should not be discharged directly into lakes, streams or ponds.

Contents 6 7.33-oz tablets
 Net Weight 2 lbs 12 oz
 EPA Reg No 9087-4
 Product No 180