

ACCEPTED

9087-4

JUN 1 1983

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No.

DIRECTIONS FOR USE.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Tabex Slow Dissolving Chlorination Tablets are an extremely concentrated form of swimming pool sanitizing agent. Tabex Chlorination Tablets are designed to dissolve slowly and at a uniform rate thereby eliminating the nuisance of frequent hand feeding of granular or liquid chlorination. We suggest the use of the TABEX Aquatic Feeder for the ultimate convenience of continuous automatic drip chlorination.

FOR CONTINUOUS DAILY CHLORINATION

Before the first use of TABEX SLOWLY DISSOLVING CHLORINATION TABLETS be sure to take the following steps:

A. Adjust the pool water pH value to between 7.2 and 7.8 as determined by the TABEX Swimming Pool Test Kit. Further adjust your Total Alkalinity to a minimum of 100 ppm (parts per million).

B. Stabilize the water to at least 50 ppm with TABEX Conditioner as directed on container label.

C. Superchlorinate (SHOCK TREAT) your pool water to a 4 ppm residual chlorine level using TABEX Super Shock Tablets as directed on the container label.

D. Butyl stop use of this canister, remove bottom cap to permit water circulation.

E. Maintain a FREE AVAILABLE CHLORINE residual of at least 1.0 to 1.5 ppm by inserting one (1) TABEX tablet into the Aquatic Feeder for each 5,000 to 10,000 gallons (25,000 to 50,000 liters) of pool water. Immerse tablets by inserting one or two Feeder Tablets into the Aquatic Feeder (the actual number of tablets and duration required is based upon the length of time you pack pump operation).

F. If the water level is low, use your back cover remaining opening in Feeder to add water to the pool. Position and the leader float cover 13 to 15 inches from the pool deck. Tablets in the Feeder Canister will be immediately available to chlorinate the pool water.

G. After the daily test, you pool water with the TABEX Swimming Pool Test Kit and adjust the Feeder as described above to maintain a FREE AVAILABLE CHLORINE residual of 1.0 to 1.5 ppm.

H. Superchlorinate (SHOCK TREAT) your pool water at least once monthly (more often in Southern regions) during hot weather and peak bather loads. Allow 24 hours for the chlorine to stabilize. Adjust the Feeder as described above to maintain a FREE AVAILABLE CHLORINE residual of 1.0 to 1.5 ppm.

EPA Reg. No. 9087-4

EPA Est. No. 9087-NY-12

More Fun/Less Work For Pool Owners




Tabex

Chlorination Tablets Feeder Canister

contains no calcium

DANGER: KEEP OUT OF REACH OF CHILDREN
See other precautions and antidotes on side panel

Active Ingredients
TRICHLORO S TRIAZINETRIONE 95%
Inert Ingredients 5%
Available Chlorine 86.5%



1. Replace canister when last tablet drops into the feeder.
NEVER ADD THESE SLOW DISSOLVING CHLORINATION TABLETS DIRECTLY INTO THE SWIMMING POOL AS DISCOLORATION OF THE POOL LINER OR FINISH WILL OCCUR.

DANGER

Highly Corrosive: causes skin and eye damage. May be fatal if swallowed. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. Irritating to nose and throat. Avoid breathing dust. Remove & wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not discharge into lakes, streams, ponds or public waters unless in accordance with an NPDES Permit. For guidance contact the regional office of the EPA.

PHYSICAL AND CHEMICAL HAZARDS

Strong oxidizing agent. Mix only with water. Use clean dry utensils. Do not add this product to any dispensing device containing remnants of any other product. Such use may cause a violent reaction leading to fire or explosion. Contamination with moisture, organic matter, or other chemicals may start a chemical reaction with generation of heat, liberation of hazardous gases and possible generation of fire and explosion. In case of contamination or decomposition, do not reuse container. If possible, isolate container in open or well-ventilated area. Flood with large volumes of water, if necessary.

PRACTICAL TREATMENT

If swallowed: Do not induce vomiting. If on skin, brush off excess chemical and flush skin with cold water for at least 15 minutes. If irritation persists, get medical attention. If in eyes, flush with cold water for at least 15 minutes. GET MEDICAL ATTENTION.

STORAGE AND DISPOSAL

Keep container tightly closed when not in use. Store in a cool, well-ventilated area away from heat or open flame. In case of decomposition isolate container, if possible, and flood with large amounts of water to dissolve all material before discarding. Place in trash collection or dispose in an approved landfill area or bury in a safe place.

Product No. 180 (11182)
Net Weight: 2.75 lbs (1.26 Kg)
Contents: 6.7.33 ounce (210 gram) tablets

Aqua

11