

When **ALGAECIDE** is used as directed, it kills and prevents development of:

- a. *Chlorella pyrenoidosa* (green algae)
- b. *Phormidium inundatum* (black algae)
- c. *Phormidium retzii* (square D algae)
- d. *Lyngaba versicolor* (blue-green algae) and other swimming pool algae.

NOTE: Kill, Control and/or Destroy refer to initial dose. Prevent development, suppress growth, inhibit, refer to maintenance dose.

ALGAECIDE is compatible with most commonly used swimming pool chemicals but should not be mixed with soap or other cleaners or pre-mixed with other water-treating chemicals. ALGAECIDE works best when added by itself directly to pool water. ALGAECIDE at use dilution is NON-CORROSIVE to inanimate surfaces.

ESL ALGAECIDE CONCENTRATED SWIMMING POOL ALGAECIDE

Keeps Pool Water Clear and Free of Visible Algae, Controls Algae Slimes

Active Ingredients:

n - Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈) dimethyl benzyl ammonium chloride 8.4%

n - Di-alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈) Methyl benzyl ammonium chloride 1.6%

Inert Ingredients: water 90.0%
Total 100.0%

Net Contents one (1) quart
EPA Reg. No. 9096-7

DANGER
Keep Out of Reach of Children.
See other precautionary & First Aid statements on back panel.

eastern shore  laboratories incorporated
P. O. Box 657 Laurel, Delaware 19956

DIRECTIONS FOR USE:

Initial Dose: One (1) gallon ALGAECIDE to 50,000 gallons of water.

Maintenance Dose: One (1) quart ALGAECIDE to 50,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance.

Maintenance dose should be added at 3 - 5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If make up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of ALGAECIDE for each 10,000 gallons of fresh make up water added per week.

When swimming season is over, add 1 gallon of ALGAECIDE to 50,000 gallons of water in treated water left in pool. This does help provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

~~PRECAUTIONARY STATEMENTS FOR PRODUCTS PRODUCING LOCAL IRRITATION~~

Danger:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

First Aid:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Apply this product only as specified on this label.

Note To Physician:

Probable mucosal damage may contraindicate the use of gastric lavage. Measure against circulatory shock, respiratory depression and convulsion may be needed.

ACCEPTED

NOV 30 1977
UNDER THE
FUNGICIDE
FOR ECONOMIC PEST CONTROL
ED UNDER NO. 9096-7
TO ATTACHED COMMENTS.