

WAYNE™

RAT & MOUSE KILLER


Warfarin Rodenticide
Contains Prolin*

Ready-to-Use Bait in "Pelletized" Form

ACTIVE INGREDIENTS
NET WEIGHT

EPA Reg. No. 9114-1

CAUTION:



**WAYNE
ANIMAL
HEALTH
AIDS**



DIRECTIONS FOR USE

RATS Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

MICE Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait Stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

CAUTION Place baits in locations not accessible to children, pets, domestic animals and wildlife. All outdoor settings should be made in tamper-proof boxes. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce Vitamin K, an antidote for warfarin.

Do not reuse container. Destroy when empty.

THIS CONTAINER CAN PROVIDE AN EXCELLENT
BAIT STATION CUT OPENINGS

