



WAYNE™

Dairy and Livestock Spray Concentrate

RESIDUAL DAIRY AND LIVESTOCK SPRAY
(Containing Ciodrin)

Controls: Face Flies, Stable Flies, Horn Flies, Ticks, Lice on Dairy and Beef Cattle,
Sheep, Goats and Swine

Controls: House Flies in Dairy Barns and other Farm Buildings



ACTIVE INGREDIENTS EPA Reg. No. 9214-2
Dimethyl Phosphate of alpha-methylbenzyl 14.40%
3-Hydroxy-cis-crotonate 75.60%
Petroleum Distillates
INERT INGREDIENTS: 10.00%

WAYNE
ANIMAL
HEALTH
AIDS



WARNING: Keep out of reach of children. See other warnings on back panel.

Manufactured for Animal Health Chemicals, Inc., St. Louis, Mo.

DIRECTIONS FOR USE

DAIRY CATTLE: To control Face Flies, Stable Flies, Horn Flies, Ticks and Lice. Dilute 1 pint of Wayne Dairy and Livestock Spray Concentrate in 3½ gallons of water (1 to 28). Apply one to two pints of dilute spray per adult animal and proportionately less for younger animals. Repeat as required but not more often than once every seven days.

LOOSE PENNED AND FEED LOT CATTLE: To control Face Flies, Stable Flies, Horn Flies and Lice. Dilute 1 pint of Wayne Dairy and Livestock Spray Concentrate in 3½ gallons of water (1 to 28). Spray at the rate of 1 to 2 pints of diluted spray over and on adult animals and proportionately less for younger animals. Repeat as required but not more often than once every seven days. Do not apply to calves under 6 months of age.

SHEEP, GOATS & SWINE: To control Lice on Swine, Lice and Ticks on Sheep and Goats. Dilute 1 pint in 6 gallons of water (1 to 50) and spray at the rate of one gallon of diluted spray per animal. Repeat as required but not more often than once every seven days.

BACK RUBBERS: For control of Flies on Beef and Dairy Cattle. Dilute 1 pint of Wayne Dairy and Livestock Spray Concentrate with 2½ gallons of suitable oil. Charge back rubber as needed. Do not spray animals with this oil solution.

RESIDUAL SURFACE SPRAY FOR CONTROL OF HOUSE FLY AND RESTING STABLE FLY. Dilute 1 pint of Wayne Dairy and Livestock Spray Concentrate in 1½ gallons of water (1 to 14). Spray at the rate of 1 gallon of diluted spray per 1000 square feet of surface (wet surfaces). Do not to point of excessive run off. Spray walls of dairy barns, on fences, hog pens and other similar places where flies are seen to congregate. Repeat as required or at least twice monthly.

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Avoid contact with skin, eyes or clothing. Do not breathe spray mist. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or

SPRAY DILUTION RATES

For (1 to 28)		For (1 to 48)	
Wayne Dairy and Livestock Spray Concentrate	Water	Wayne Dairy and Livestock Spray Concentrate	Water
8 Fluid Ounces (1 measuring cup)	1½ Gallons	5 Tablespoons (1 measuring cup)	1 Gallon
1 Pint	3½ Gallons	8 Fluid Ounces (1 measuring cup)	3 Gallons
1 Quart	7 Gallons	1 Pint	6 Gallons
1 Gallon	28 Gallons	1 Quart	12 Gallons
		1 Gallon	48 Gallons

respirator of a type found suitable by the U. S. Bureau of Mines for protection against Ciodrin insecticide. Apply this product only as specified on this label and at recommended dosages. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Keep away from heat and open flame. Keep out of reach of children.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate feed, food stuffs or water. Food utensils such as teaspoons and table spoons should not be used for food purposes after use with product. Do not reuse empty container. Destroy it by perforating or crushing. Bury or incinerate in safe place.

