

WAYNE DISINFECTANT

For cleaning, disinfecting and sanitizing animal housing and equipment; cleaning sows prior to farrowing; sanitizing poultry drinking water; as a shoe bath for barns and poultry houses; as a deodorant; sanitizing food processing equipment.

ACTIVE INGREDIENTS:

a-(p-Nonylphenyl)-omega-hydroxypoly(oxyethylene)-iodine complex\* 18.05%  
(providing 1.75% titratable iodine)

Phosphoric acid 16.00%

INERT INGREDIENTS Total 65.95%  
100.00%


\* West Chemical's Famed Iodine<sup>®</sup>  
Manufactured under the following U. S. Patents: 2,931,777  
2,977,315  
3,028,299

CAUTION: Keep out of the reach of children.  
See additional warnings on rear panel.

Manufactured for

ALLIED MILLS, INC.,  
Chicago, Ill.

Net Contents: One U.S. Gallon.



REAR PANEL

Wayne Disinfectant has a built-in indicator of germicidal activity. When the color disappears, a fresh solution should be prepared.

DIRECTIONS FOR USE:

1. For cleaning and disinfecting animal housing and equipment on the farm. Clean and disinfect in the early morning with Wayne Disinfectant. Use 1/2 ounce per gallon of water. Use a soft-bristle brush to clean and disinfect with a good brush or with a water sprayer. Start at the highest point of the building or equipment and work down. Let solution drain dry without rinsing.

2. For cleaning sows.

Before placing sows in the farrowing box, scrub sow's flanks and udder with a lukewarm solution of Wayne Disinfectant per gallon of water. Use a soft-bristle brush. Rinse with clean water, and do not rinse until just before farrowing.

3. Sanitizing poultry drinking water.

Add 1 ounce of Wayne Disinfectant per gallon of drinking water. Use regular use on the farm. When used in the waterers, the waterers first