



WAYNE®

Fly Block MEDICATED

(FOR RUMINANTS ONLY)

9114-18
ACCEPTED
NOV 20 1972
UNDER THE FEDERAL MICROFILME
FUNCTIONS OF THE NATIONAL ARCHIVES
FOR SUBJECTS
TO ATTACHED COMMENTS.

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle. Aids in the control of stomach worms (*Haemonchus*, *Ostertagia* and *Trichostrongylus* spp.), nodular worms (*Fasciola hepatica* spp.), large-mouth bowelworms (*Chabertia* spp.) and pinworms (*Oxyuris equi* spp.) in cattle. Aids in the prevention of lungworms (*Dictyocaulus viverrinx*), *Parapheraphus necrophorus*, and in soft tissue lesions caused by *Clostridium botulinum* and *Clostridium perfringens* in cattle. A nutritional supplement.

Pesticidally Active Drug Ingredient
Phenothiazine (Biologically assayed) 1.5%
Pesticidally Inert Ingredients* 98.5%
Other Active Drug Ingredient
Ethylene Diamine Dihydrochloride 1.0%
*Refers to feed and minerals which are neither active nor inert.

Guaranteed Analysis

Crude Protein (This includes not more than 6.75% equivalent protein from non protein nitrogen) 16.0%
Crude Fat 12.0%
Crude Fiber 12.0%
Calcium (Ca) 0.50%
Phosphorus (P) 0.50%
Iodine 0.01%
Salt 0.50%
Vitamin A 100,000 I.U./lb.
Vitamin D₃ 100,000 I.U./lb.

Ingredients

Ground yellow corn, cottonseed meal, brewers dried grains, wheat middlings, dehydrated alfalfa meal, urea, cane molasses, vitamin A palmitate, irradiated dried yeast (source of vitamin D₃), ethylene diamine dihydrochloride, ground lime, dicalcium phosphate, calcium sulfate, salt, copper oxide, cobalt carbonate, manganese oxide, zinc oxide, iron oxide, magnesium oxide, and zinc oxide.

with ethoxyquin) urea, cane molasses, vitamin A palmitate, irradiated dried yeast (source of vitamin D₃), ethylene diamine dihydrochloride, ground lime, dicalcium phosphate, calcium sulfate, salt, copper oxide, cobalt carbonate, manganese oxide, zinc oxide, iron oxide, magnesium oxide, and zinc oxide.

CAUTION: Keep Out of Reach of Children.

Feeding Directions

For treatment of horn flies and face flies, feed 0.25 pounds of block per head daily. For control of stomach worms, nodular worms, large-mouth bowelworms, pinworms, lungworms, and *Parapheraphus necrophorus*, feed 0.25 pounds of block per head daily. For control of *Clostridium botulinum* and *Clostridium perfringens*, feed 0.25 pounds of block per head daily. For control of *Dictyocaulus viverrinx*, *Parapheraphus necrophorus*, and *Oxyuris equi*, feed 0.25 pounds of block per head daily. For control of *Chabertia* spp., feed 0.25 pounds of block per head daily. For control of *Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp., feed 0.25 pounds of block per head daily. For control of *Fasciola hepatica*, feed 0.25 pounds of block per head daily. For control of *Dictyocaulus viverrinx*, *Parapheraphus necrophorus*, and *Oxyuris equi*, feed 0.25 pounds of block per head daily. For control of *Chabertia* spp., feed 0.25 pounds of block per head daily. For control of *Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp., feed 0.25 pounds of block per head daily. For control of *Fasciola hepatica*, feed 0.25 pounds of block per head daily.

WARNING: Do not administer to horses. Do not administer to dogs. Do not administer to cats. Do not administer to birds. Do not administer to swine. Do not administer to fish. Do not administer to reptiles. Do not administer to amphibians. Do not administer to invertebrates. Do not administer to plants. Do not administer to fungi. Do not administer to bacteria. Do not administer to viruses. Do not administer to parasites. Do not administer to other organisms.

FPA Reg. No. 9114-18

Allied Mills, Inc. - General Offices - Chicago, Ill. 60606

NET WEIGHT 33.33 LBS.

8129
2013

