



WAYNE[®] ORAL LARVICIDE [CONTAINS RABON[®] ORAL LARVICIDE]

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated Cattle and House Flies in the manure of treated Swine

ACTIVE INGREDIENT	BY WEIGHT
100% Rabon Oral Larvicide	100%
INERT INGREDIENTS	Total
	100.00%

← **KEEP OUT OF REACH OF CHILDREN
CAUTION** →

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When fed to cattle RABON[®] Oral Larvicide passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, RABON[®] Oral Larvicide helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

Start feeding RABON[®] larvicide feed early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall, until cold weather restricts fly activity.

Rations containing RABON[®] Oral Larvicide may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

RABON[®] Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and barns to control adult house flies and stable flies which can breed not only in cattle manure but other decaying vegetable matter or sludge on the premises.

In order to achieve optimum fly control, RABON[®] Oral Larvicide should be used in conjunction with other good management and sanitation practices.

NET WT. 45 LBS.
(20.45 kg)

H233-3N

Wayne Feed Division
Continental Grain Co.
Chicago, IL 60606

(5532-R2)
822
EPA Reg. No. 101-107-01
EPA Reg. No. 101-107-01

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 29 1982

Wayne Feed Division, Inc.
Pesticide Registration Act
Responsible
EPA Reg. No.

9119-26

142

WAYNE ORAL LARVICIDE

DIRECTIONS FOR USE (Continued)

Wayne Oral Larvicide is an effective fly and tick repellent when used in accordance with the directions. This product is not intended for use on birds, fish, or aquatic animals. It is not intended for use on horses, dogs, cats, or other domestic animals. It is not intended for use on ornamental plants, lawns, or gardens. It is not intended for use on any other animals or plants. It is not intended for use on any other animals or plants. It is not intended for use on any other animals or plants.

For more information on the use of Wayne Oral Larvicide, contact your local Wayne Feed Division office or the manufacturer.

CATTLE: IN A COMPLETE RATION - No Other Ration Fed

Wayne Oral Larvicide can be used to protect cattle from fly and tick infestation when fed in a complete ration.

To prepare a larvicide ration, mix 1.3 lbs. of Wayne Oral Larvicide per ton of meal type feed and offer free choice.

For a rate to sustain milk production, mix 2.6 lbs. of Wayne Oral Larvicide per ton of meal type feed and offer 4-6 lbs. of feed per animal per day. This is equivalent to 45.4 mg. of RABON[®] per lb. of feed.

CATTLE: IN A CONCENTRATED FEED - Roughage Fed Separately

Wayne Oral Larvicide can be used to protect cattle from fly and tick infestation when fed in a concentrated feed.

To prepare a larvicide concentrate feed, mix 1.3 lbs. of Wayne Oral Larvicide per ton of concentrate feed and offer 4-6 lbs. of feed per animal per day. This is equivalent to 45.4 mg. of RABON[®] per lb. of feed.

MIXING DIRECTIONS

Pounds of Concentrate Consumed per Animal per day	RABON [®] In the Concentrate		Pounds of Wayne Oral Larvicide per Ton of Concentrate
	mg	lb	
10	454	0.18	4.0
15	681	0.27	6.0
20	908	0.36	8.0
50	2270	0.90	20.0
100	4540	1.80	40.0
150	6810	2.70	60.0
200	9080	3.60	80.0
250	11350	4.50	100.0
300	13620	5.40	120.0

Feed the appropriate larvicide concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200-1700 pounds, increase the amount of concentrate fed to 1.5 times that recommended.

SWINE: All Swine should be treated

Pigs (weaners to market weight): Mix 1.3 lbs. of this product per ton of meal type feed and offer free choice. This is equivalent to 22.7 mg. of RABON[®] per lb. of feed.

Sows, Boars and Breeding Gilt: Mix 2.6 lbs. of this product per ton of meal type feed and offer 4-6 lbs. of feed per animal per day. This is equivalent to 45.4 mg. of RABON[®] per lb. of feed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to federal, state or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved federal, state or local procedures under the Resource Conservation and Recovery Act.

RABON[®] is a registered trademark of Shell Oil Company.



NET WT. 45 LBS.
(20.41 kg)

Wayne Feed Division
Centennial Feeds Co.
Chicago, IL 60606

H233-3N

(8502-R2)
826
EAST...
CHICAGO, ILL.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

NOV 29 1982

OFFICE OF PESTICIDE REGISTRATION

Continental Grain Co.
10 South Riverside Plaza
Chicago, IL 60606

310/759-3
18/2

Attn: Mr. ML A. Werner

Dear Mr. Werner:

Subject: Amendment - Add Swine
Wayne Oral Larvicide
EPA Reg. No: 9114-26
Appl. Date: June 23, 1982

The amendment referred to above, submitted in connection with registration under FIFRA Sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA Sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before your release the product for shipment bearing the amended labeling:

Change the container disposal statement in accordance with the enclosure (page 2).

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

G