

Reg. 9123-18

Baldwin fertilizer Baldwin

LAWN FOOD & WEED CONTROL

13-2-3

ACCEPTED
MAY 30 1985
9123-18

A quick greening fertilizer
for a luxurious lawn

.... combined with
herbicide to control
most common broadleaf weeds.

LAWN FOOD
&
WEED CONTROL

Covers 5000 sq. ft. 26 lbs. (11.8 kg) net wt.	Active Ingredients *Dimethylamine Salt of 2,4-Dichlorophenoxy Acetic Acid Inert Ingredients *Equivalent to 0.84% 2,4-Dichlorophenoxy Acetic Acid	1.01% 98.99%
CAUTION: KEEP OUT OF REACH OF CHILDREN See back panel for additional precautionary statements		EPA EST. NO. 9123 WA 1 EPA REG. NO. 9123

Magnolia Products, Inc.
Kirkland, WA 98033

Balance

LAWN FOOD & WEED CONTROL

A quick greening fertilizer combined with herbicide to control most common broadleaf weeds.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply in spring or summer when broadleaf weeds are actively growing and temperatures are above 60°F.

1. For best results, before applying, moisten lawn very lightly (mist only) with hose, so weed killer particles will adhere to the moist weed leaves.
2. Apply fertilizer evenly, selecting spreader setting below.
3. Avoid overlapping when applying.
4. Do not water for 24 hours after application, to avoid washing weed killer particles off the broad leaf weeds. Then, water thoroughly.

NOTE: If temperature remains over 70°F, water thoroughly 12 hours after application.

Do not apply on newly seeded lawns.

Weed control does not occur overnight. Broadleaf weeds will begin to wilt in a few days, and will disintegrate in about 3 to 4 weeks.

Weed seeds presently in the lawn and those deposited by the wind, will not be controlled by this application.

Common broadleaf weeds controlled: chickweed, dandelion, docks, mustard, sorrel, plantain, dog fennel, clover, Japanese clover.

SPREADER SETTING

Scotts Spreader

Setting #6

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling. Vapors from this product may injure susceptible plants in immediate vicinity. Do not use on St. Augustine grass, centipede grass, carpet grass or dicanders, nor on lawns where desirable bent grass or clover is present. Do not apply on or near desirable flowers, shrubs, fruits or vegetable beds. Do not use on newly seeded lawns. Do not apply within drip-line of desirable trees and shrubs. Avoid drift of dust to desirable plants.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water used for irrigation or domestic purposes. NOTE: Birds feeding over treated areas may be harmed.

STORAGE AND DISPOSAL: Do not store near and avoid contamination of food, seeds, insects, cades, fungicides and other fertilizers. Wash out fertilizer spreader after use with this fertilizer. Do not reuse empty container. Wrap container and put in trash collection.

ANALYSIS

Total Nitrogen (N)	13%
Available Phosphoric Acid (P ₂ O ₅)	2%
Soluble Potash (K ₂ O)	3%

INGREDIENTS

Urea, Dinitrogen, Muriate of Potash, Ammonium Phosphate, Organic Soil Conditioner, Herbicide

For an attractive, healthy lawn year round, apply the proper fertilizer every 4 weeks during spring, summer and fall (approximately 3-4 times per year).

Your home deserves a beautiful lawn
USE A BALANCED LAWN CARE PLAN

SPRING	SUMMER	SUMMER	FALL
MOSS CONTROL & LAWN FOOD	LAWN FOOD	LAWN FOOD & WEED CONTROL	MOSS CONTROL & LAWN FOOD

BEST AVAILABLE COPY

Dalman fertilizer

Balan

BEST AVAILABLE COPY

LAWN FOOD & WEED CONTROL

13-2-3

ACCEPTED
MAY 30 1985
9123 18

A quick greening fertilizer for a luxurious lawn

.... combined with herbicide to control most common broadleaf weeds.

LAWN FOOD & WEED CONTROL

Covers
5000 sq. ft.
26 lbs
(11.8 kg)
net wt

Active Ingredients
*Dimethylamine Salt of 2,4-Dichlorophenoxy Acetic Acid
Inert Ingredients
*Equivalent to 0.84% 2,4-Dichlorophenoxy Acetic Acid

1.01%
98.99%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional precautionary statements

EPA EST. NO. 9123 WA 1
EPA REG. NO. 9123

Balance

LAWN FOOD & WEED CONTROL

A quick greening fertilizer combined with herbicide to control most common broadleaf weeds.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply in spring or summer when broadleaf weeds are actively growing and temperatures are above 60°F.

1. For best results, before applying, moisten lawn very lightly (mist only) with hose, so weed killer particles will adhere to the moist weed leaves.
2. Apply fertilizer evenly, selecting spreader setting below.
3. Avoid overlapping when applying.
4. Do not water for 24 hours after application, to avoid washing weed killer particles off the broad leaf weeds. Then, water thoroughly.

NOTE: If temperature remains over 70°F, water thoroughly 12 hours after application.

Do not apply on newly seeded lawns.

Weed control does not occur overnight. Broadleaf weeds will begin to wilt in a few days, and will disintegrate in about 3 to 4 weeks.

Weed seeds presently in the lawn and those deposited by the wind, will not be controlled by this application.

Common broadleaf weeds controlled: chickweed, dandelion, docks, mustard, sorrel, plantain, dog fennel, clover, Japanese clover.

SPREADER SETTING

Scotts Spreader

Setting #6

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling. Vapors from this product may injure susceptible plants in immediate vicinity. Do not use on St. Augustine grass, centipede grass, carpet grass or dicentra, nor on lawns where desirable bent grass or clover is present. Do not apply on or near desirable flowers, shrubs, fruits or vegetable plants. Do not use on newly seeded lawns. Do not apply within dripline of desirable trees and shrubs. Avoid drift of dust to desirable plants.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water used for irrigation or domestic purposes. NOTE: Birds feeding over treated areas may be harmed.

STORAGE AND DISPOSAL: Do not store near and avoid contamination of food, seeds, insecticides, fungicides and other fertilizers. Wash out fertilizer spreader after use with this label. Do not reuse empty container. Wrap container and put in trash collection.

ANALYSIS

Total Nitrogen (N)	13%
Available Phosphoric Acid (P ₂ O ₅)	2%
Soluble Potash (K ₂ O)	3%

INGREDIENTS

Urea, Dolomite, Murate of Potash, Ammonium Phosphate, Organic Soil Conditioner, Herbicide

For an attractive, healthy lawn year round, apply the proper fertilizer every 4 weeks during spring, summer and fall (approximately 3-4 times per year).

Your home deserves a beautiful lawn
USE A BALANCED LAWN CARE PLAN

SPRING	SUMMER	SUMMER	FALL
MOSS CONTROL & LAWN FOOD	LAWN FOOD	LAWN FOOD & WEED CONTROL	MOSS CONTROL & LAWN FOOD

BEST AVAILABLE COPY