

[9129-3]

DIRECTIONS FOR USE

Southland NR-4 Insecticide will effectively control flies, mosquitoes, wasps, flying moths, roaches, ants, crickets, silverfish and exposed stages of stored products insects, when used as directed.

For a fog: use Southland NR-4 Insecticide undiluted.

For space and contact spray: use Southland NR-4 Insecticide diluted with an equal part of deodorized kerosene prior to use.

As a fog for space spray:

For Flies, Mosquitoes, Wasps, and Gnats: Close all doors and windows. Point nozzle and direct the fog or mist to all parts of the room, especially windows and other insect sources which attract these insects. For best results, make sure fog or mist is evenly distributed throughout control area. Apply 5 to 10 seconds fogging time per 1,000 cubic feet of space. In case of space sprays, spray until control area is filled with mist. Keep room closed for one hour after application. Repeat treatment when necessary.

For Roaches, Ants, Silverfish and Crickets: Close all inside doors, cabinets, closets, etc. Direct the fog or spray toward insects, into cracks, crevices, and trails, on doorsills and window frames. For initial application, when fogging apply for 10 to 15 seconds for each 100 square feet of floor space. Follow up with periodic space applications at 5 to 10 seconds per 1,000 cubic feet. Keep room closed for one hour after application. Repeat treatment when necessary.

Southland NR-4

(A non-residual insecticide for fog or spray application)

Active Ingredients:

Pyrethrins	0.350%w/w
n-Octyl stearoxide of L-cinchona	1.540%
Related compounds	0.210%
Methylated cephalinics	2.525%
Petroleum distillate	95.275%
	100.000%

For use in dairies, delis, restaurants, or wherever food is processed or sold.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

When using this product as a space spray or fog in these areas, apply only when the plant is not in operation. Avoid using this product in edible products areas of meat packing plants.

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

For Stored Products Insects (Exposed Stages): For such insects as flour beetles and granary weevils, spray or fog insects, into cracks, crevices, around machinery, around containers of stored products and other areas wherever these insects may hide. It is important to contact as many insects as possible. When fogging a minimum of 30 seconds treatment time should be used per 100 square feet of floor space. Treated area should be kept tightly closed for at least 45 minutes. Repeat treatment when necessary.

For Such Insects As Flour Moths and Indian Meal Moths: Close all doors and windows. Direct fog or spray to all parts of the room, especially into dark areas, around stored products containers, where adult moths may be resting. When fogging a minimum treatment time of 8 to 10 seconds should be used for each 1,000 cubic feet of space. Keep treated area closed for at least 30 minutes. Repeat weekly or when flying moths reappear.

As a Contact Spray:

For Roaches, Ants, Silverfish, Crickets and Stored Products Insects (Exposed Stages): Treat cracks and crevices in woodwork, dark places behind pipes, and all places which these insects inhabit, hitting as many insects as possible. Repeat treatment whenever reinfestation occurs.

USDA Reg. No.

ACCEPTED

JUN 14 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 9129-3 SUBJECT