


[9143-52]



DIRECTIONS FOR USE
IN FOGGING EQUIPMENT AND HAND SPRAYERS TO KILL ACCESSIBLE STAGES OF INSECTS* INFESTING DRIED FRUIT PRODUCTS IN STORAGE OR BEING PROCESSED: Close room or warehouse and shut off all ventilating systems. Dispense this product as a fine mist in the air above trays or shelves of dried fruit. Use a spraying device similar to the Microsol machine, or use fixed ceiling sprayers that produce a very fine spray. Do not apply to fruit directly. If overhead sprayers are used, be sure that the spray is discharged from a position of at least 6 feet over the top trays of fruit. Use a dosage of 1 gallon of this spray per 50,000 cubic feet of room space. Repeat applications at this dosage in any given month. More than ten applications may also be made if this spray is diluted with more petroleum distillate, so that the pyrethrins and synergists in the final spray are reduced proportionately. Do not remain in treated areas. Ventilate the area after treatment is completed. Maximum dilution is 1 part F-6630 plus 5.6 parts oil.

Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-Toothed Grain Beetle

TO KILL THE ADULT AND ACCESSIBLE FORMS OF INSECT PESTS* OF STORED FOOD (STORED IN MULTIWALL PAPER BAGS OR IN CLOTH BAGS) IN WAREHOUSES, STORAGE ROOMS AND SIMILAR LOCATIONS: Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Microsol machine, or use fixed ceiling sprayers that produce a very fine spray. Use a dosage of one fluid ounce of this spray per 1000 cubic feet of room space. Do not remain in treated areas. Ventilate the area after treatment is completed. Repeat application as necessary for insect control.

*Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Mediterranean Flour Moth and Indian Meal Moth.

TO KILL FLIES, MOSQUITOES AND GNATS: Close room and discharge spray mist as described above. Use a dosage of 1/2 to 1 fluid ounce per 1000 cubic feet of room space. Leave room closed for at least ten minutes. Sweep up and destroy fallen insects. Repeat as necessary for control. Do not remain in treated areas. Ventilate the area after treatment is completed.

IN PRESSURIZED TANK SPRAYERS FOR WETTING CONTACT SPRAYS AS A SURFACE SPRAY TO KILL THE ACCESSIBLE STAGES OF GRANARY INSECTS: RICE WEEVILS, CONFUSED FLOUR BEETLES, RUSTY RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, CADELLES, MEAL MOTH LARVAE AND ADULTS, CIG-

CHEMSCOPE

FOOD PLANT INSECTICIDE

SUPER STRENGTH

ACTIVE INGREDIENTS:

Pyrethrins	.50%
*Piperonyl butoxide, technical	1.00%
†N-octyl bicycloheptene dicarboximide	1.67%
Petroleum distillate	96.83%
	<hr/>
	100.00%

*Equivalent to .80% (butylcarbityl) (6-propylpiperonyl) ether and .20% related compounds.
†MGK 264, Insecticide Synergist.

E.P.A. REG. NO. 9143-52

E.P.A. EST. 9143-TX-1

NET VOLUME 1 GALLON
CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. SEE DIRECTIONS AND

to all areas infested by these insects. Use a dosage of 1 gallon per 1000 square feet. Pay particular attention to cracks and crevices in floor, walls and ceilings. Spray thoroughly around and into floor drains, conveyors and other areas where granary insects may be found. (For Flying Moths and Beetles, treat as for Flies, Mosquitoes and Gnats. See above.) **DO NOT APPLY THIS SPAY TO SURFACES THAT MAY COME IN CONTACT WITH FOOD; EXCESSIVE RESIDUES IN FOOD MAY RESULT.**

TO KILL ROACHES, SILVERFISH AND WATERBUGS IN SHELTERED SITUATIONS: Apply this spray with a sprayer that produces coarse, wetting droplets into all cracks and crevices in woodwork, floors, underneath sinks, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control. Do not allow spray to contact any stored food.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not apply directly to food. Cover or remove all food and food processing equipment during application. Do not apply space sprays while food processing is underway. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above. After spraying in meat packing plants, bakeries and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In case washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours. Remove pets, birds, and cover fish aquariums before spraying.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty containers. Destroy when empty and discard in a safe place. Keep container closed. Do not pour, store or spill near heat or open flame. NOTE: This spray when applied per the directions and cautions above, conforms with Food Additive Regulations 121.1074, 121.1075 and 121.1205 and the food additive tolerances established thereunder.

CHEMSCOPE CORPORATION

1909 Hi-Line Drive

P. O. Box 10752

Dallas, Texas

75207