

APR 15 1991

Mr. Rodger N. Cotman
CHEMSCOPE CORPORATION
3200 East Randol Mill Road
Arlington, TX 76011

Dear Mr. Cotman:

Subject: Label Revision
RID-ALL BRAND SPACE AND RESIDUAL INSECTICIDE
EPA Registration Number 9143-65
SWAT FLYING INSECT KILLER
EPA Registration Number 9143-60
CHEMSCOPE TOTAL RELEASE FOGGER
EPA Registration Number 9143-68
Submission dated March 21, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

1. For product 9143-65, revise the total inert ingredient percentage declaration on the label to include the "Related Compounds" percentage. Do not list the "Related Compounds" separately as an active ingredient.
2. For product 9143-65, ensure that the signal word "CAUTION" and the statement "Keep out of reach of children" are promptly displayed on the front panel and not obscured with other text.
3. For product 9143-60, the statement "CONTAINS PETROLEUM DISTILLATES" must appear on the label directly below the ingredients statement.
3. For product 9143-68, revise the "INERT INGREDIENTS" total percentage to include the "Related compounds" from d-trans a' ethrin. Do not combine the "Related compounds" percentage with the "d-trans ALLETHRIN" percentage as an active ingredient.

- 25
4. For product 9143-60, reinstate the isomer ratio substatement for d-cis/trans phenothrin.

Sincerely,

RFM Richard F. Mountfort
Product Manager (10)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

1 D-ALL
BRAND

SPACE & RESIDUE
INSECTICIDE

ACCEPTED
RECOMMENDATIONS
in EPA Letter Dated
APR 15 1981

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
Registered for the pesticide
9143-65
under EPA Reg. No.

ACTIVE INGREDIENTS:

* Resmethrin[5-(phenylmethyl)-3-furanyl]methyl- 3-(2-methyl-1-propenyl)cyclopropanecarboxylate	0.350% w/w
Related Compounds	0.048%
INERT INGREDIENTS	99.602%

	100.000% w/w

*Cis/trans isomers ratio: max. 30%(+)cis and min. 70%(+)trans.
Roussel Bio's SBP-1382 brand of resmethrin insecticide.
Roussel Bio Corp. 400 Sylvan Avenue, Englewood Cliffs, NJ 07632

KILLS: House flies, mosquitoes, gnats, wasps, roaches, earwigs, grain
mites, sowbugs, spiders, centipedes, ants, biting flies, aphids (plant
lice), leathoppers, exposed thrips, adult whiteflies, plant bugs,
Japanese beetles, spittlebugs.

FOR USE IN HOMES, MOTELS, INSTITUTIONS, RESTAURANTS.

SHAKE WELL BEFORE USING

E.P.A. EST. 9143-TX-1 E.P.A. REG. NO. 9143-65

VOLUME 20 FL. OZ.

NET WT. 17 OZ. (1 LB. 1 OZ)
METRIC WT. 482 GRAMS
KEEP OUT OF REACH OF CHILDREN
CAUTION
CONTENTS UNDER PRESSURE
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

BEST AVAILABLE COPY

CHEMSCOPE CORPORATION
3200 EAST RANDOL MILL ROAD
ARLINGTON, TX 76011
METRO (817) 640-5653

745

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ENVIRONMENTAL HAZARDS: The persticide is highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marches, and potholes). Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130 F may cause bursting. Never throw container into fire or incinerator.

STORAGE: Keep in locked storage area.

CONTAINER DISPOSAL: Replace cap and discard container in trash.

DOUBLE ACTION INSECT KILLER: This is a dual purpose formulation that can be used as a contact spray against flying and crawling insects and as a residual insecticide providing up to 3 to 4 weeks of control of roaches.

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

HOUSE FLIES, MOSQUITOES, AND GNATS: INDOORS: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

TO REDUCE ANNOYANCE FROM FLYING INSECTS: OUTDOORS: Reduce mosquitoes, gnats, midges, house flies, biting flies and flying moths (millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic feet of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

ROACHES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS AND CENTIPEDES: CONTACT SPRAY: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible. Repeat as necessary.

ROACHES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS AND CENTIPEDES: RESIDUAL SPRAY: Spray baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in the kitchen and bathroom. In basements, spray hiding places in cracks and crevices, lower walls and dark places. Spray at the rate of 3 to 4 seconds per linear foot at a distance of 12 to 14 inches from the surface. Repeat as necessary.

WASPS: Applications should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

ANTS: Apply to ant trails and around door sills, window frames and other places where ants may enter.

ORNAMENTAL PLANT PESTS: Spray aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittle bugs, Japanese beetles, mimosa webworm larvae, and oakworm larvae. Direct spray on insects by holding the container 12 to 18 inches from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides, covering foliage until slightly moist. Avoid heavy wetting. Repeat applications as necessary.

USE WITH CAUTION on the foliage of Flowering Quince, Juniper Cypress, Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus, as certain varieties have shown some leaf browning. When infested, spray smaller evergreens, deciduous trees and shrubs as rhododendrons, pine, spruces, hemlock, yews, arborvitae, junipers, holly, podocarpus, boxwood, laurel, lilac, spirea, dogwood, palm tree, mimosa, toyon, viburnum, oaks, maples locust, forsythia, althea and roses for control of aphids, and other pests listed above. Also spray infested flowering plants as peonies, iris zinnia, snapdragons, ageratums, chrysanthemums, nasturtium, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sago, broadleaf acacia for the same listed pests.

