DIRECTIONS

FOR DAIRY FARM USE

Teat Cups and liners should be dipped in a solution of Udder-San before being placed on the cow. It is recommended that cows' udders be washed before each milking and that teats be dipped immediately after milking.

Use one (1) ounce of Udder-San to each five (5) gallons of lukewarm water, or one-half (2) ounce of Lazarus Udder-San to a ten (10) quart pail of lukewarm water. Both dilutions provide 25 ppm available iodine.

It is recommended that this solution be sprayed onto udders with a "soft" garden-type nozzle or that it be applied by disposable paper towel (preferably Lazarus Roll Cow Towels where available).

DIRECTIONS

Udder-San is its own indicator of germicidal activity. The amber or yellow color shows the presence of the active ingredient - iodine. When the color fades, prepare a fresh solution. Lazarus Udder-San is not affected by water hardness or cold water.

CAUTION

In case of contact of the concentrate with the eye, flush with plenty of water. If irritation persists, get medical attention.

MADE IN U.S.A.

R.P. CONCENTRATE

FOR USE IN FORMULATING ECONOMIC POISONS (GERMICIDES) ONLY

ACTIVE INGREDIENTS:

α -(p-nonylphenyl)-omega hydroxypoly(oxyethylene)-iodine complex	
Polyethoxypolypropoxypolyethoxyethanol-iodine complex (Provides 3.85% Minimum Titratable Iodine)	
INERT INGREDIENT: U.S. Patent 3028299, 2977315, 2931777, 2759869	25.5%

NET CONTENTS: 50 GALLONS U.S.

DANGER: Keep Out Of Reach Of Children.

Harmful if swallowed. Do not take internally. Causes eye damage or skin irritation. Do not get in eyes. Wear goggles or protective shield when handling this product. Avoid contact with skin. In case of contact, flush thoroughly with water. For eyes, immediately get medical attention.

Rinse empty container thoroughly with water and discard it.



West agro-chemical, inc.

A SUBSIDIARY OF WEST CHEMICAL PRODUCTS, INC.

NEW YORK, N.Y. 11101 EPA Reg. No. 9148-8

7-05**83-A**092

Lot No.

0583