

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

September 14, 2005

Jane M. Reise
Regulatory Assistant
International Dioxide, Inc.
554 Ten Rod Road
North Kingstown, RI 02852

Subject: Carnebon 200
EPA Registration No. 9150-3
Application Date: July 20, 2005
Receipt Date: August 18, 2005

Dear Ms. Reise:

This acknowledges receipt of your notification, submitted under the provision of PR Notice 98-10, FIFRA section 3(c)9.

Proposed Notification

- clarification of Directions For Use

General Comments

Based on a review of the material submitted, the following comments apply:

The notification application is acceptable and a copy has been inserted in your file for future reference.

Should you have any questions or comments concerning this letter, please contact me at (703) 308-6345.

Sincerely,

Wanda Y. Henson
Product Reviewer (32)
Regulatory Management Branch II
Antimicrobials Division (7510C)

CONCURRENCES

MBOL	750C	750C						
NAME	E. Berg	Henson						
TE	9/14/05	9-14-05						



International Dioxide, Inc.
554 Ten Rod Road
North Kingstown, RI 02852
401-295-8800 Tel 401-295-7108 Fax

July 20, 2005

Document Processing Desk
Ms. Emily Mitchell
Antimicrobials Division (7510C)
US Environmental Protection Agency
Office of Pesticide Programs
Ariel Rios Building
1200 Pennsylvania Avenue NW
Washington, DC 20460

Reference: Carnebon 200®, EPA Registration No. 9150-3; NOTIFICATION

Dear Ms. Mitchell,

Please find the attached EPA Form 8570-1 and two copies of revised labeling submitted in support of notification of label change in accordance with PR Notice 98-10 II.M.3. for the subject product. Also attached is a one copy of the label highlighting the change (refer to page 3) in the directions.

The revised labeling incorporates, under "Preparation of Use-Solution" Tuberculocidal and Virucidal, a 10% acid activator solution.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Should you have any questions, feel free to call.

Sincerely,

Jane M. Reise
Regulatory Assistant
International Dioxide, Inc.



Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 9150-3	2. EPA Product Manager Emiy Mitchell	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Carnebon 200 (R)	PM#	
5. Name and Address of Applicant (Include ZIP Code) International Dioxide Inc. 554 Ten Rod Road North Kingstown, RI 02852 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Correction of acid used to activate the Camobon 200 product. "This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container		<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 55 gal 5 gal 1 gal		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product				<input checked="" type="checkbox"/> Lithograph Paper glued Stenciled <input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jane M. Reise	Title Regulatory Assistant	Telephone No. (Include Area Code) 401 294 6803
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 		
3. Title REGULATORY ASSISTANT		
4. Typed Name JANE M. REISE		5. Date July 20, 2005

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

CHEMICAL AND PHYSICAL HAZARDS

Chlorine dioxide is a strong oxidizing agent. Contamination with other materials such as acids, chlorine, organic chemicals, etc. may cause a chemical reaction resulting in evolution of chlorine dioxide and heat. Explosion and/or fire could result. Chlorine dioxide is a poisonous explosive gas. Keep all chemical and foreign materials away from this solution.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


STORAGE: Do not store with easily oxidizable materials, acids, reducers, and combustible material. Avoid heat or freezing conditions. Store upright and do not stack drums over two high on pallets or partially filled drums. Use of a drum pump is suggested. Keep drum tightly closed when not withdrawing liquid. In case of spills, dilute with large quantities of water. Do not allow liquid to dry because this could present a fire hazard. Store only in the original container and take care to prevent cross-contamination with other pesticides, fertilizers, food and feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EMERGENCY HANDLING: In case of contamination or decomposition, do not reseal container. Isolate in open, well-ventilated area. Flood with large volumes of water. Cool unopened drums in vicinity by water spray.

NOTICE: Seller expressly warrants that the product conforms to its chemical description. There are no warranties associated with the sale of the product either express or implied, including but not limited to the warranties of fitness for a particular purpose or use.



Active Ingredient:
Chlorine Dioxide..... 2.0%
Inert Ingredients..... 98.0%
100%

[Hospital Disinfectant - Food-Contact Surface Sanitizer - Food Processing Water Sanitizer - Disinfectant - Sanitizer - Virucide - Tuberculocide

Hospital Use- Farm Premise Sanitation - Food Processing Water - No rinse Sanitizer - Institutional Use - Non-Flammable - Eliminates Mold and Mildew - Mold and Mildew Control Combats Mold and Mildew - Concentrated Broad Spectrum Biocide - Farm Premise Sanitatic Poultry Premise Sanitation - Animal Laboratory Disinfection

**KEEP OUT OF THE REACH OF CHILDREN
CAUTION**

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

<i>First Aid</i>
<u>If on skin or clothing:</u> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
<u>If in eyes:</u> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
<u>If swallowed:</u> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
<u>If inhaled:</u> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
For 24 hour emergency information on this product, call Chemtrec at 1-800-424-9300 (US, Canada, Puerto Rico, Virgin Islands) 1-703-627-3887 (All Other Areas)
Hotline: You may also contact the National Poison Control Center at 1-800-222-1212 for Emergency Medical Advice Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

REG. NO. 9150-3

E.P.A. EST. NO. 9150-RI-01

NET WT. _____ LBS.

Carnebon 200® is a registered trademark of International Dioxide Inc., a DuPont Company.

© 2005. E.I. du Pont de Nemours and Company. All rights reserved.

Manufactured by:



International Dioxide, Inc.
554 Ten Rod Road
North Kingstown, RI 02852

4/22

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Carnebon 200® is a specially designed formulation of chlorine dioxide and is a uniquely versatile biocide. It controls microbial contamination in animal research facilities, food-processing and industrial waters, pulp and papermaking processing waters and cutting oils. It also disinfects environmental surfaces in hospitals and institutions and sanitizes food-contact surfaces. **Carnebon 200®** is highly effective against mold and mildew.

Carnebon 200® delivers a non-corrosive disinfectant and cleaning performance in an economical concentrate.

Carnebon 200® meets AOAC efficacy standards for hospital disinfectants and food-contact surface sanitizing solutions.

Carnebon 200® can be used in federally inspected meat and poultry plants as both a disinfectant and food-contact surface sanitizer.

Carnebon 200® is the product of choice for HAACP programs.

The efficacy of **Carnebon 200®** depends on the degree of activation. Unactivated **Carnebon 200®** effectively controls microbes in processing waters and mold and mildew. For disinfection and sanitization, **Carnebon 200®** must be activated. Read the activation instructions carefully prior to using **Carnebon 200®**.

Carnebon 200® can be used to treat hard, non-porous surfaces and water systems in: hospitals, medical and dental offices, food processing facilities, bottling plants, breweries, meat-packing plants, poultry-processing plants, fish-processing plants, food storage areas, institutional kitchens, dairy and poultry farms and production facilities, mushroom production facilities, animal research facilities, agricultural storage facilities (including containers, trailers, rail cars, vessels and bins), animal transport vehicles and equipment, animal confinement and rearing facilities, animal handling facilities, egg processing plants, livestock facilities, hatcheries, hotels, business and office buildings, institutional facilities, public facilities.

Carnebon 200® is an effective disinfectant against the following bacteria at a 300 ppm activated use-solution of **Carnebon 200®** (~ 30 ppm free chlorine dioxide) in 10 minutes in the presence of 5% organic serum.

- Pseudomonas aeruginosa (Pseudomonas)
- Staphylococcus aureus (Staph)
- Salmonella cholerasuis (Salmonella)

Carnebon 200® is tuberculocidal (effective against Mycobacterium bovis, BCG) at a 1200 ppm activated use-solution of **Carnebon 200®** (~200 ppm free chlorine dioxide) in 10 minutes at 20 deg C.

Carnebon 200® is an effective virucide against the following viruses at a 800 ppm activated use-solution of **Carnebon 200®** (~100 ppm free chlorine dioxide) in 10 minutes, 15 minutes for Canine parvovirus ATCC VR-2017.

- HIV-1 (AIDS Virus) HTLV-III_B
- Canine parvovirus ATCC VR-2017
- Rat coronavirus RCV-SDA-681
- Mouse hepatitis virus MHV-A59
- Minute virus of mice MVM-P
- Parainfluenza virus, Type 1 ATCC VR-105 SENDEI/52

Carnebon 200® is an effective sanitizer against Salmonella typhi at a 100-200 ppm activated use-solution of **Carnebon 200®** in 30 seconds.

An unactivated use-solution of 1000 ppm of **Carnebon 200®** effectively controls mold and mildew in 60 seconds.

Preparation of Carnebon 200® Use-Solutions

Disinfectant Use-Solution

Prepare an activated 300 ppm use-solution of Carnebon 200® by using **one** of the three procedures described below.

- 1) Add 1 part Carnebon 200® to 64 parts water and then adjust the pH of the diluted Carnebon 200® to 2.6 with either acetic, citric, phosphoric, sulfuric, hydrochloric or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Prepare in a well-ventilated area and avoid breathing any fumes which may be produced during activation.

Alternatively to minimize worker handling, an automated system can also be utilized that will safely activate the concentrate of Carnebon 200® with any of the various acids listed to deliver the proper pH and safety dilute the material to the 300 ppm working solutions.

- 2) Add 2 fl. oz. of Carnebon 200® to one (1) gallon of water into a clean plastic pail and add 1.2 grams of Activator C or 8.6 grams of Activator K. Allow 15 minutes reaction time and for the activator to completely dissolve. Prepare in a well-ventilated area and avoid breathing any fumes, which may be produced during activation.
- 3) An activated 300 ppm use-solution of Carnebon 200® can also be prepared electrolytically by adding Carnebon 200® directly to the OXYCHLOR® e-generator. For proper operation, of the OXYCHLOR® e-generator, consult the OXYCHLOR® e-generator manual or your IDI or authorized representative.

Tuberculocidal Use-Solution

Prepare an approximate 1200 ppm use-solution of Carnebon 200® (200 ppm of free chlorine dioxide) by adding 1 part of Carnebon 200® into a clean, plastic pail and then add 5 parts of a 10% acid activator solution. The acid activator can be either acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic, or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Allow 60 minutes for reaction time and for the activator to completely dissolve. Then dilute the activated solution with 12 parts of water. Prepare in a well-ventilated area and avoid breathing any fumes that may be produced during activation.

Virucidal Use-Solution

Prepare an approximate 800 ppm use-solution of Carnebon 200® (100 ppm of free chlorine dioxide) by adding 1 part of Carnebon 200® into a clean, plastic pail and then add 5 parts of the 10% acid activator solution. The acid activator can be either acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic, or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Allow 15 minutes for reaction time and for the activator to completely dissolve. Then dilute the activated solution with 20 parts of water.

Food-Contact Surface Sanitizing Solution

Prepare a 200 ppm activated use-solution of Carnebon 200® by using **one** of the three procedures described below:

- 1) Add 1 part Carnebon 200® to 3 parts water and then activate by adding food-grade citric, phosphoric, acetic or other equivalent food-grade acid (of at least 99% purity) to a pH of 2.6. Please contact your IDI or authorized representative regarding equivalent acids. Agitate for 5 minutes and then allow to stand for 15 minutes. Then dilute 1 part of the activated solution with 24 parts of water.

Alternatively to minimize worker handling, an automated system can also be utilized that will safely activate the concentrate of Carnebon 200® with any of the various acids listed to deliver the proper pH and safety dilute the material to the 200 ppm working solutions.

- 2) Add 1 gallon of Carnebon 200® to 20 gallons of water followed by 12 grams of Activator K. Allow to stand for 15 minutes after agitation for 5 minutes and then dilute 1 part with 4 parts of water.
- 3) An electrolytically activated use-solution can be prepared by adding Carnebon 200® directly to the OXYCHLOR® e-generator. The activated use-solution prepared by the OXYCHLOR® e-generator must contain between 50-100 ppm of an activated use-solution of Carnebon 200®. For proper operation of the OXYCHLOR® e-generator, consult the OXYCHLOR® e-generator system manual or your IDI or authorized representative.

Mold & Mildew Use-Solution

Prepare a 1000 ppm use-solution of Carnebon 200[®], by placing 1 part Carnebon 200[®] per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail or drum. Dilute with clean, potable water.

APPLICATION INSTRUCTIONS:

FOOD PROCESSING PLANTS, FOOD-HANDLING ESTABLISHMENTS AND RESTAURANTS

Carnebon 200[®] can be used to:

- To control microbial contamination, slime and odor in food processing waters.
- To sanitize food processing equipment and surfaces in food processing and food-handling establishments.
- To sanitize food-contact surfaces and utensils in food-handling establishments.
- To disinfectant non-food contact surfaces in food-processing plants, food handling establishments and restaurants.
- For use as a terminal food-contact surface sanitizer rinse conforming to 40 CFR 180.940 (b) and (c) Food Contact Surface Sanitizing Solutions.

Specific Applications

Use Carnebon 200[®] as a Terminal Sanitizing Rinse for Stainless Steel Tanks, Transfer Lines, On-line Equipment, Recirculating and Clean-in-Place (CIP) systems, Food-contact surfaces and similar surfaces, such as tables, trays, bins, etc., utensils and Food-Processing Equipment in Poultry, Meat, Fish & Meat Processing Plants, Dairies, Bottling Plants, Restaurants, Canneries and Breweries

- 1) Prior to sanitization, remove all gross food particles and soil by use of a pre-flush, pre-scrape or pre-soak treatment. Then clean all lines, tanks, or surfaces with a suitable detergent followed by a potable water rinse.
- 2) Prepare the **Food-Contact Surface Sanitizing Solution** as described above.
- 3) Fill, immerse, circulate, wipe or spray the target surface with the sanitizing solution making sure the surface area is thoroughly wet for at least one minute. Hard to reach in-place equipment, pipes, closed vessels, etc., should be filled with the sanitizing solution to ensure contact of all surfaces. Use suitable protective breathing apparatus when spraying the solution on external equipment.
- 4) Allow the sanitizing solution to drain from all treated surfaces and air dry. Do not rinse treated surface.
- 5) The above solution may not be reused for sanitizing but may be diluted to 1:5 with water and used for cleaning of walls, floors and drains of the plant.

Use Carnebon 200® in Food Processing Plants to Control Odor and Slime Forming Bacteria in Cooling and Warming Waters Such as Canning Retort and Pasteurizer Cooling Water Used to Decrease or Increase Packaged Product Temperature By Immersion in or by Spraying with the Treated Process Waters

- 1) All tanks, tunnels, conveyor chains, heat exchangers, heat exchange towers, lines, spray bars, and nozzles should be thoroughly cleaned when possible, and completely rinsed using clean, potable water prior to treatment.
- 2) **Preparation and Application of Use-Solution:** Water systems, including the cooling or warming tanks or spray systems, towers, lines and all water containing parts of the system may be batch loaded at start-up with one part Carnebon 200® per 4000 parts potable water (5 ppm available chlorine dioxide). To maintain the 5 ppm available chlorine dioxide in the water system, a timed or electronically controlled chemical feed pump or injector system can be used for additions to the system or for treating the make-up water. Make up new Carnebon 200® solutions daily.
- 3) **Preparation and Application of Optional Activated Use-Solution (acid activation):** If heavy use of cooling or warming water or introduction of additional bacteria loads are expected, or if slime buildup is heavy, an additional activation step may be used in preparation of the use-solution. Prepare the activated use-solution in a well-ventilated area and avoid breathing any fumes which may be produced while crystals are dissolving. For each one thousand (1,000) gallons of system water to be treated, measure one quart (0.95 liters) of Carnebon 200® and pour into a clean plastic container, pail or drum. To this Carnebon 200® amount, add food grade citric acid of no less than 99% purity, at the rate of 3.3 oz. (95 grams or 0.2 lbs.) of crystals per quart (0.95 liters) of Carnebon 200®. Allow five (5) minutes reaction time for crystals to dissolve. Cooling or warming water systems may be batch loaded at start up using one quart (0.95 liters) of the activated solution per one thousand (1,000) gallons of potable water (5 ppm available chlorine dioxide). Batch or timed additions of the activated solution can be made or an electronically controlled chemical feed pump or injector system can be used for additions of the activated solution to the process water to maintain 5 ppm available chlorine dioxide. Make up new Carnebon 200® solutions daily.
- 4) **Preparation and Application of Optional Activated Use-Solution (Oxychlor e-generator):**
An activated use-solution can also be prepared electrolytically by adding Carnebon 200® directly to the Oxychlor e-generator. Add the activated use-solution prepared by the Oxychlor e-generator to water systems, including cooling or warming tanks, or spray systems, tower lines and to all water containing parts of the system. Batch load these systems at start-up and maintain a concentration of 5 ppm available chlorine dioxide in the system. For proper operation of the Oxychlor e-generator, consult the Oxychlor e-generator system manual or your International Dioxide representative.

Note: Chemical feed pumps and injectors must be chlorine resistant for best operation. Available chlorine dioxide levels should be confirmed using an International Dioxide chlorine dioxide test kit.

Use Carnebon 200® in Food Processing Plants to Control the Build-Up of Odor and Slime Forming Bacteria in Stainless Steel Transfer Lines and On-Line Equipment Such as Hydrocoolers, Pasteurizers and the Like Overnight and Over Weekends.

- 1) Clean equipment or line thoroughly using a suitable detergent followed by a clean, potable water rinse before treatment.
- 2) **Preparation and application of solution:** For each 1000 parts of solution in lines and/or equipment, add 1 part of Carnebon 200® (20 ppm available chlorine dioxide) to potable make-up water. Mix and fill lines and equipment overnight. Drain and allow to air dry just prior to next run start-up.

Carnebon 200® to Control the Buildup of Odor and Slime Forming Bacteria in Process Waters for Vegetable Rinses and Associated Tanks, Flumes and Lines.

- 1) All tanks, flumes and lines should be thoroughly cleaned with a suitable detergent and completely rinsed using clean, potable water prior to treatment.
- 2) Preparation and Application of Use-Solution: Chill tanks or vegetable rinse tanks may be batch loaded at start-up with 1 part Carnebon 200® per 4000 parts of potable water (5 ppm available chlorine dioxide). Make-up waters should be treated using a chemical feed pump or injector system and applied at the rate of 1 part Carnebon 200® per 4000 parts potable water. Make up new Carnebon 200® solutions daily.
- 3) Preparation and Application of Optional Activated Use-Solution: If heavy use of rinse water is expected, or if slime buildup is extreme, an additional activation step may be used in preparation of solution. Prepare the activated use-solution in a well-ventilated area and avoid breathing any fumes which may be produced while crystals are dissolving. For each 10 gallons of rinse water to be used, measure 1/3 fl. oz. (10 ml) of Carnebon 200® and pour into a clean, plastic container containing 1 gallon of water. Activate this solution by:
 - 1) Adding 0.002 grams of Activator-C or
 - 2) Adding 2.2 grams of Activator K or
 - 3) Adjusting the pH to 2.6 with acetic acid, citric acid, phosphoric acid, sulfuric acid or hydrochloric acid.

Allow this solution to stand for 15 minutes and then add to 9 gallons of water to give 5 ppm available chlorine dioxide. Chill tanks or vegetable rinse tanks may be batch loaded at start-up with the activated Carnebon 200® solution 1/3 fl. oz. (10 ml) per ten (10) gallons of potable water (5 ppm available chlorine dioxide). Make-up waters should be treated using a chemical feed pump. In order to ensure accurate delivery, a 1 to 10 dilution of the active concentration should be made and the feed rate of 3 1/3 fl. oz. of activated Carnebon 200® solution per ten (10) gallons should be maintained. Make up fresh Carnebon 200® solutions daily.

- 4) Preparation and Application of Optional Activated Use-Solution (Oxychlor e-generator): An activated use-solution can also be prepared electrolytically by adding Carnebon 200® directly to the Oxychlor e-generator. Add the activated use-solution prepared by the Oxychlor e-generator to chill tanks or vegetable rinse tanks. Batch load these systems at start-up and maintain a concentration of 5 ppm available chlorine dioxide in the system. For proper operation of the Oxychlor e-generator, consult the Oxychlor e-generator system manual or your IDI representative.

Note: Chemical feed pumps and injectors must be chlorine resistant for best operation. Available chlorine dioxide levels should be confirmed using an International Dioxide chlorine dioxide test kit.

Use Carnebon 200® to Extend Freshness and Shelf Life of Fruits and Vegetables

- 1) Before treatment, whole fruits and vegetables should be washed and thoroughly rinsed with clean, potable water.
- 2) In a one (1) gallon container, add 1/3 fl. oz. (10 ml) of Carnebon 200® and add 0.002 grams of Activator-C or adjust the pH to 2.6 with vinegar. Allow to stand for 15 minutes then add to 9 gallons of water.
- 3) Pretreatment for Uncut, Unpeeled Fruits and Vegetables: Dip uncut, unpeeled fruits and vegetables in treatment solution for about ten (10) to twenty (20) seconds, then follow with a potable water rinse.

Use Carnebon 200® To Disinfect Non-Food Contact Surfaces (Walls, Ceilings, Drains and Floors) in Food Processing Plants and Food-Handling Establishments.

- 1) Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
- 2) Prepare the Disinfectant Use-Solution as described above.
- 3) Apply the disinfectant use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge or sprayer, or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled surfaces, a pre-cleaning is recommended.
- 4) For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface rub with a brush, sponge or cloth. Do not breathe spray. Make sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.
- 5) After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.

Use Carnebon 200® To Control Mold & Mildew and Slime Forming Bacteria on Non-Food Contact Surfaces (Floors, Walls, Ceilings and Drains) in Food-Processing Plants and Food-Handling Establishments

- 1) Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
- 2) Follow the directions for **Mold & Mildew Use-Solution**: as described above
- 3) **Application**: Drench, spray or fog solution on walls, floors, ceilings and surfaces using a suitable watering, spraying or fogging device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying or fogging, the area should be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.
- 4) Repeat application as needed.

TRANSPORT VEHICLES

To Disinfect Hard, Non-porous Surfaces in Vehicles Including Animal Transport Vehicles, Rail Cars, Trailers and Vessels. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.

1. Prior to application of Carnebon 200®, clean all vehicles with high-pressure water and a suitable detergent.
2. Follow directions for **Disinfectant Use-Solution** as described above.
3. Then apply the disinfectant use-solution to all surfaces to be treated. All treated surfaces must remain wet for at least 10 minutes.

SEAFOOD USE

Use Carnebon 200® as an Bacteriostat for Treating Ice Used for Icing Fish in the Round

Carnebon 200® may be batch loaded or metered into makeup water used to produce ice for icing fish in the round. Prepare a non-activated working solution containing 20 ppm of available chlorine dioxide by adding 1 part of Carnebon 200® to 1000 parts of water.

HOSPITALS, INSTITUTIONS, MEDICAL AND DENTAL CLINICS, and VETERINARY CLINICS.

Note: The Oxychlor e-generator has not been tested against *Pseudomonas aeruginosa*. The Oxychlor e-generator is not approved for use in hospitals, laboratories, morgues, and other institutions.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Carnebon 200® can be used to:

- To disinfectant environmental surfaces.
- To control mold and mildew on environmental surfaces.
- To control animal viruses on environmental surfaces.
- To control odor and slime forming bacteria.

Special Applications

To Disinfect Walls, Ceilings and Floors and other Environmental Surfaces in Hospitals, Institutions Veterinary Clinics, and Animal Research Facilities

- 1) Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
- 2) Prepare the **Disinfectant Use-Solution** as described above.
- 3) Apply the disinfectant use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge or sprayer, or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled surfaces, a pre-cleaning is recommended.
- 4) For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface rub with a brush, sponge or cloth. Do not breathe spray. Make sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.
- 5) After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OR SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS that involve healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS). Carnebon 200® destroys HIV-1 (AIDS Virus) HTLV-III_B on precleaned environmental surfaces/objects previously soiled with blood or other body fluids in ten minutes contact

Personal Protection: The worker should wear protective equipment such as disposable latex or rubber gloves, gowns, masks and eye protection to prevent contamination from items soiled with blood or body fluids.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of Carnebon 200®.

Contact Time: Allow Carnebon 200® to contact treated items for 10 minutes to kill HIV-1. This time may not control other common types of viruses and bacteria.

Disposal Of Infectious Material: Any blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

To Control Mold & Mildew and Slime Forming Bacteria on Walls, Floors, Ceilings, and Surfaces and other Environmental Surfaces

- 1) Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
- 2) Follow the directions for **Mold & Mildew Control Use-Solution** as described above
- 3) **Application:** Drench, spray or fog solution on walls, floors, ceilings and surfaces using a suitable watering, spraying or fogging device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying or fogging, the area should be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.
- 4) Repeat application as needed.

To Disinfect Non-Porous, Hard Surfaces Such as Tile Floors, Walls and Ceilings and Stainless Steel Cold Rooms and Walk-In Incubators

- 1) Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above.
- 3) **Application of Activated Disinfection-Solution:** Activated solutions may be sprayed, mopped or sponged onto surfaces to be disinfected. All surfaces must be thoroughly wetted for at least ten (10) minutes. When spraying disinfectant solutions, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.

Use In Dental Offices and Laboratories as a Dental Pumice Disinfectant

- 1) Prepare solution in a well-ventilated area. To make one (1) liter of solution, pour 1.0 fl oz (approximately 25 mls.) of Carnebon 200® into a clean glass or plastic container. To this, add 2 ½ grams (1/2 teaspoon) of citric acid crystals (included) and mix slightly, allowing 5 minutes reaction time and for crystals to dissolve completely. Avoid breathing any fumes which may be produced during activation. Once solution has yellowed, diluted to one (1) liter with clean potable water, for a working solution of 500 ppm activated ClO₂.
- 2) To apply: The working solution can be conveniently contained in a one (1) liter plastic "squeeze" bottle for up to five days. Apply to dry pumice powder exactly as water to produce the pumice slurry. Apply additional working solution as needed to reconstitute dried out slurry to appropriate viscosity. Carnebon 200® solutions should be made up fresh, preferably on Monday and discarded on Friday or 5 days after preparation.

As a Virucide to Kill Animal Viruses (Rat Coronavirus RCV-SDA-681, Mouse Hepatitis Virus MHV-A59, Minute Virus of Mice MVM-P and Canine Parvovirus ATCC VR-2017) Parainfluenza Virus Type 1 ATCC VR-105 SENDAI/52, HIV-1 HTLV-III_B) on Non-Porous, Hard Surfaces Such as Tile Floors, Walls and Ceilings and Stainless Steel Cold Rooms and Walk-In Incubators.

- 1) Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
- 2) Follow the directions for **Virucidal Use-Solution** as described above.
- 3) **Application of Activated Use-Solution:** Activated solutions may be sprayed, mopped or sponged onto surfaces to be treated. All surfaces must be thoroughly wetted for at least ten (10) minutes (15 minutes contact time for canine parvovirus ATCC VR-2017). When spraying the virucidal solution, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.

To Disinfect Bench Tops, Biological Hoods, Incubators, Stainless Steel Equipment and Instruments.

- 1) Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above
- 3) **Application of Activated Use-Solution:** Activated solutions may be squirted directly onto surfaces from a plastic squeeze bottle or may be used as a soak solution. All contact surfaces must be thoroughly wetted for at least ten (10) minutes. Allow to air dry. Activated solutions of Carnebon 200®, stored in plastic squirt bottles, may be held up to one (1) week before replacement with fresh solution. Soak solutions of Carnebon 200® should be changed daily.

To Disinfect Surfaces of Water Baths and Tubs

- 1) Prior to disinfection, thoroughly clean the bath or tub with a suitable detergent and rinse with clean water.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above.
- 3) **To apply:** Turn circulating motor on and allow the water to circulate for at least (10) minutes. Drain tub completely. After the draining is finished, tub is ready for use.

To Disinfect Water Bath Incubators

- 1) Prior to disinfection, thoroughly clean the reservoir with a suitable detergent and rinse with clean water.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above.
- 3) **To apply:** Activated solution should be poured into waterbath reservoir and allowed to stand one (1) hour at room temperature. Drain reservoir and fill with fresh water.

To Control Odor and Slime Forming Bacteria in Water Bath Incubators

- 1) When using Carnebon 200® in waterbath incubators, always begin with a freshly cleaned and disinfected reservoir.
- 2) **Application:** Fill water bath with clean, potable water to near capacity. Add 1 part Carnebon 200® for each 400 parts water (50 ppm available chlorine dioxide). When water becomes cloudy, discard water and repeat procedure.

Control Odors Resulting from the Sterilization of Spent Biologicals in Steam Autoclaves

- 1) To reduce autoclave odors of used biologicals, Carnebon 200[®] should be sprayed or poured directly into the stainless steel autoclave buckets.
- 2) Preparation of Use-Solution: Place 1 part Carnebon 200[®] per 20 parts working solutions (1,000 ppm available chlorine dioxide) into a clean glass or plastic container and mix
- 3) Application: Spray or pour Carnebon 200[®] solution into or onto the autoclave buckets just prior to autoclaving.

To Deodorize Animal Holding Rooms, Sick Rooms, Morgues and Work Rooms

- 1) Rooms to be deodorized should be in a clean condition prior to Carnebon 200[®] application.
- 2) Preparation of Use-Solution Place 1 part Carnebon 200[®] per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean glass or plastic container. 3) Application: Spray solution using a suitable spraying device onto walls, ceilings and floors, lightly dampening all surfaces. Avoid breathing mist of solutions by using an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide. Allow to air dry, and then ventilate the area. Treat as required.

IN ANIMAL REARING & CONFINEMENT FACILITIES**To Disinfect Hard, Non-Porous Surfaces in Commercial Animal Confinement Facilities such as Poultry Houses, Swine Pens, Calf Barns and Kennels and for use in Laboratory Animal Breeding and Research Quarters for Controlling Cross-Contamination of Microorganisms Infectious to these Animals and Humans from Treated Surfaces**

- 1) Remove all animals and feed from premises, vehicles, enclosures, coops and crates.
- 2) Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
- 3) Empty all troughs, racks and other feeding and watering appliances.
- 4) Thoroughly clean all surfaces with soap and detergent and rinse with water.
- 5) Follow the directions for Disinfectant Use-Solution as described above.
- 6) Application: Using a commercial sprayer, saturate all surfaces with the activated Carnebon 200[®] solution for a period of ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure.
- 7) After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate when solution has dried.
- 8) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.

As a Virucide to Kill Animal Viruses (Rat Coronavirus RCV-SDA-681, Mouse Hepatitis Virus MHV-A59, Minute Virus of Mice MVM-P and Canine Parvovirus ATCC VR-2017) on Non-Porous, Hard Surfaces in Commercial Animal Confinement Facilities Such as Poultry Houses, Swine Pens, Calf Barns, and Kennels and in Laboratory Animal and Research Quarters

- 1) Remove all animals and feed from premises, vehicles, enclosures, coops and crates.
- 2) Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures traversed by animals.
- 3) Empty all troughs, racks and other feeding and watering appliances.
- 4) Thoroughly clean all surfaces with soap and detergent and rinse with water.
- 5) Follow the directions for Virucidal Use-Solution as described above.
- 6) Application of Activated Use-Solution: Activated solutions may be sprayed, mopped or sponged onto surfaces to be treated. All surfaces must be thoroughly wetted for at least ten (10) minutes (15 minutes contact time for canine parvovirus ATCC VR-2017). When spraying virucidal solution, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.
- 7) After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate when solution has dried.
- 8) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.

To Control the Build-up of Odor and Slime Forming Bacteria in Animal Confinement Areas

- 1) Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, cases and other facilities and fixtures occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with clean water.
- 2) Preparation of Use-Solution: Place 1 part Carnebon 200[®] per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail.
- 3) Application: Using a commercial sprayer; saturate all surfaces with the Carnebon 200[®] solution. When spraying Carnebon 200[®] solutions, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide to avoid breathing mist.

To Control Animal Odors on Pets and in Litter Boxes, Carpets and Concrete Floors

- 1) For litter boxes: Wash out litter boxes with suitable detergent and rinse with clean, potable water. Soak overnight in solution of one part Carnebon 200[®] per 32 parts of water (625 ppm available chlorine dioxide). Add litter, sprinkle surface liberally with Carnebon 200[®] solution.
- 2) For controlling odors in carpets: Add 1 part Carnebon 200[®] per 40 parts (500 ppm available chlorine dioxide) of rug shampoo mix. Shampoo carpets. Allow to air dry. **NOTE:** Carnebon 200[®] may bleach some carpets and fabrics, especially if applied on top of another chemical agent. Do not apply until a sample test has been tried and observed for at least 24 hours.
- 3) For concrete floors: Clean floor thoroughly using a suitable detergent; rinse with clean water. Prepare solution by adding 1 part of Carnebon 200[®] per 16 parts of water (1,250 ppm available chlorine dioxide). Mop or spray solution liberally onto floor. Allow to air dry.
- 4) For animal baths: Wash animal well with appropriate pet shampoo; rinse with clean water. Prepare solution by adding 1 part Carnebon 200[®] per 200 parts of water (100 ppm available chlorine dioxide). Rinse animal thoroughly with prepared solution. Allow to air dry. Avoid direct contact with animal's eyes, nose and ears.
- 5) For treating animal odors with high levels of ammonia: Wash area thoroughly with suitable detergent and rinse with clean water. Preparation of solution: For each 32 parts of solution place 1 part Carnebon 200[®] into a clean, plastic container. To this concentrate, add 1 tablespoon household bleach and allow to react for five (5) minutes. Dilute with 32 parts of clean, potable water. Apply by mopping or spraying solution liberally onto area. Allow to air dry. Additional applications may be necessary.

To Control Mold & Mildew and Slime Forming Bacteria on Walls, Floors, Ceilings, Bins, Boxes, Pens, Barns, Kennels and other Animal Health Surfaces

- 1) Remove animals and feed from area to be treated
- 2) Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
Follow the directions for Mold & Mildew Use-Solution as described above
- 4) Application: Drench, spray or fog solution on walls, floors, ceilings and surfaces using a suitable watering, spraying or fogging device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying or fogging, the area should be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces.
- 5) Follow treatment with a potable water rinse
- 6) Repeat application as needed.

ANIMAL TRANSPORT VEHICLES

To Disinfect Hard, Non-Porous Surfaces in Vehicles Including Animal Transport Vehicles, Rail Cars, Trailers and Vessels. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.

1. Prior to application of Carnebon 200[®], clean all vehicles with high-pressure water and a suitable detergent.
2. Follow directions for Disinfectant Use-Solution as described above.
3. Then apply the disinfectant use-solution to all surfaces to be treated. All treated surfaces must remain wet for at least 10 minutes.

15/22

T1. TREATMENT OF WATER STORAGE SYSTEMS AND POTABLE WATER

To Disinfect Water Storage Systems Aboard Aircraft, Trains, Buses, Boats, RV's, Off-Shore Oil Rigs, etc.

- 1) Prior to disinfection, tanks should be cleaned using a suitable detergent and thoroughly flushed with clean, potable water.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above.
- 3) Pour activated solution into tank, filling the tank completely. Bleed air out of lines and allow to stand at least ten (10) minutes. Drain tank and lines and flush with potable water.

To Control Build-Up of Slime and Odor Causing Bacteria and Enhance the Taste of Stored Potable Water

- 1) Prior to treatment of potable water, thoroughly clean and disinfect the water storage system to ensure a sanitary condition. Thoroughly rinse with clean, potable water.
- 2) Potable water should be treated at a rate of one part of Carnebon 200[®] per 4000 parts potable water (5 ppm available chlorine dioxide) and may be injected or batch treated.
- 3) Water storage tank should be sufficiently sealed to prevent outside contamination and direct sunlight.

Using an International Dioxide test kit, confirm the chlorine dioxide to be 5 ppm and check to see this level does not fall below 1 ppm.

To Help Remove Off-Odors and Tastes from Municipal Well Waters

- 1) Carnebon 200[®] should be injected into the incoming water main using a chemical proportioning pump, or injector, at a rate of one 1 part Carnebon 200[®] per 20,000 parts water (1 ppm available chlorine dioxide).
- 2) Confirm pump or injector accuracy using an International Dioxide test kit and adjust accordingly.
- 3) Carnebon 200[®] levels should be checked weekly.

INDUSTRIAL WATER SYSTEMS AND INDUSTRIAL BIOCIDES USE

TO INHIBIT BACTERIAL SLIME FORMING BACTERIAL BUILDUP IN COMMERCIAL WATER FILTRATION SYSTEMS, SAND BEDS, GRAVEL BEDS, CHARCOAL FILTERS AND COOLING WATER SYSTEMS.

Filters:

- 1) Carefully back-flush filters with potable water, where possible, to remove any accumulated solid residue and contamination.
- 2) Fill system with potable water and adjust pH to 6.0 with citric acid, phosphoric acid, or acetic acid (vinegar) or equivalent.
- 3) Add 1 part of Carnebon 200[®] per 64 parts of (300 ppm of available chlorine dioxide) of filter system volume to the access hatch and circulate the system for 1 hour. Check the pH and bring back to 6.0 if it has drifted. Bring the available chlorine dioxide concentration back to 300 ppm.
- 4) Circulate the solution for 1 additional hour, discharge and then water wash for 30 minutes with potable water to remove the chlorine dioxide.

For Enclosed And Recirculating Cooling Water Systems

- 1) Add 2.5-10 gallons of Carnebon 200[®] per 10,000 gallons of cooling water (5-20 ppm of available chlorine dioxide) every week.
- 2) Depending on the degree and type of contamination, addition frequency may be reduce to every 2-3 weeks when contamination is under control.

10/22

TO INHIBIT THE GROWTH OF SLIME AND ODOR CAUSING BACTERIA IN WATER BASED CUTTING OILS

- 1) *Batch Method* - Add 80 oz. of Carnebon 200[®] per thousand gallons to fresh system and repeat weekly or on first indication of increased bacterial contamination (odor, slime, bacterial count). Alkaline systems may require higher concentration of Carnebon 200[®].
- 2) *Continuous Method* - Proportion in 5 gallons of Carnebon 200[®] per million gallons per day used in the system. Alkaline systems may require higher concentration.
- 3) *Badly Contaminated Systems* - Slug dose system with 25 gallons of Carnebon 200[®] per million gallons of cutting oil. Then start the continuous procedure described above.

Adjust quantities in any of the above systems to compensate for levels of contamination, pH, type of contamination etc., as necessary.

AS A SLIMICIDE IN PAPER MILLS TO PREVENT SLIME, TAR SPOTS, AND PITCH SPOTS IN WHITE WATER SYSTEMS

By maintaining a chlorine dioxide concentration in the white water, the microorganisms cannot produce the nodules which result in slime.

- 1) If the pH of the white water is below 7.0, add 11¼ gallons of Carnebon 200[®] per hundred tons of paper produced.
- 2) If the pH of the white water is above 7.0, then add ½ gallon of 5% sodium hypochlorite as an activator with each 11 ¼ gallons of Carnebon 200[®].

Continuous proportioning of the Carnebon 200[®] feed is recommended for best results. In many cases, the amount can be reduced after the system is clean.

TO PREVENT CORROSION AND SLIME BACTERIA IN OIL WELLS DURING SECONDARY RECOVERY OPERATIONS

- 1) Prepare a working solution of 5,000 ppm of available chlorine dioxide by diluting each gallon of Carnebon 200[®] used to 4 gallons of solution with the injection water.
- 2) Proportion 1 part of the above solution into each 150 parts of reinjected acidified (3.0 - 4.0 pH) water.
- 3) Monitor microbial content of the water and increase or decrease the addition rate of the working solution as necessary.

FOR CONTROL OF MOLLUSKS IN ONCE THROUGH WATER COOLING SYSTEMS AND INTAKES

- 1) Add 10 gallons of Carnebon 200[®] to 100 gallons of water and add 1 lb. of Activator-C (or 6.9 lbs. of Activator K) to the solution with mild stirring for 15 minutes. This produces an activated solution containing 2,000 ppm available chlorine dioxide. (Use respirator approved for chlorine dioxide).
- 2) As an alternate activation method, reduce the pH of the above solution to 3.0 with a mineral or organic acid and allow to slowly stir for ½ hour before use.

SLUG DOSE: Add between 2.5 gallons and 12.5 gallons of the above solution per 1,000 gallons of water (5-25 ppm of available chlorine dioxide).

CONTINUOUS DOSE: Add between 0.125 gallons and 1 gallon of the above solution per 1,000 gallons of water (0.25 to 2.0 ppm of available chlorine dioxide).

MUSHROOM FACILITIES

Carnebon 200[®] can be used in mushroom facilities such as mushroom production, spawn production, mushroom processing and cannery operations:

- As a food-contact surface sanitizer.
- To disinfect non-food contact surfaces.
- To control mold and mildew on environmental surfaces.

Irrigation Water

Dilute 1 part Carnebon 200® with 400 parts water to obtain optimum deodorizing and whitening effects in irrigation water. This solution is designed to be added to irrigation water to control odors and whiten product growing in plastic or stainless steel vessels.

As An Area Deodorizer

Dilute 1 part Carnebon 200® with 7 parts water to obtain optimum deodorizing effects. To eliminate gaseous malodors, spray or mist until odor disappears. Three (3) seconds of spraying or fogging is need for each 1500 cu. feet. Where a fogging is used in very large areas, set device to run 1-2 minutes each hour or less as area is cleared of malodors. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide.

For Deodorizing Sludge Waste Pile or Land Areas

Spray until surface is well saturated. Repeat daily or upon reoccurrence of odor.

Specific Applications

Use Carnebon 200® As a Terminal Sanitizing Rinse for Stainless Steel Tanks, Transfer Lines, On-line Equipment, Picking Baskets, Picking Utensils and Other Food Contact Surfaces

- 1) All gross food particles and soil should be removed prior to sanitizing by use of a pre-flush, pre-scrape or pre-soak treatment.
- 2) Clean picking baskets, line equipment or other surfaces thoroughly using a suitable detergent and rinse with water before sanitizing.
- 3) Follow instructions for **Food-Contact Surface Sanitizing Solution** as described above.
- 4) **Application:** Flush picking baskets, line equipment or other food-contact surfaces with the sanitizing solution making sure surface area is thoroughly wet for at least one (1) minute. After sanitizing, drain baskets or equipment and allow to air dry. Treat after each use or production run. Discard solution after each use.
- 5) **Optional Activated Use-Solution (Oxychlor e-generator)** An activated use-solution can also be prepared electrolytically by adding Carnebon 200 directly to the Oxychlor e-generator. The activated use-solution prepared by the Oxychlor e-generator must contain between 50-100 ppm of total available chlorine dioxide. For proper operation of the Oxychlor e-generator, consult the Oxychlor e-generator system manual or your IDI representative.

To Disinfect Walls, Ceilings and Floors

- 1) Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
- 2) Follow the directions for **Disinfectant Use-Solution** as described above.
- 3) **Application:** Spray disinfectant solution onto surface using a suitable spraying device and making sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.

To Control Mold and Slime Forming Bacteria on Walls, Floors, Ceilings, and Post-Crop Mushroom Growing Surfaces

- 1) Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
- 2) **Preparation of Use-Solution:** Place 1 part Carnebon 200® per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail or drum and dilute with clean, potable water.
- 3) **Application:** Drench, spray or fog solution on walls, floors, ceilings and post-crop mushroom growing surfaces using a suitable watering, spraying or fogging device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying or fogging, the area should be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.
- 4) Repeat application as needed.

POULTRY

To Treat Poultry Chiller Water

A) Carnebon 200® Plus Chlorine

In order to control the microorganism population in poultry chiller water, target the addition of available chlorine dioxide at 20-40 ppm level so that a residual of 0.5-3 ppm is measured in the exiting chilled water.

This is easily accomplished by activating Carnebon 200®, a mixture of oxychlorine species capable of generating 95%+ of chlorine dioxide, with chlorine which is already available in all poultry chiller water systems. The feed rates of the various streams is set forth below for the reactants, chlorine and Carnebon 200®.

PPM ClO ₂	CARNEBON 200® FEED RATE	Cl ₂ FEED RATE LBS./GAL	Cl ₂ PPM
20	1.0 gal/1000 gal H ₂ O	0.0083	10
30	1.5 gal/1000 gal H ₂ O	0.01245	15
40	2.0 gal/1000 gal H ₂ O	0.0166	20

B) Carnebon 200® Plus Acid

This antimicrobial agent may be used as a component of (1) a carcass spray or dip solution prior to immersion of the carcass in a prechiller or chiller tank or (2) in a prechiller or chiller solution.

- 1) When used as a carcass spray or dip solution, dilute 1 part of Carnebon 200® to 28 parts with water. The solution is then acidified to a pH between 2.5 and 2.9 with an acid selected from the following acids: phosphoric, citric, acetic, hydrochloric, lactic, malic or sulfuric.
- 2) When used in a prechiller or chiller tank, Carnebon 200® is diluted 1:280 (i.e. 2.5 gallons of Carnebon 200® diluted to 700 gallons with water). This solution is activated by addition of an acid such as phosphoric, citric, acetic, hydrochloric, lactic, malic or sulfuric to a pH of between 2.8 to 3.2.
- 3) As an alternate activation method, the Oxychlor e system, may be used to generate 700 ppm available chlorine dioxide solution for a carcass spray or dip solution or 70 ppm for use in the prechiller. For proper operation of the Oxychlor e-generator, consult the Oxychlor e-generator manual or your IDI representative.

To Control Bacteria, Taste and Odor in the Water Supply System

- 1) If the water supply is badly fouled with biofilm, then add 5 ppm of available chlorine dioxide to the water supply by adding 1 part of Carnebon 200® to each 4000 parts of poultry drinking water.
- 2) After 24 hours, the addition rate can be reduced to 1 ppm of available chlorine dioxide by adding 1 part Carnebon 200® to each 20,000 parts of poultry drinking or cooling comfort water.
- 3) If the microbiological content of the water is eliminated by this rate of addition, the concentration of available chlorine dioxide can be reduced to 0.5 ppm (1 part of Carnebon 200® per 40,000 parts of water); if the microbiological control is not adequate at 1 ppm available chlorine dioxide, then add 1.5 ppm of available chlorine dioxide to the poultry drinking or cooling comfort water (1 part of Carnebon 200® per 14,000 parts of water).

ANIMAL REARING AND CONFINEMENT FACILITIES**To Control Bacteria, Taste and Odor in the Water Supply System**

- 1) If the water supply is badly fouled with boil, then add 5 ppm of available chlorine dioxide to the water supply by adding 1 part Carnebon 200[®] to each 4,000 parts animal drinking and animal cooling / comfort water.
- 2) After 24 hours, the addition rate can be reduced to 1 ppm of available chlorine dioxide by adding 1 part of Carnebon 200[®] to each 20,000 parts of animal drinking and animal cooling / comfort water.
- 3) If the microbiological content of the water is eliminated by this rate of addition, the concentration of available chlorine dioxide can be reduced to 0.5 ppm (1 part of Carnebon 200[®] per 40,000 parts of water); if the microbiological control is not adequate at 1 ppm available chlorine dioxide, then add 1.5 ppm of available chlorine dioxide to the animal drinking and animal cooling / comfort water (1 part of Carnebon 200[®] per 14,000 parts of water).

20/22

To Control Bacteria and Odor in the Egg Room

- 1) Wash down the entire egg room with high pressure water containing 20 ppm of available chlorine dioxide (1 part Carnebon 200® diluted to 1,000 parts with water) to remove gross filth or heavy soil.
- 2) Spray the entire area for 5 minutes with a Tri-Jet Fogmaster (or equivalent) with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200® diluted to 20 parts with water), being sure to cover walls, ceiling, floors, work tables and benches. Allow to dry for 1 hour or if possible overnight before resuming operations.

The washing and fogging operations should be conducted once per week (or more frequently in cases of heavy contamination during operations).

- 3) If it is necessary to clean the floors by mopping, then use 390 ppm of available chlorine dioxide (1 part Carnebon 200® per 50 parts water). Allow to dry on the floor.
- 4) A shoe or boot bath of 1,000 ppm of available chlorine dioxide (1 part Carnebon 200® per 20 parts water) is placed at the entrance to the egg room. Doors to the room should be kept closed at all times.
- 5) A glove dip, or rinse tank or basin, containing 50 ppm of available chlorine dioxide (1 part Carnebon 200® per 400 parts water) is used on entering and exiting the room.

Both the shoe and boot bath and glove dip should be replaced daily (sooner if traffic is heavy).

- 6) Humidification water is treated with 40 ppm of available chlorine dioxide (1 part of Carnebon 200® per 500 parts water) to prevent the build-up and airborne spread of odor-causing microorganisms.
- 7) Provide 20 ppm of available chlorine dioxide (1 part Carnebon 200® per 1,000 parts water) to the water supply in the egg washing machine.

To Control Bacteria and Odor in the Hatching Area

- 1) As soon as chicks are separated from Hatch, remove all trash containers with eggshells, down, etc. from the hatching area.
- 2) Remove all poultry and feeds from premises, trucks, coops and crates.
- 3) Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
- 4) Empty all troughs, racks and other feeding and watering appliances.
- 5) Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 6) Spray or fog the entire area for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200® to 20 parts with water), using a Tri-Jet Fogmaster (or equivalent). Allow a 10 minute contact time.
- 7) Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
- 8) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.
- 9) All workers in this area should use a hand dip or rinse containing 50 ppm of available chlorine dioxide (1 part Carnebon 200® diluted to 400 parts with water).

To Control Odor and Bacteria when Separating Chicks in the Chick Room, Chick Grading Box and Sexing Room

- 1) Remove all poultry and feeds from premises, trucks, coops and crates.
- 2) Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
- 3) Empty all troughs, racks and other feeding and watering appliances.
- 4) Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5) Spray or fog the entire area for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] to 20 parts with water), using a Tri-Jet Fogmaster (or equivalent). Allow a 10 minute contact time.
- 6) Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
- 7) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.
- 8) All workers in this area should use a hand dip or rinse containing 50 ppm of available chlorine dioxide (1 part Carnebon 200[®] diluted to 400 parts with water).
- 9) After use, wash area with high-pressure water to remove gross filth and soil.
- 10) Use a spray bottle containing a solution of 1,000 ppm of available chlorine dioxide (1 part Carnebon 200[®] diluted to 20 parts with water), on hands, wire mesh and in empty chick boxes to control contamination and odors from litter.
- 11) To clean the floor by mopping daily, use a solution containing 390 ppm of available chlorine dioxide (1 part Carnebon 200[®] per 50 parts water). Allow to air dry.

To Control Bacteria and Odor in the Incubator Room

- 1) The area is sprayed or fogged at least once per week for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] diluted to 20 parts with water), after removing gross filth or soil with a high pressure water wash. Wet all surfaces and allow to dry.
- 2) The floor should be mopped daily with a solution containing 390 ppm of available chlorine dioxide (1 part of Carnebon 200[®] diluted to 50 parts with water).
- 3) A shoe and boot bath containing 1,000 ppm of available chlorine dioxide (1 part Carnebon 200[®] diluted to 50 parts with water) should be placed at all entrances to the incubator room.
- 4) 20 ppm of available chlorine dioxide (1 part Carnebon 200[®] diluted to 1,000 parts with water) is added to water in the humidification system or the air filters are sprayed with a 100 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] diluted to 200 part with water) to reduce airborne bacterial contamination.
- 5) Each time the eggs are removed from the incubator, a prior glove dip at 50 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] diluted to 400 parts with water) is recommended, followed by a spray of 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] diluted to 20 parts with water) on the eggs from a spray bottle.
- 6) Where containers are used to discard bad eggs, 1 part of Carnebon 200[®] per 7 parts of water (3,125 ppm of available chlorine dioxide) will control obnoxious odors and bacterial contamination.

The doors to the area should be kept closed as much as possible to avoid airborne contamination.

22/22

To Prevent Airborne and Surface Contamination of the Hatchery from the Tray Washing Room and Loading Platform

- 1) Close all doors in the tray washing room to avoid contamination of other hatchery operations. Discard all chick downs, egg shells, and cast-off chicks into the trash barrels and transfer the covered containers to the loading platform for disposal.
- 2) Wash the trays, carriages and other working equipment in a tray washing machine with 300-500 psi water to remove gross filth and soil.
- 3) As a final rinse in the tray washing machine, use a solution containing 20 ppm of available chlorine dioxide (1 part of Carnebon 200[®] diluted to 1,000 part with water) in high pressure water. Allow the trays, carriers and other working equipment to air dry. The walls, floors and carrying stands must also be sanitized with the same solution. Allow the equipment to air dry. Hold the sanitized equipment in a closed area for reuse.
- 4) Entrance and exit from the tray washing room must be through a foot rinse containing a solution of 1,000 ppm of available chlorine dioxide (1 part Carnebon 200[®] diluted to 20 parts with water). The rinse must be at least ½ inch deep and should be changed daily unless traffic is heavy.
- 5) After use, the tray washing room is washed with high pressure water to remove gross filth and soil. It is then decontaminated by spraying or fogging with a solution containing 1,000 ppm of available chlorine dioxide (1 part of Carnebon 200[®] diluted to 20 parts with water) for 15 minutes and allowed to air dry. This treatment is repeated after each use of the tray wash room.
- 6) The Loading Platform is washed from time to time to remove gross filth and soil. The trash containers are washed after discarding the contents to remove gross filth and soil. They are then sprayed with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon 200[®] diluted to 20 parts with water) and stored.