



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 12, 2020

Christina M. Swick
Agent
International Dioxide, Inc.
40 Whitecap Drive
North Kingstown, RI 02852

Subject: Label Amendment: Emerging Viral Pathogens Claim
Product Name: Carnebon 200 2% Aqueous Stabilized Chlorine Dioxide
EPA Registration Number: 9150-3
Application Date: 03/26/2020
Decision Number: 561171

Dear Ms. Swick:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Because you have opted to add statements pertaining to emerging viral pathogens to your label as described in the August 19, 2016, Guidance to Registrants: Process For Making Claims Against Emerging Viral Pathogens Not On EPA-Registered Disinfectant Labels ("Guidance"), https://www.epa.gov/sites/production/files/2016-09/documents/emerging_viral_pathogen_program_guidance_final_8_19_16_001_0.pdf, you are subject to the following additional terms of registration:

1. You may make statements pertaining to emerging viral pathogens only through the following communications outlets: technical literature distributed exclusively to health care facilities, physicians, nurses and public health officials, "1-800" consumer information services, social media sites and company websites (non-label related). These statements shall not appear on marketed (final print) product labels.


2. Your statements pertaining to emerging viral pathogens must adhere to the format approved on the Agency-accepted master label.
 3. You may make statements pertaining to emerging viral pathogens only upon a disease outbreak that meets all the following criteria:
 - a. The causative organism must be a virus that causes an infectious disease that has appeared in a human or animal population in the U.S. for the first time, or that may have existed previously but is rapidly increasing in incidence or geographic range.
 - i. For human disease, the outbreak is listed in one of the following Centers for Disease Control (CDC) publications:
 - A. CDC Current Outbreak List for “U.S. Based Outbreaks” (www.cdc.gov/outbreaks),
 - B. CDC Current Outbreak List for “Outbreaks Affecting International Travelers” with an “Alert” or “Advisory” classification (www.cdc.gov/outbreaks) (also released through the CDC’s Health Alert Network (HAN) notification process)
 - C. Healthcare-Associated Infections (HAIs) Outbreaks and Patient Notifications page (www.cdc.gov/hai/outbreaks)
 - ii. For animal disease, the outbreak is identified as an infectious disease outbreak in animals within the U.S. on the World Organization for Animal Health (OIE) Weekly Disease Information page (www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/WI).
- A. The CDC or OIE has identified the taxonomy, including the viral family and/or species, of the pathogen and provides notice to the public of the identity of the emerging virus that is responsible for an infectious disease outbreak. Based on the taxonomy of the outbreak pathogen identified by the CDC or OEI, the pathogen's viral subgroup are large non-enveloped, and enveloped.
 - B. The virus can be transmitted via environmental surfaces (non-vector transmission), and environmental surface disinfection has been recommended by the CDC, OIE or EPA to control the spread of the pathogen.
4. You may begin communicating statements pertaining to emerging viral pathogens only upon CDC or OIE’s publication per term 3.a. of an outbreak of an emerging viral pathogen meeting all of the criteria of term 3. You must cease and remove all such non-label communications intended for consumers no later than 24 months after the original publication of the outbreak per term 3.a., unless the Agency issue written guidance to the contrary due to continued public health concerns. The emerging pathogen claim language may remain on the master label.

5. Terms from points 1 through 4 above shall become immediately void and ineffective if registration for use against Canine parvovirus ATCC VR-2017 is suspended or cancelled or no longer meets the criteria for a disinfectant claim (see EPA Product Performance Test Guideline 810.2200). In addition, terms B.1 through B.4 above shall become immediately void and ineffective upon your receipt of evidence of ineffectiveness against any pathogen in a less-resistant Spaulding category.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, you may contact the disinfectants list at disinfectantslist@epa.gov.

Sincerely,




Steven Snyderman, Acting Product Manager 33
Regulatory Management Branch 1
Antimicrobials Division (7510P)
Office of Pesticide Programs

Enclosure: stamped label

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL
TEXT IN {BRACES} ARE COMMENTS TO REVIEWER
Alternate Brand Name: Anthium™ BCD-200

Carnebon™ 200
2% AQUEOUS STABILIZED CHLORINE DIOXIDE

Active Ingredient:
Chlorine Dioxide ----- 2%
Other Ingredients----- 98%
Total: 100%



KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCION

See side panels for additional Precautionary Statements
[See [attached] pamphlet for Directions for Use and Applications]

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment.

If inhaled: Move person to fresh air. If person is not breathing call 911 or an ambulance then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.

For Transport & Medical Emergencies: CHEMTREC: (800) 424-9300 (outside the U.S. (703) 527-3887). [For Product Information: [(800) 477-6071 (outside the U.S. (401) 295-8800)]]].

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 9150-3
EPA EST. No. XXXXXX-YY-ZZZ

NET CONTENTS: _____ GAL.

Manufactured For:
INTERNATIONAL DIOXIDE, INC.
40 Whitecap Drive
North Kingstown, RI 02852

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL
TEXT IN {BRACES} ARE COMMENTS TO REVIEWER
Alternate Brand Name: Anthium™ BCD-200

[© 2017 International Dioxide, Inc. All rights reserved. Anthium is a trademark of International Dioxide, Inc. [or its affiliates.]]

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

CHEMICAL AND PHYSICAL HAZARDS

Chlorine dioxide is a strong oxidizing agent. Contamination with other materials such as acids, chlorine, organic chemicals, etc. may cause a chemical reaction resulting in evolution of chlorine dioxide and heat. Explosion and/or fire could result. Chlorine dioxide is a poisonous explosive gas. Keep all chemical and foreign materials away from this solution.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store with easily oxidizable materials, acids, reducers, and combustible material. Avoid heat or freezing conditions. Store upright and do not stack drums over two high on pallets or partially filled drums. Use of a drum pump is suggested. Keep drum tightly closed when not withdrawing liquid. In case of spills, dilute with large quantities of water. Do not allow liquid to dry because this could present a fire hazard. Store only in the original container and take care to prevent cross-contamination with other pesticides, fertilizers, food and feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

[[Containers equal to or less than 5 gallons:] Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

[[Containers over 5 gallons:] Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

EMERGENCY HANDLING: In case of contamination or decomposition, do not reseal container. Isolate in open, well-ventilated area. Flood with large volumes of water. Cool unopened drums in vicinity by water spray.

EMERGENCY HANDLING: In case of contamination or decomposition, do not reseal container. Isolate in open, well-ventilated area. Flood with large volumes of water. Cool unopened drums in vicinity by water spray.

WARRANTY: Seller expressly warrants that the product conforms to its chemical description. There are no warranties associated with the sale of the product either express or implied, including but not limited to the warranties of fitness for a particular purpose or use.

DIRECTIONS FOR USE

[† Not approved for use in California]

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[Carnebon™ 200 is a specially designed formulation of chlorine dioxide and is a very versatile biocide. It controls microbial contamination in animal research facilities, food-processing and industrial waters, pulp and papermaking processing waters and cutting oils. It also disinfects environmental surfaces in hospitals and institutions and sanitizes food-contact surfaces. Carnebon™ 200 is highly effective against mold and mildew.]

[Carnebon™ 200 delivers a non-corrosive disinfectant and cleaning performance in an economical concentrate.]

[Carnebon™ 200 meets AOAC efficacy standards for hospital disinfectants and food-contact surface sanitizing solutions.]

[Carnebon™ 200 can be used in federally inspected meat and poultry plants as both a disinfectant and food-contact surface sanitizer.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

[The efficacy of Carnebon™ 200 depends on the degree of activation. Unactivated Carnebon™ 200 effectively controls microbes in processing waters and mold and mildew. For disinfection and sanitization, Carnebon™ 200 must be activated. Read the activation instructions carefully prior to using Carnebon™ 200.]

[Carnebon™ 200 can be used to treat hard, non-porous surfaces and water systems in: hospitals, medical and dental offices, food processing facilities, bottling plants, breweries, meat-packing plants, poultry-processing plants, fish-processing plants, food storage areas, institutional kitchens, dairy and poultry farms and production facilities, mushroom production facilities, animal research facilities, agricultural storage facilities (including containers, trailers, rail cars, vessels and bins), animal transport vehicles and equipment, animal confinement and rearing facilities, animal handling facilities, egg processing plants, livestock facilities, hatcheries, hotels, business and office buildings, institutional facilities, public facilities.]

[Carnebon™ 200 is an effective disinfectant against the following bacteria at a 300 ppm activated use-solution of Carnebon™ 200 (~ 30 ppm free chlorine dioxide) in 10 minutes in the presence of 5% organic serum.

- Pseudomonas aeruginosa (Pseudomonas)
- Staphylococcus aureus (Staph)
- Salmonella enterica (Salmonella)]

[Carnebon™ 200 is tuberculocidal (effective against Mycobacterium bovis, BCG) at a 1200 ppm activated use-solution of Carnebon™ 200 (~200 ppm free chlorine dioxide) in 10 minutes at 20 deg C.]

[+] [Carnebon™ 200 is an effective virucide against the following viruses at an 800 ppm activated use-solution of Carnebon™ 200 (~100 ppm free chlorine dioxide) in 10 minutes, 15 minutes for Canine parvovirus ATCC VR-2017.

- HIV-1 (AIDS Virus) HTLV-IIIB*
- Canine parvovirus ATCC VR-2017*
- Rat coronavirus RCV-SDA-681*
- Mouse hepatitis virus MHV-A59*
- Minute virus of mice MVM-P*
- Parainfluenza virus, Type 1 ATCC VR-105 SENDAI/52*]

[Carnebon™ 200 is an effective sanitizer against Salmonella typhi at a 100-200 ppm activated use-solution of Carnebon™ 200 in 30 seconds.]

[+] [An unactivated use-solution of 1000 ppm of Carnebon™ 200 effectively controls mold and mildew in 60 seconds.]

[Preparation of Carnebon™ 200 Use-Solutions]

[Virucidal Use-Solution*]

Prepare an approximate 800 ppm use-solution of Carnebon™ 200 (100 ppm of free chlorine dioxide) by adding 1 part of Carnebon™ 200 into a clean, plastic pail and then add 5 parts of the 10% acid activator solution. The acid activator can be acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic, or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Allow 15

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

minutes for reaction time and for the activator to completely dissolve. Then dilute the activated solution with 20 parts of water.]

[†] [Tuberculocidal Use-Solution

Prepare an approximate 1200 ppm use-solution of Carnebon™ 200 (200 ppm of free chlorine dioxide) by adding 1 part of Carnebon™ 200 into a clean, plastic pail and then add 5 parts of a 10% acid activator solution. The acid activator can be acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic, or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Allow 60 minutes for reaction time and for the activator to completely dissolve. Then dilute the activated solution with 12 parts of water. Prepare in a well-ventilated area and avoid breathing any fumes that may be produced during activation.]

[Disinfectant Use-Solution

Prepare an activated 300 ppm use-solution of Carnebon™ 200 by using one of the three procedures described below.

1. Add 1 part Carnebon™ 200 to 64 parts water and then adjust the pH of the diluted Carnebon™ 200 to 2.6 with acetic, citric, phosphoric, sulfuric, hydrochloric or other equivalent acid. Please contact your IDI or authorized representative regarding equivalent acids. Prepare in a well-ventilated area and avoid breathing any fumes which may be produced during activation.
Alternatively to minimize worker handling, an automated system can also be utilized that will safely activate the concentrate of Carnebon™ 200 with any of the various acids listed to deliver the proper pH and safety dilute the material to the 300 ppm working solutions.
2. Add 2 fl. oz. of Carnebon™ 200 to one (1) gallon of water into a clean plastic pail and add 1.2 grams of Activator C or 8.6 grams of Activator K. Allow 15 minutes reaction time and for the activator to completely dissolve. Prepare in a well-ventilated area and avoid breathing any fumes, which may be produced during activation.
3. An activated 300 ppm use-solution of Carnebon™ 200 can also be prepared electrolytically by adding Carnebon™ 200 directly to the e-generator. For proper operation, of the e-generator, consult the e-generator manual or your IDI or authorized representative.]

[†] [Mold & Mildew Use-Solution

Prepare a 1000 ppm use-solution of Carnebon™ 200 by placing 1 part Carnebon™ 200 per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail or drum. Dilute with clean, potable water.]

[Food-Contact Surface Sanitizing Solution

Prepare a 200 ppm activated use-solution of Carnebon™ 200 by using one of the three procedures described below:

1. Add 1 part Carnebon™ 200 to 3 parts water and then activate by adding food-grade citric, phosphoric, acetic or other equivalent food-grade acid (of at least 99% purity) to a pH of 2.6. Please contact your IDI or authorized representative regarding equivalent acids. Agitate for 5 minutes and then allow to stand for 15 minutes. Then dilute 1 part of the activated solution with 24 parts of water.
Alternatively to minimize worker handling, an automated system can also be utilized that will safely activate the concentrate of Carnebon™ 200 with any of the various acids listed to deliver the proper

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

pH and safely dilute the material to the 200 ppm working solutions.

2. Add 1 gallon of Carnebon™ 200 to 20 gallons of water followed by 12 grams of Activator K. Allow to stand for 15 minutes after agitation for 5 minutes and then dilute 1 part with 4 parts of water.
3. An electrolytically activated use-solution can be prepared by adding Carnebon™ 200 directly to the e-generator. The activated use-solution prepared by the e-generator must contain between 50-100 ppm of an activated use-solution of Carnebon™ 200. For proper operation of the e-generator, consult the e-generator system manual or your IDI or authorized representative.]

[Non-Food Contact Surface Sanitizing Solution

1. Prior to sanitization, remove all gross food particles and soil.
2. Prepare a 30 ppm activated use-solution of Carnebon™ 200:
3. To 10 gallons of water add 1.9 oz. of Carnebon™ 200 and 0.38 oz. of sodium hypochlorite (6%) for a nominal 30 ppm solution. Adjust the pH to between 2.5 and 3.5 with acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic or other suitable acid. Hold the solution for 15 minutes before applying. The efficiency of the conversion can be affected by the quality of the water. Conditions may be adjusted to accommodate the quality of the water.]

[Prepare a Non-Food Contact Door Foam Sanitizer Solution:

Using a dilution and delivery device, add Carnebon™ 200 at the rate of 1.9 oz. per 10 gallons of water, add sodium hypochlorite (6%) at the rate of 0.38 oz. per 10 gallons of water, followed by a foaming acidic activator at a rate of 0.38 oz. per 10 gallons of water. Hold the solution for 15 minutes before spraying.]

[APPLICATION INSTRUCTIONS:]

[FOOD PROCESSING PLANTS, FOOD-HANDLING ESTABLISHMENTS AND RESTAURANTS

Carnebon™ 200 can be used to:

- To control microbial contamination, slime and odor in food processing waters.
- To sanitize food processing equipment and surfaces in food processing and food-handling establishments.
- To sanitize food-contact surfaces and utensils in food-handling establishments.
- To disinfectant non-food contact surfaces in food-processing plants, food handling establishments and restaurants.
- For use as a terminal food-contact surface sanitizer rinse conforming to 40 CFR 180.940 (b) and (c) Food Contact Surface Sanitizing Solutions.]

[Specific Applications]

[Use Carnebon™ 200 in Food Processing Plants to Control the Build-Up of Odor and Slime Forming Bacteria in Stainless Steel Transfer Lines and On-Line Equipment Such as Hydrocoolers, Pasteurizers and the Like Overnight and Over Weekends.

1. Clean equipment or line thoroughly using a suitable detergent followed by a clean, potable water rinse before treatment.
2. Preparation and application of solution: For each 1000 parts of solution in lines and/or equipment, add 1 part of Carnebon™ 200 (20 ppm available chlorine dioxide) to potable make-up water. Mix and fill lines and equipment overnight. Drain and allow to air dry just prior to next run start-up.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

[Use Carnebon™ 200 as a Terminal Sanitizing Rinse for Stainless Steel Tanks, Transfer Lines, On-line Equipment, Recirculating and Clean-in-Place (CIP) systems, Food-contact surfaces and similar surfaces, such as tables, trays, bins, etc., utensils and Food-Processing Equipment in Poultry, Meat, Fish & Meat Processing Plants, Dairies, Bottling Plants, Restaurants, Canneries and Breweries

1. Prior to sanitization, remove all gross food particles and soil by use of a pre-flush, pre-scrape or pre-soak treatment. Then clean all lines, tanks, or surfaces with a suitable detergent followed by a potable water rinse.
2. Prepare the **Food-Contact Surface Sanitizing Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Fill, immerse, circulate, wipe or spray the target surface with the sanitizing solution making sure the surface area is thoroughly wet for at least one minute. Hard to reach in-place equipment, pipes, closed vessels, etc., must be filled with the sanitizing solution to ensure contact of all surfaces. Use suitable protective breathing apparatus when spraying the solution on external equipment.
4. Allow the sanitizing solution to drain from all treated surfaces and air dry. Do not rinse treated surface.
5. The above solution may not be reused for sanitizing but may be diluted to 1:5 with water and used for cleaning of walls, floors and drains of the plant.]

[+] [Use Carnebon™ 200 To Disinfect Non-Food Contact Surfaces (Walls, Ceilings, Drains and Floors) in Food Processing Plants and Food-Handling Establishments.

1. Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
2. Prepare the **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Apply the disinfectant use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge or sprayer, or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. Preclean surfaces if heavily soiled.
4. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface rub with a brush, sponge or cloth. Do not breathe spray. Make sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.
5. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[+] [Use Carnebon™ 200 to Control Mold & Mildew and Slime Forming Bacteria on Non-Food Contact Surfaces (Floors, Walls, Ceilings and Drains) in Food-Processing Plants and Food-Handling Establishments

1. Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
2. Follow the directions for **Mold & Mildew Use-Solution**: as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application: Drench or spray solution on walls, floors, ceilings and surfaces using a suitable watering or spraying device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying, the area must be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.

4. Repeat application as needed.]

[This product may be used in accordance with FDA regulations in Title 21 CFR 173.300 and 21 CFR 173.325; Secondary direct food additives permitted in food for human consumption.]

[Use Carnebon™ 200 in Food Processing Plants to Control Odor and Slime Forming Bacteria in Cooling and Warming Waters Such as Canning Retort and Pasteurizer Cooling Water Used to Decrease or Increase Packaged Product Temperature By Immersion in or by Spraying with the Treated Process Waters

1. All tanks, tunnels, conveyor chains, heat exchangers, heat exchange towers, lines, spray bars, and nozzles must be thoroughly cleaned when possible, and completely rinsed using clean, potable water prior to treatment.
2. Preparation and Application of Use-Solution: Water systems, including the cooling or warming tanks or spray systems, towers, lines and all water containing parts of the system may be batch loaded at start-up with one part Carnebon™ 200 per 4,000 parts potable water (5 ppm available chlorine dioxide). To maintain the 5 ppm available chlorine dioxide in the water system, a timed or electronically controlled chemical feed pump or injector system can be used for additions to the system or for treating the make-up water. Make up new Carnebon™ 200 solutions daily.
3. Preparation and Application of Optional Activated Use-Solution (acid activation): If heavy use of cooling or warming water or introduction of additional bacteria loads is expected, or if slime buildup is heavy, an additional activation step may be used in preparation of the use-solution. Prepare the activated use-solution in a well-ventilated area and avoid breathing any fumes which may be produced while crystals are dissolving. For each one thousand (1,000) gallons of system water to be treated, measure one quart (0.95 liters) of Carnebon™ 200 and pour into a clean plastic container, pail or drum. To this Carnebon™ 200 amount, add food grade citric acid of no less than 99% purity, at the rate of 3.3 oz. (95 grams or 0.2 lbs.) of crystals per quart (0.95 liters) of Carnebon™ 200. Allow five (5) minutes reaction time for crystals to dissolve. Cooling or warming water systems may be batch loaded at start up using one quart (0.95 liters) of the activated solution per one thousand (1,000) gallons of potable water (5 ppm available chlorine dioxide). Batch or timed additions of the activated solution can be made or an electronically controlled chemical feed pump or injector system can be used for additions of the activated solution to the process water to maintain 5 ppm available chlorine dioxide. Make up new Carnebon™ 200 solutions daily.
4. Preparation and Application of Optional Activated Use-Solution (e-generator): An activated use-solution can also be prepared electrolytically by adding Carnebon™ 200 directly to the e-generator. Add the activated use-solution prepared by the e-generator to water systems, including cooling or warming tanks, or spray systems, tower lines and to all water containing parts of the system. Batch load these systems at start-up and maintain a concentration of 5 ppm available chlorine dioxide in the system. For proper operation of the e-generator, consult the e-generator system manual or your International Dioxide representative.

Note: Chemical feed pumps and injectors must be chlorine resistant for best operation. Confirm levels of available chlorine dioxide by using a chlorine dioxide test kit.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

[Use Carnebon™ 200 to Control the Buildup of Odor and Slime Forming Bacteria in Process Waters for Vegetable Rinses and Associated Tanks, Flumes and Lines.

1. Clean all tanks, flumes, and lines thoroughly with a suitable detergent and completely rinsed using clean, potable water prior to treatment.
2. Preparation and Application of Use-Solution: Chill tanks or vegetable rinse tanks may be batch loaded at start-up with 1 part Carnebon™ 200 per 4000 parts of potable water (5 ppm available chlorine dioxide). Treat make-up waters using a chemical feed pump or injector system and applied at the rate of 1 part Carnebon™ 200 per 4,000 parts potable water. Make up new Carnebon™ 200 solutions daily.
3. Preparation and Application of Optional Activated Use-Solution: If heavy use of rinse water is expected, or if slime buildup is extreme, an additional activation step may be used in preparation of solution. Prepare the activated use-solution in a well-ventilated area and avoid breathing any fumes which may be produced while crystals are dissolving. For each 10 gallons of rinse water to be used, measure 1/3 fl. oz. (10 ml) of Carnebon™ 200 and pour into a clean, plastic container containing 1 gallon of water. Activate this solution by:
 - a. Adding 0.002 grams of Activator-C or
 - b. Adding 2.2 grams of Activator K or
 - c. Adjusting the pH to 2.6 with acetic acid, citric acid, phosphoric acid, sulfuric acid or hydrochloric acid.

Allow this solution to stand for 15 minutes and then add to 9 gallons of water to give 5 ppm available chlorine dioxide. Chill tanks or vegetable rinse tanks may be batch loaded at start-up with the activated Carnebon™ 200 solution 1/3 fl. oz. (10 ml) per ten (10) gallons of potable water (5 ppm available chlorine dioxide). Treat make-up waters using a chemical feed pump. In order to ensure accurate delivery, a 1 to 10 dilution of the active concentration must be made and the feed rate of 3 1/3 fl. oz. of activated Carnebon™ 200 solution per ten (10) gallons must be maintained. Make up fresh Carnebon™ 200 solutions daily.

4. Preparation and Application of Optional Activated Use-Solution (e-generator): An activated use-solution can also be prepared electrolytically by adding Carnebon™ 200 directly to the e-generator. Add the activated use-solution prepared by the e-generator to chill tanks or vegetable rinse tanks. Batch load these systems at start-up and maintain a concentration of 5 ppm available chlorine dioxide in the system. For proper operation of the e-generator, consult the e-generator system manual or your IDI representative.

Note: Chemical feed pumps and injectors must be chlorine resistant for best operation. Confirm levels of available chlorine dioxide by using a chlorine dioxide test kit.]

[Use Carnebon™ 200 as a Non-Food Contact Door Foam Solution

1. Prepare **Non-Food Contact Door Foam Sanitizing Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
2. Foam the generated solution on the surface to a depth of 1 inch minimum or onto the floor in the doorway to achieve complete wetting of equipment wheels.]

[Use of Carnebon™ 200 to Sanitize Non-Food Contact Surfaces (Walls, Ceilings, Drains and Floors) in Food Processing Plants and Food-Handling Establishments.

1. Before sanitizing, all gross filth must be removed from areas to be sanitized and thoroughly cleaned with a suitable detergent.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

2. Prepare the **Non-Food Contact Sanitizer Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Apply the sanitizer use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge, or sprayer or immersion. Treated surfaces must remain wet for five (5) minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled surfaces, a pre-cleaning is recommended.
4. For sprayer applications, use a coarse spray device. Spray 6 – 8 inches from the surface; rub with a brush, sponge or cloth. Do not breathe spray. Make sure that the area is thoroughly wet for at least five (5) minutes. Active solutions may be irritating if inhaled; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.
5. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[TRANSPORT VEHICLES]

[+] [To Disinfect Hard, Non-porous Surfaces in Vehicles Including Animal Transport Vehicles, Rail Cars, Trailers and Vessels. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.

1. Prior to application of Carnebon™ 200 clean all vehicles with high-pressure water and a suitable detergent.
2. Follow directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Then apply the disinfectant use-solution to all surfaces to be treated. All treated surfaces must remain wet for at least 10 minutes.]

[SEAFOOD USE]

[+] [Use Carnebon™ 200 as a Bacteriostat for Treating Ice Used for Icing Fish in the Round

Carnebon™ 200 may be batch loaded or metered into makeup water used to produce ice for icing fish in the round. Prepare a non-activated working solution containing 20 ppm of available chlorine dioxide by adding 1 part of Carnebon™ 200 to 1000 parts of water.]

[HOSPITALS, INSTITUTIONS, MEDICAL AND †DENTAL CLINICS, and VETERINARY CLINICS.

Note: The e-generator has not been tested against *Pseudomonas aeruginosa*. The e-generator is not approved for use in hospitals, laboratories, morgues, and other institutions.

[+] This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

[+] Carnebon™ 200 can be used to:

- To disinfectant environmental surfaces.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

- To control mold and mildew on environmental surfaces.
- To control animal viruses on environmental surfaces.
- To control odor and slime forming bacteria.]

[Specific Applications]

[†] [To Disinfect Walls, Ceilings and Floors and other Environmental Surfaces in Hospitals, Institutions, Veterinary Clinics, and Animal Research Facilities

1. Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
2. Prepare the **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Apply the disinfectant use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge or sprayer, or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled surfaces, a pre-cleaning is recommended.
4. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface rub with a brush, sponge or cloth. Do not breathe spray. Make sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.
5. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OF SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS that involve healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type I (HIV-I) (associated with AIDS). Carnebon™ 200 destroys HIV-1 (AIDS Virus) HTLV-IIIB on precleaned environmental surfaces/objects previously soiled with blood or other body fluids in ten minutes contact

Personal Protection: The worker must wear protective equipment such as disposable latex or rubber gloves, gowns, masks and eye protection to prevent contamination from items soiled with blood or body fluids.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of Carnebon™ 200

Contact Time: Allow Carnebon™ 200 to contact treated items for 10 minutes to kill HIV-1. This time may not control other common types of viruses and bacteria.

Disposal of Infectious Material: Any blood and other body fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.]

[†] [To Control Mold & Mildew and Slime Forming Bacteria on Walls, Floors, Ceilings, and Surfaces and other Environmental Surfaces

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

1. Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
2. Follow the directions for **Mold & Mildew Control Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application: Drench or spray solution on walls, floors, ceilings and surfaces using a suitable watering or spraying device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying, the area must be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.
4. Repeat application as needed.]

[To Disinfect Non-Porous, Hard Surfaces Such as Tile Floors, Walls and Ceilings and Stainless Steel Cold Rooms and Walk-In Incubators

1. Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application of Activated Disinfection-Solution: Activated solutions may be sprayed, mopped or sponged onto surfaces to be disinfected. All surfaces must be thoroughly wetted for at least ten (10) minutes. When spraying disinfectant solutions, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[To Sanitize Non-Porous, Non-Food Contact Hard Surfaces Such as Glazed Tile Floors, Walls and Ceilings and Stainless Steel Cold Rooms and Walk-In Incubators

1. Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to sanitizing.
2. Follow the directions for **Non-Food Contact Sanitizer Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application of Activated Sanitizer-Solution: Activated solutions may be sprayed, mopped or sponged onto surfaces to be sanitized. All surfaces must be thoroughly wetted for at least five (5) minutes. When spraying disinfectant solutions, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[†] [For Use in Dental Offices and Laboratories as a Dental Pumice Disinfectant

1. Prepare solution in a well-ventilated area. To make one (1) liter of solution, pour 1.0 fl oz (approximately 25 mls.) of Carnebon™ 200 into a clean glass or plastic container. To this, add 2 ½ grams (1/2 teaspoon) of citric acid crystals (included) and mix slightly, allowing 5 minutes reaction time and for crystals to dissolve completely. Avoid breathing any fumes which may be produced during activation. Once solution has yellowed, dilute to one (1) liter with clean potable water, for a working solution of 500 ppm activated ClO₂.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

2. To apply: The working solution can be conveniently contained in a one (1) liter plastic “squeeze” bottle for up to five days. Apply to dry pumice powder exactly as water to produce the pumice slurry. Apply additional working solution as needed to reconstitute dried out slurry to appropriate viscosity. Make fresh Carnebon™ 200 solutions, preferably on Monday and discarded on Friday or 5 days after preparation.]

[As a Virucide* to Kill Animal Viruses (Rat Coronavirus RCV-SDA-681, Mouse Hepatitis Virus MHV-A59, Minute Virus of Mice MVM-P and Canine Parvovirus ATCC VR-2017) Parainfluenza Virus Type 1 ATCC VR-105 SENDAI/52, HIV-1 HTLV-IIIB) on Non-Porous, Hard Surfaces Such as Tile Floors, Walls and Ceilings and Stainless Steel Cold Rooms and Walk-In Incubators.

1. Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
2. Follow the directions for **Virucidal Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application of Activated Use-Solution: Activated solutions may be sprayed, mopped or sponged onto surfaces to be treated. All surfaces must be thoroughly wetted for at least ten (10) minutes (15 minutes contact time for canine parvovirus ATCC VR-2017). When spraying the virucidal solution, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[To Disinfect Bench Tops, Biological Hoods, Incubators, Stainless Steel Equipment and Instruments.

1. Clean all surfaces thoroughly with a suitable detergent and rinse with water prior to disinfection.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application of Activated Use-Solution: Activated solutions may be squirted directly onto surfaces from a plastic squeeze bottle or may be used as a soak solution. All contact surfaces must be thoroughly wetted for at least ten (10) minutes. Allow to air dry. Activated solutions of Carnebon™ 200 stored in plastic squirt bottles, may be held up to one (1) week before replacement with fresh solution. Soak solutions of Carnebon™ 200 must be changed daily.]

[+] [To Disinfect Surfaces of Water Baths and Tubs

1. Prior to disinfection, thoroughly clean the bath or tub with a suitable detergent and rinse with clean water.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. To apply: Turn circulating motor on and allow the water to circulate for at least (10) minutes. Drain tub completely. After the draining is finished, tub is ready for use.]

[To Disinfect Water Bath Incubators

1. Prior to disinfection, thoroughly clean the reservoir with a suitable detergent and rinse with clean water.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

3. To apply: Pour the activated solution into water bath reservoir and allow to stand one (1) hour at room temperature. Drain reservoir and fill with fresh water.]

[+] [To Control Odor and Slime Forming Bacteria in Water Bath Incubators

1. When using Carnebon™ 200 in water bath incubators, always begin with a freshly cleaned and disinfected reservoir.
2. Application: Fill water bath with clean, potable water to near capacity. Add 1 part Carnebon™ 200 for each 400 parts water (50 ppm available chlorine dioxide). When water becomes cloudy, discard water and repeat procedure.]

[+] [To Control Odors Resulting from the Sterilization of Spent Biologicals in Steam Autoclaves

1. To reduce autoclave odors of used biologicals, spray or pour Carnebon™ 200 directly into the stainless steel autoclave buckets.
2. Preparation of Use-Solution: Place 1 part Carnebon™ 200 per 20 parts working solutions (1,000 ppm available chlorine dioxide) into a clean glass or plastic container and mix
3. Application: Spray or pour Carnebon™ 200 solution into or onto the autoclave buckets just prior to autoclaving.]

[+] [To Deodorize Animal Holding Rooms, Sick Rooms, Morgues and Work Rooms

1. Rooms to be deodorized must be in a clean condition prior to Carnebon™ 200 application.
2. Preparation of Use-Solution: Place 1 part Carnebon™ 200 per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean glass or plastic container.
3. Application: Spray solution using a suitable spraying device onto walls, ceilings and floors, lightly dampening all surfaces. Avoid breathing mist of solutions by using an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide. Allow to air dry, and then ventilate the area. Treat as required.]

[IN ANIMAL REARING & CONFINEMENT FACILITIES]

[+] [To Control the Build-up of Odor and Slime Forming Bacteria in Animal Confinement Areas

1. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, cases and other facilities and fixtures occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with clean water.
2. Preparation of Use-Solution: Place 1 part Carnebon™ 200 per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail.
3. Application: Using a commercial sprayer; saturate all surfaces with the Carnebon™ 200 solution. When spraying Carnebon™ 200 solutions, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide to avoid breathing mist.]

[To Disinfect Hard, Non-Porous Surfaces In Commercial Animal Confinement Facilities such as Poultry Houses, Swine Pens, Calf Barns and Kennels and for use in Laboratory Animal Breeding and Research Quarters for Controlling Cross-Contamination of Microorganisms Infectious to these Animals and Humans from Treated Surfaces

1. Remove all animals and feed from premises, vehicles, enclosures, coops and crates.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with water.
5. Prepare an activated 300-400 ppm use-solution of Carnebon™ 200: Add 2 ounces of Carnebon™ 200 to 1 gallon of water. Once the Carnebon™ 200 has been diluted, add 0.5-1.0 oz. "Biosolve™ AcidEdge™" per one gallon of solution OR 0.5-1.0 oz. "Biosolve™ Acidic Cleaner – low foaming formula" per gallon of solution. Prepare in a well ventilated area and avoid breathing any fumes which may be produced during activation
 - a. 2 ounces of Carnebon™ 200
 - b. 1 gallon of water
 - c. To the solution of step b. add 0.5 – 1.0 oz. "Biosolve™ AcidEdge™"
OR
0.5 – 1.0 oz. "Biosolve™ Acidic Cleaner – low foaming formula"Always add Carnebon™ 200 to water. Followed by acidic cleaner.
6. Application: Using a commercial sprayer, saturate all surfaces with the activated Carnebon™ 200 solution for a period of ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure.
7. After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate when solution has dried.
8. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.]

[To Sanitize Hard, Non-Porous Surfaces In Commercial Animal Confinement Facilities such as Poultry Houses, Swine Pens, Calf Barns and Kennels and for use in Laboratory Animal Breeding and Research Quarters for Controlling Cross-Contamination of Microorganisms Infectious to these Animals and Humans from Treated Surfaces

1. Remove all animals and feed from premises, vehicles, enclosures, coops and crates.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with water.
5. Prepare a 30 ppm activated use-solution of Carnebon™ 200: To 10 gallons of water add 1.9 ounces of Carnebon™ 200 and 0.38 oz. sodium hypochlorite (6%) for a nominal 30 ppm solution. Adjust pH to between 2.5 and 3.5 with acetic, citric, phosphoric, sulfuric, hydrochloric, glycolic or other suitable acid. Hold the solution for 15 minutes before applying. Alternatively, the following cleaners can be used instead of acid: add 0.5 – 1.0 oz "Biosolve™ AcidEdge™" per one gallon of solution OR 0.5-1.0 oz. "Biosolve™ Acidic Cleaner – low foaming formula" per gallon of solution. The efficiency of the conversion can be affected by the quality of water. Conditions may be adjusted to accommodate the quality of water.
6. Application: Using a commercial sprayer, saturate all surfaces with the activated Carnebon™ 200 solution for a period of five (5) minutes. Active solutions may be irritating when breathed;

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure.

7. After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate when solution has dried.
8. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.]

[As a Virucide* to Kill Animal Viruses (Rat Coronavirus RCV-SDA-681, Mouse Hepatitis Virus MHV-A59, Minute Virus of Mice MVM-P and Canine Parvovirus ATCC VR-2017) on Non-Porous, Hard Surfaces in Commercial Animal Confinement Facilities Such as Poultry Houses, Swine Pens, Calf Barns, and Kennels and in Laboratory Animal and Research Quarters

1. Remove all animals and feed from premises, vehicles, enclosures, coops and crates.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with water.
5. Follow the directions for **Virucidal Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
6. Application of Activated Use-Solution: Activated solutions may be sprayed, mopped or sponged onto surfaces to be treated. All surfaces must be thoroughly wetted for at least ten (10) minutes (15 minutes contact time for canine parvovirus ATCC VR-2017). When spraying virucidal use-solution, use an appropriate spraying device. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.
7. After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate when solution has dried.
8. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.]

[+] [To Control Mold & Mildew and Slime Forming Bacteria on Walls, Floors, Ceilings, Bins, Boxes, Pens, Barns, Kennels and other Animal Health Surfaces

1. Remove animals and feed from area to be treated
2. Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
3. Follow the directions for **Mold & Mildew Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
4. Application: Drench or spray solution on walls, floors, ceilings and surfaces using a suitable watering or spraying device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying, the area must be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces.
5. Follow treatment with a potable water rinse

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

6. Repeat application as needed.]

[+] [To Control Animal Odors on Pets and in Litter Boxes, Carpets and Concrete Floors

1. For litter boxes: Wash out litter boxes with suitable detergent and rinse with clean, potable water. Soak overnight in solution of one part Carnebon™ 200 per 32 parts of water (625 ppm available chlorine dioxide). Add litter, sprinkle surface liberally with Carnebon™ 200 solution.
2. For controlling odors in carpets: Add 1 part Carnebon™ 200 per 40 parts (500 ppm available chlorine dioxide) of rug shampoo mix. Shampoo carpets. Allow to air dry. **NOTE:** Carnebon™ 200 may bleach some carpets and fabrics, especially if applied on top of another chemical agent. Do not apply until a sample test has been tried and observed for at least 24 hours.
3. For concrete floors: Clean floor thoroughly using a suitable detergent; rinse with clean water. Prepare solution by adding 1 part of Carnebon™ 200 per 16 parts of water (1,250 ppm available chlorine dioxide). Mop or spray solution liberally onto floor. Allow to air dry.
4. For animal baths: Wash animal well with appropriate pet shampoo; rinse with clean water. Prepare solution by adding 1 part Carnebon™ 200 per 200 parts of water (100 ppm available chlorine dioxide). Rinse animal thoroughly with prepared solution. Allow to air dry. Avoid direct contact with animal's eyes, nose and ears.
5. For treating animal odors with high levels of ammonia: Wash area thoroughly with suitable detergent and rinse with clean water. Preparation of solution: For each 32 parts of solution place 1 part Carnebon™ 200 into a clean, plastic container. To this concentrate, add 1 tablespoon household bleach and allow to react for five (5) minutes. Dilute with 32 parts of clean, potable water. Apply by mopping or spraying solution liberally onto area. Allow to air dry. Additional applications may be necessary.]

[ANIMAL TRANSPORT VEHICLES]

[+] [To Disinfect Hard, Non-Porous Surfaces in Vehicles Including Animal Transport Vehicles, Rail Cars, Trailers and Vessels. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions.

1. Prior to application of Carnebon™ 200 clean all vehicles with high-pressure water and a suitable detergent.
2. Follow directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Then apply the disinfectant use-solution to all surfaces to be treated. All treated surfaces must remain wet for at least 10 minutes.]

[TREATMENT OF WATER STORAGE SYSTEMS AND POTABLE WATER]

[To Disinfect Water Storage Systems Aboard Aircraft, Trains, Buses, Boats, RV's, Off-Shore Oil Rigs, etc.

1. Prior to disinfection, clean tanks using a suitable detergent and thoroughly flush with clean, potable water.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

3. Pour activated solution into tank, filling the tank completely. Bleed air out of lines and allow to stand at least ten (10) minutes. Drain tank and lines and flush with potable water.]

[+] [To Control Build-Up of Slime and Odor Causing Bacteria and Enhance the Taste of Stored Potable Water

1. Prior to treatment of potable water, thoroughly clean and disinfect the water storage system to ensure a sanitary condition. Thoroughly rinse with clean, potable water.
2. Potable water must be treated at a rate of one part of Carnebon™ 200 per 4000 parts potable water (5 ppm available chlorine dioxide) and may be injected or batch treated.
3. Water storage tank must be sufficiently sealed to prevent outside contamination and direct sunlight.

Using a test kit, confirm the chlorine dioxide to be 5 ppm and check to see this level does not fall below 1 ppm.]

[+] [To Help Remove Off-Odors and Tastes from Municipal Well Waters

1. Inject Carnebon™ 200 into the incoming water main using a chemical proportioning pump, or injector, at a rate of 1 part Carnebon™ 200 per 20,000 parts water (1 ppm available chlorine dioxide).
2. Confirm pump or injector accuracy using an International Dioxide test kit and adjust accordingly.
3. Check the Carnebon™ 200 levels weekly.]

[INDUSTRIAL WATER SYSTEMS AND INDUSTRIAL BIOCIDES USE]

[+] [TO INHIBIT BACTERIAL SLIME FORMING BACTERIAL BUILDUP IN COMMERCIAL WATER FILTRATION SYSTEMS, SAND BEDS, GRAVEL BEDS, CHARCOAL FILTERS AND COOLING WATER SYSTEMS.

Filters:

1. Carefully back-flush filters with potable water, where possible, to remove any accumulated solid residue and contamination.
2. Fill system with potable water and adjust pH to 6.0 with citric acid, phosphoric acid, or acetic acid (vinegar) or equivalent.
3. Add 1 part of Carnebon™ 200 per 64 parts of (300 ppm of available chlorine dioxide) of filter system volume to the access hatch and circulate the system for 1 hour. Check the pH and bring back to 6.0 if it has drifted. Bring the available chlorine dioxide concentration back to 300 ppm.
4. Circulate the solution for 1 additional hour, discharge and then water wash for 30 minutes with potable water to remove the chlorine dioxide.]

[+] [For Enclosed and Recirculating Cooling Water Systems:

1. Add 2.5-10 gallons of Carnebon™ 200 per 10,000 gallons of cooling water (5-20 ppm of available chlorine dioxide) every week.
2. Depending on the degree and type of contamination, addition frequency may be reduced to every 2-3 weeks when contamination is under control.
3. For very high levels of microbial contamination of the cooling water, add an activated solution to the cooling water. First, dilute 1 part Carnebon™ 200 with 3 parts water and follow with acidification to a pH of 2.6 with phosphoric, citric or acetic acid. This forms an activated solution of 5,000 ppm available chlorine dioxide. Dilute the 5,000 ppm activated solution to the indicated feed solution

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

ppm in the table below by selecting the desired concentration in cooling water. Then add the feed solution to the cooling water at a rate of 1 part of feed solution to 13 parts of cooling water.

Desired Concentration – Available Chlorine Dioxide – Cooling Water	Feed Solution – Available Chlorine Dioxide (ppm)	Dilution – Activated Solution
5 ppm	70	1:70
10 ppm	140	1:35
15 ppm	210	1:23
20 ppm	280	1:17

- Alternatively, an activated use-solution can be prepared electrolytically by adding Carnebon™ 200 directly to the e-generator. Add the activated use-solution to the cooling water system so that a concentration of 5-20 ppm available chlorine dioxide is achieved. For proper operation of the e-generator, consult the e-generator manual or your International Dioxide, Inc. representative.]

[+] [TO PREVENT CORROSION AND SLIME BACTERIA IN OIL WELLS DURING SECONDARY RECOVERY OPERATIONS

- Prepare a working solution of 5,000 ppm of available chlorine dioxide by diluting each gallon of Carnebon™ 200 used to 4 gallons of solution with the injection water.
- Proportion 1 part of the above solution into each 150 parts of reinjected acidified (3.0 - 4.0 pH) water.
- Monitor microbial content of the water and increase or decrease the addition rate of the working solution as necessary.]

[+] [TO INHIBIT THE GROWTH OF SLIME AND ODOR CAUSING BACTERIA IN WATER BASED CUTTING OILS

- Batch Method** - Add 80 oz. of Carnebon™ 200 per thousand gallons to fresh system and repeat weekly or on first indication of increased bacterial contamination (odor, slime, bacterial count). Alkaline systems may require higher concentration of Carnebon™ 200.
- Continuous Method** - Proportion in 5 gallons of Carnebon™ 200 per million gallons per day used in the system. Alkaline systems may require higher concentration.
- Badly Contaminated Systems** - Slug dose system with 25 gallons of Carnebon™ 200 per million gallons of cutting oil. Then start the **Continuous Method**.

Adjust quantities in any of the above systems to compensate for levels of contamination, pH, type of contamination etc., as necessary.]

[+] [AS A SLIMICIDE IN PAPER MILLS TO PREVENT SLIME, TAR SPOTS, AND PITCH SPOTS IN WHITE WATER SYSTEMS

By maintaining a chlorine dioxide concentration in the white water, the microorganisms cannot produce the nodules which result in slime.

- If the pH of the white water is below 7.0, add 11¼ gallons of Carnebon™ 200 per hundred tons of paper produced.
- If the pH of the white water is above 7.0, then add ½ gallon of 5% sodium hypochlorite as an activator with each 11 ¼ gallons of Carnebon™ 200.

Continuous proportioning of the Carnebon™ 200 feed is recommended for best results. In many cases, the amount can be reduced after the system is clean.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

[+] [FOR CONTROL OF MOLLUSKS IN ONCE THROUGH WATER COOLING SYSTEMS AND INTAKES

1. Add 10 gallons of Carnebon™ 200 to 100 gallons of water and add 1 lb. of Activator-C (or 6.9 lbs. of Activator K) to the solution with mild stirring for 15 minutes. This produces an activated solution containing 2,000 ppm available chlorine dioxide. (Use respirator approved for chlorine dioxide).
2. As an alternate activation method, reduce the pH of the above solution to 3.0 with a mineral or organic acid and allow to slowly stir for ½ hour before use.

SLUG DOSE: Add between 2.5 gallons and 12.5 gallons of the above solution per 1,000 gallons of water (5-25 ppm of available chlorine dioxide).

CONTINUOUS DOSE: Add between 0.125 gallons and 1 gallon of the above solution per 1,000 gallons of water (0.25 to 2.0 ppm of available chlorine dioxide).]

[MUSHROOM FACILITIES

Carnebon™ 200 can be used in mushroom facilities such as mushroom production, spawn production, mushroom processing and cannery operations:

- As a food-contact surface sanitizer.
- To disinfect non-food contact surfaces.
- To sanitize non-food contact surfaces.
- To control mold and mildew on environmental surfaces.]

[+] [Irrigation Water

Dilute 1 part Carnebon™ 200 with 400 parts water to obtain optimum deodorizing and whitening effects in irrigation water. This solution is designed to be added to irrigation water to control odors and whiten product growing in plastic or stainless steel vessels.]

[+] [As an Area Deodorizer

Dilute 1 part Carnebon™ 200 with 7 parts water to obtain optimum deodorizing effects. To eliminate gaseous malodors, spray or mist until odor disappears. Three (3) seconds of spraying is need for each 1500 cu. feet. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide.]

[+] [For Deodorizing Sludge Waste Pile or Land Areas

Spray until surface is well saturated. Repeat daily or upon reoccurrence of odor.]

[Specific Applications]

[Use Carnebon™ 200 as a Terminal Sanitizing Rinse for Stainless Steel Tanks, Transfer Lines, On-Line equipment, Picking Baskets, Picking Utensils and Other Food Contact Surfaces

1. Remove all gross food particles and soil prior to sanitizing by use of a pre-flush, pre-scrape or pre-soak treatment.
2. Clean picking baskets, line equipment or other surfaces thoroughly using a suitable detergent and rinse with water before sanitizing.
3. Follow instructions for **Food-Contact Surface Sanitizing Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

4. Application: Flush picking baskets, line equipment or other food-contact surfaces with the sanitizing solution making sure surface area is thoroughly wet for at least one (1) minute. After sanitizing, drain baskets or equipment and allow to air dry. Treat after each use or production run. Discard solution after each use.]

[To Disinfect Walls, Ceilings and Floors

1. Before disinfection, all gross filth must be removed from areas to be disinfected and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
2. Follow the directions for **Disinfectant Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application: Spray disinfectant solution onto surface using a suitable spraying device and making sure that the area is thoroughly wet for at least ten (10) minutes. Active solutions may be irritating when breathed; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[To Sanitize Walls, Ceilings and Floors

1. Before sanitizing, all gross filth must be removed from areas to be sanitized and thoroughly cleaned with a suitable detergent followed by a clean, potable water rinse.
2. Follow the directions for **Sanitizer Use-Solution** as described in the Preparation of Carnebon™ 200 Use-Solutions section.
3. Application: Spray Sanitizer solution onto surface using a suitable spraying device and making sure that the area is thoroughly wet for at least five (5) minutes. Active solutions may be irritating if inhaled; therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

[+] [To Control Mold and Slime Forming Bacteria on Walls, Floors, Ceilings, and Post-Crop Mushroom Growing Surfaces

1. Before treatment, all soil and gross filth must be removed from areas to be treated and cleaned with detergent followed by a potable water rinse.
2. Preparation of Use-Solution: Place 1 part Carnebon™ 200 per 20 parts working solution (1,000 ppm available chlorine dioxide) into a clean, plastic pail or drum and dilute with clean, potable water.
3. Application: Drench or spray solution on walls, floors, ceilings and post-crop mushroom growing surfaces using a suitable watering or spraying device and making sure all surface areas are wet. During application, area must be closed as tightly as possible and sealed. After spraying, the area must be opened and aired for one (1) hour before repopulating. Avoid breathing solution mist by use of an applicable NIOSH/MSHA respirator appropriate for chlorine dioxide. Avoid contact with food or food-contact surfaces. Allow to air dry.
4. Repeat application as needed.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL
 TEXT IN {BRACES} ARE COMMENTS TO REVIEWER
 Alternate Brand Name: Anthium™ BCD-200

[POULTRY]

[+] [To Treat Poultry Chiller Water]

[+] [A) Carnebon™ 200 Plus Chlorine

In order to control the microorganism population in poultry chiller water, target the addition of available chlorine dioxide at 20-40 ppm level so that a residual of 0.5-3 ppm is measured in the exiting chilled water.

This is easily accomplished by activating Carnebon™ 200 a mixture of oxychlorine species capable of generating 95%+ of chlorine dioxide, with chlorine which is already available in all poultry chiller water systems. The feed rates of the various streams are set forth below for the reactants, chlorine and Carnebon™ 200

ClO ₂ PPM	CARNEBON™ 200 FEED RATE	Cl ₂ FEED RATE LBS. / GAL.	Cl ₂ PPM
20	1.0 gal/1000 gal H ₂ O	0.0083	10
30	1.5 gal/1000 gal H ₂ O	0.01245	15
40	2.0 gal/1000 gal H ₂ O	0.0166	20

[+] [B) Carnebon™ 200 Plus Acid

This antimicrobial agent may be used as a component of (1) a carcass spray or dip solution prior to immersion of the carcass in a rechiller or chiller tank or (2) in a prechiller or chiller solution.

1. When used as a carcass spray or dip solution, dilute 1 part of Carnebon™ 200 to 28 parts with water. The solution is then acidified to a pH between 2.5 and 2.9 with an acid selected from the following acids: phosphoric, citric, acetic, hydrochloric, lactic, malic or sulfuric.
2. When used in a prechiller or chiller tank, Carnebon™ 200 is diluted 1:280 (i.e. 2.5 gallons of Carnebon™ 200 diluted to 700 gallons with water). This solution is activated by addition of an acid such as phosphoric, citric, acetic, hydrochloric, lactic, malic or sulfuric to a pH of between 2.8 to 3.2.
3. As an alternate activation method, the e-generator system may be used to generate 700 ppm available chlorine dioxide solution for a carcass spray or dip solution or 70 ppm for use in the prechiller. For proper operation of the e-generator, consult the e-generator manual or your IDI representative.]

[+] [To Control Bacteria, Taste and Odor in the Water Supply System

1. If the water supply is badly fouled with biofilm, then add 5 ppm of available chlorine dioxide to the water supply by adding 1 part of Carnebon™ 200 to each 4000 parts of poultry drinking water.
2. After 24 hours, the addition rate can be reduced to 1 ppm of available chlorine dioxide by adding 1 part Carnebon™ 200 to each 20,000 parts of poultry drinking or cooling comfort water.
3. If the microbiological content of the water is eliminated by this rate of addition, the concentration of available chlorine dioxide can be reduced to 0.5 ppm (1 part of Carnebon™ 200 per 40,000 parts of water); if the microbiological control is not adequate at 1 ppm available chlorine dioxide, then add 1.5 ppm of available chlorine dioxide to the poultry drinking or cooling comfort water (1 part of Carnebon™ 200 per 14,000 parts of water).]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL
TEXT IN {BRACES} ARE COMMENTS TO REVIEWER
Alternate Brand Name: Anthium™ BCD-200

[ANIMAL REARING AND CONFINEMENT FACILITIES]

[To Disinfect Waterlines and Associated Fixtures in Animal Confinement Facilities such as Poultry Houses, Swine Pens, Calf Barns and Kennels When Animals are not Present

1. Remove all animals from premises.
2. Drain waterlines and watering appliances.
3. Prepare an activated 300 ppm use-solution of Carnebon™ 200 by:
 - a. Injection Using Metering Equipment: Add Carnebon™ 200 at the rate of 1 part to 65 parts of water. Once the Carnebon™ 200 has been diluted; inject 0.5 – 1.0 oz. per gallon “Biosolve™ AcidEdge™” or 0.5 – 1.0 oz. per gallon “Biosolve™ Acidic Cleaner – low foaming formula”.
 - b. Header Tank: Add 1.9 ounces of Carnebon™ 200 per gallon of water in a clean plastic header tank sufficient to refill water lines to deliver 300 ppm. Thoroughly mix solution, and then add 0.5 – 1.0 oz. per gallon “Biosolve™ AcidEdge™” or 0.5 – 1.0 oz. per gallon “Biosolve™ Acidic Cleaner – low foaming formula”. Trigger each nipple drinker to ensure contact with solution.
4. Turn on water supply or open filling valve to fill entire drinking water supply. Allow solution to remain in water lines for 4 – 8 hours.
5. Drain waterlines and flush with clean water.]

[DRINKING WATER FOR POULTRY, SWINE, CATTLE AND OTHER LIVESTOCK]

[To Control Taste and Odor in the Water Supply System

1. Prepare a solution with 5 ppm available chlorine dioxide by adding 1 part of Carnebon™ 200 per 4,000 parts of water (a 1:4,000 dilution) (1 fl. oz. Carnebon™ 200 to each 31.5 gallons). Allow 15 minutes before delivery to livestock or poultry.
2. If the water supply has heavy contamination prepare a solution of 11 ppm available chlorine dioxide by adding 1 part of Carnebon™ 200 per 1818 parts of water (a 1:1818 dilution) (1 fl. oz. Carnebon™ 200 to each 14 gallons). Allow 15 minutes before delivery to livestock or poultry.
3. After 24 hours, the addition rate can be reduced to 1 ppm of available chlorine dioxide by adding 1 gallon Carnebon™ 200 to each 20,000 gallons of animal drinking water as long as terminal concentration at end of waterline is not less than 0.5 ppm.
4. Treat water continuously from day one. Remove Carnebon™ 200 from drinking water 24 hours prior to vaccination, then resume treatment 24 hours after vaccinations. **Note:** *When treating drinking water for poultry, swine, cattle and other livestock, this product is not intended for use in human drinking water and treated water must not be made available for human consumption.*]

[To Disinfect Drinking Water Supply for Poultry, Swine, Cattle and Other Livestock

Use Carnebon™ 200 with a chlorine dioxide generator to generate an aqueous chlorine dioxide solution. Alternatively, Carnebon™ 200 can be mixed manually to generate an aqueous chlorine dioxide solution. The chlorine dioxide generator and manual mixing methods react Carnebon™ 200 with either a chlorine solution and acid or an acid. The generated chlorine dioxide solution can be added at a point in the system which ensures uniform mixing and distribution of up to 5 ppm of chlorine dioxide.

Follow all instructions for the chlorine dioxide generator carefully. Always prepare and use chlorine dioxide solutions in a well-ventilated area. Treat water continuously from day one. Remove Carnebon™ 200 from

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

drinking water 24 hours prior to vaccination, then resume treatment 24 hours after vaccinations.

Note: *When treating drinking water for poultry, swine, cattle and other livestock, this product is not intended for use in human drinking water and treated water must not be made available for human consumption.*

1. [Manual Mixing Method A

- A. For a 5 ppm chlorine dioxide solution add 1 part Carnebon™ 200 to 4,000 parts water; approximately 1.0 fl. oz. Carnebon™ 200 per 32 gallons of water. Use more water for lower chlorine dioxide concentrations.
- B. Add 2-5 ppm sodium hypochlorite; 1 – 3 parts of 12.5% bleach to 4,000 parts water.
- C. Using an appropriate acid add sufficient acid to lower solution pH to 5.0 to 6.5
- D. Allow 15 minutes before delivery to livestock water lines.
- E. After 24 hours, the addition rate can be reduced to 1 ppm of available chlorine dioxide by adding 1.0 fl. oz. of Carnebon™ 200 to approximately 160 gallons of animal drinking water as long as terminal concentration at the end of the water line is not less than 0.5 ppm.]

2. [Manual Mixing Method B

- A. Add 1 part Carnebon™ 200 to 3 parts water.
- B. Activate by adding phosphoric, hydrochloric, acetic or other food grade acid to a pH of 2.5-3.5.
- C. Mix and allow to stand for at least 15 minutes before delivery to livestock water lines.
- D. Then dilute 1 part of the activated solution with 1,000 to 5,000 parts water for a 1 to 5 ppm chlorine dioxide solution.]]

[+] [To Control Bacteria and Odor in the Egg Room

1. Wash down the entire egg room with high pressure water containing 20 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 1,000 parts with water) to remove gross filth or heavy soil.

Conduct the washing operations once per week (or more frequently in cases of heavy contamination during operations).

2. If it is necessary to clean the floors by mopping, then use 390 ppm of available chlorine dioxide (1 part Carnebon™ 200 per 50 parts water). Allow to dry on the floor.
3. A shoe or boot bath of 1,000 ppm of available chlorine dioxide (1 part Carnebon™ 200 per 20 parts water) is placed at the entrance to the egg room. Doors to the room must be kept closed at all times.
4. A glove dip, or rinse tank or basin, containing 50 ppm of available chlorine dioxide (1 part Carnebon™ 200 per 400 parts water) is used on entering and exiting the room.

Replace both the shoe and boot bath and glove dip daily (sooner if traffic is heavy).

5. Humidification water is treated with 40 ppm of available chlorine dioxide (1 part of Carnebon™ 200 per 500 parts water) to prevent the build-up and airborne spread of odor-causing microorganisms.
6. Provide 20 ppm of available chlorine dioxide (1 part Carnebon™ 200 per 1,000 parts water) to the water supply in the egg washing machine.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

[+] [To Control Odor and Bacteria when Separating Chicks in the Chick Room, Chick Grading Box and Sexing Room

1. Remove all poultry and feeds from premises, trucks, coops and crates.
2. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Spray the entire area for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 to 20 parts with water). Allow a 10 minute contact time.
6. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
7. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.
8. All workers in this area must use a hand dip or rinse containing 50 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 400 parts with water).
9. After use, wash area with high-pressure water to remove gross filth and soil.
10. Use a spray bottle containing a solution of 1,000 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 20 parts with water), on hands, wire mesh and in empty chick boxes to control contamination and odors from litter.
11. To clean the floor by mopping daily, use a solution containing 390 ppm of available chlorine dioxide (1 part Carnebon™ 200 per 50 parts water). Allow to air dry.]

[+] [To Control Bacteria and Odor in the Hatching Area

1. As soon as chicks are separated from Hatch, remove all trash containers with eggshells, down, etc. from the hatching area.
2. Remove all poultry and feeds from premises, trucks, coops and crates.
3. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
4. Empty all troughs, racks and other feeding and watering appliances.
5. Thoroughly clean all surfaces with soap or detergent and rinse with water.
6. Spray the entire area for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 to 20 parts with water). Allow a 10 minute contact time.
7. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.
9. All workers in this area must use a hand dip or rinse containing 50 ppm of available chlorinedioxide (1 part Carnebon™ 200 diluted to 400 parts with water).]

[+] [To Control Bacteria and Odor in the Incubator Room

1. The area is sprayed at least once per week for 5 minutes with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 diluted to 20 parts with water), after removing gross filth or soil with a high pressure water wash. Wet all surfaces and allow to dry.
2. Mop the floor daily with a solution containing 390 ppm of available chlorine dioxide (1 part of Carnebon™ 200 diluted to 50 parts with water).

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

3. Place a shoe and boot bath containing 1,000 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 50 parts with water) at all entrances to the incubator room.
4. 20 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 1,000 parts with water) is added to water in the humidification system or the air filters are sprayed with a 100 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 diluted to 200 part with water) to reduce airborne bacterial contamination.
5. Each time the eggs are removed from the incubator, a prior glove dip at 50 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 diluted to 400 parts with water) is recommended, followed by a spray of 1,000 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 diluted to 20 parts with water) on the eggs from a spray bottle.
6. Where containers are used to discard bad eggs, 1 part of Carnebon™ 200 per 7 parts of water (3,125 ppm of available chlorine dioxide) will control obnoxious odors and bacterial contamination.
7. Keep the doors to the area closed as much as possible to avoid airborne contamination.]

[+] [To Prevent Airborne and Surface Contamination of the Hatchery from the Tray Washing Room and Loading Platform

1. Close all doors in the tray washing room to avoid contamination of other hatchery operations. Discard all chick downs, egg shells, and cast-off chicks into the trash barrels and transfer the covered containers to the loading platform for disposal.
2. Wash the trays, carriages and other working equipment in a tray washing machine with 300-500 psi water to remove gross filth and soil.
3. As a final rinse in the tray washing machine, use a solution containing 20 ppm of available chlorine dioxide (1 part of Carnebon™ 200 diluted to 1,000 part with water) in high pressure water. Allow the trays, carriers and other working equipment to air dry. The walls, floors and carrying stands must also be sanitized with the same solution. Allow the equipment to air dry. Hold the sanitized equipment in a closed area for reuse.
4. Entrance and exit from the tray washing room must be through a foot rinse containing a solution of 1,000 ppm of available chlorine dioxide (1 part Carnebon™ 200 diluted to 20 parts with water). The rinse must be at least ½ inch deep and must be changed daily unless traffic is heavy.
5. After use, the tray washing room is washed with high pressure water to remove gross filth and soil. It is then decontaminated by spraying with a solution containing 1,000 ppm of available chlorine dioxide (1 part of Carnebon™ 200 diluted to 20 parts with water) for 15 minutes and allowed to air dry. This treatment is repeated after each use of the tray wash room.
6. The Loading Platform is washed from time to time to remove gross filth and soil. The trash containers are washed after discarding the contents to remove gross filth and soil. They are then sprayed with a 1,000 ppm solution of available chlorine dioxide (1 part Carnebon™ 200 diluted to 20 parts with water) and stored.]

[+] [Ventilation Systems

To treat non-porous hard surfaces for odor causing bacteria associated with ventilation and air conditioning duct work in residential, commercial, and institutional situations. Prior to inspecting, cleaning, treating or working on a ventilation system or its components, the system must be turned off or disconnected from any part of the system not isolated.

1. Mechanically clean, vacuum, or blow free of dirt, dust, mold and debris all duct work using a commercial duct cleaning system or service prior to treatment. The air ducts to be treated must be mechanically sound

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

and free or air leaks.

2. Preparation of Activated Use-Solution: Add 1 part Carnebon™ 200 to 40 parts water then adjust the pH of the diluted Carnebon™ 200 to 2.6 with acetic, citric, phosphoric, sulfuric, hydrochloric or other equivalent acid. Please contact you IDI or authorized representative regarding equivalent acids. Prepare in a well-ventilated area and avoid breathing any fumes which may be produced during activation. Protective eyewear is recommended.

Application of Activated Solution: Activated solutions may be sprayed into duct work. All surfaces must be thoroughly wetted for at least ten (10) minutes. When spraying use an appropriate spraying device. Spray application is the preferred method on large surfaces that are easily accessed by removing entry plates or access panels. The selected spray equipment must provide a consistent particle size (1-300 microns) and a uniform spray pattern using a 0.011" spray lip. Avoid excess wetting but be certain coverage is complete on the tops, sides and bottoms of the unlined sheet metal air ducts. All preexisting or treatment created access panels must be properly resealed or replaced in accordance with Industry Standards after servicing. During application, area must be closed as tightly as possible and sealed. After spraying, the area must be opened and aired for one (1) hour before re-entry. Active solutions may be irritating if inhaled; therefore always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. After application, allow to air dry. Treat as required. Always apply freshly made solutions. Never reuse activated solutions.]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

{OPTIONAL LOGOS/MARKINGS/3rd Party Certifications}



Certified to NSF/ANSI 60
Max. Use Level 206 mg/L



KOSHER

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL

TEXT IN {BRACES} ARE COMMENTS TO REVIEWER

Alternate Brand Name: Anthium™ BCD-200

{OPTIONAL MARKETING CLAIMS}

[Hospital Disinfectant]

[Food Contact Surface Sanitizer]

[Food Processing Water Sanitizer]

[Disinfectant]

[Sanitizer]

[Virucide*]

[Tuberculocide]

[Hospital Use]

[Food Processing Water]

[No Rinse Sanitizer]

[Institutional Use]

[Non-Flammable]

[Eliminates Mold and Mildew]

[Mold and Mildew Control]

[Combats Mold and Mildew]

[Concentrated Broad Spectrum Biocide]

[Disinfect Animal Drinking Water Lines]

[Poultry Premise Sanitation]

[Farm Premise Sanitation]

[Animal Laboratory Disinfection]

[To Control Taste and Odor in Water Supply System of Poultry, Swine, Cattle and Other Livestock]

Notes to reviewer: TEXT IN [BRACKETS] IS OPTIONAL
 TEXT IN {BRACES} ARE COMMENTS TO REVIEWER
 Alternate Brand Name: Anthium™ BCD-200

This product qualifies for emerging viral pathogen claims per the EPA’s ‘Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA- Registered Disinfectant Labels’ when used in accordance with the appropriate use directions indicated below.

This product meets the criteria to make claims against certain emerging viral pathogens from the following viral categories:

- Enveloped Viruses
- Large Non-Enveloped Viruses

For an emerging viral pathogen that is a/an...	...following the directions for use for the following supporting organisms on the label:
Enveloped virus	Canine parvovirus ATCC VR-2017
Large, non-enveloped virus	Canine parvovirus ATCC VR-2017

[Product Name] has demonstrated effectiveness against viruses similar to [name of emerging virus] on hard, non-porous surfaces. Therefore, [Product Name] can be used against [name of emerging virus] when used in accordance with the directions for use against Canine Parvovirus ATCC VR-2017 on hard, non-porous surfaces. Refer to the [CDC or OIE] website at [pathogen-specific website address] for additional information.

[Name of illness/outbreak] is caused by [name of emerging virus]. [Product Name] kills similar viruses and therefore can be used against [name of emerging virus] when used in accordance with the directions for use against Canine Parvovirus ATCC VR-2017 on hard, non-porous surfaces. Refer to the [CDC or OIE] website at [website address] for additional information.