519/2001

MAY 9 2001

Mr. Archy F. Slover Chemist Morgan-Gallackhe, Inc 8707 Millergrove Drive Santa Fe Springs, CA 90670-2001

Dear Mr. Slover:

ſ

(

ì

Subject: Updating First Aid Statements Per PR Notice 2001 ACIDISOL Acid Sanitizer EPA Registration Number 9152-18 Your Letter Dated March 22, 2001

The amended labeling referred to the above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted. Submit the following comment (s).

The size (net contents) of the product must be listed on the label. A stamped label is enclosed your records.

If you have any questions concerning this letter please contact Drusilla Copeland at (703) 308-6224.

Sincerely,

Adam Heyward Product Manager Regulatory Management Branch II Antimicrobials Division (7510C)

Enclosure

CONCURRENCES									
SYMBOL									
SURNAME									
DATE									

OFFICIAL FILE COPY

PER

12

مرمد ب	

ACCEPTED

with COMMENTS

in EPA Letter Dated:

Under the Federal Insecticide, Fungicide, and Rodenticide Act as

amended, for the pesticide,

9 2001

MAY

# ACIDISOL

**ACID SANITIZER** 

## ACTIVE INGREDIENTS

	registered under EPA Reg. No.
Phosphoric Acid	0152-18
Dodecyl Benzene Sulfonic Acid	<b>5.0%</b>
INERT INGREDIENTS:	
	<b>TOTAL 100.0%</b>

EPA REGISTRATION NO. 9152-18 EPA EST NO. 9152-CA-1

# **KEEP OUT OF REACH OF CHILDREN**

DANGER

See side panels for additional precautionary and practical treatment statements

# **NET CONTENTS:**

٤

Morgan-Gallacher, Inc.

#### FIRST AID If inhaled - Move person to fresh air. It person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, it possible. Call a poison control center or doctor for further treatment advice. It on skin Take off contaminated clothing or clothing - Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advise. Hold eye open and rinse slowly and gently with water for 15-20 minutes. If in eyes Remove contact lenses, it present, abor the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice. tf swallowed - Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by the poison control center or doctor. . Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Questions? Contact: 1-800-537-5567

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

# STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, tood, or teed by storage or disposal. Do not store on aide. Avoid impacting of side walls.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or punture and dispose of in a sanitary iandfill, or incineration or, if allowed by state and local authorities by burning. If burned, stay out of amoke. Do not discharge effluent containing this product in lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effuent containing this product to sever systems without previously notifying the local sevage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

# SPILL OR LEAK

Contain spill/selc. Place in an appropriate waste container. Flush spill area with water. Observe local, state and federal laws and regulations regarding disposal, spill, cleanup, removal or discharge.

# SANTA FE SPRINGS, CA 90670

### 10/00A

WHERE TO USE: Acidisol is recommended for sanitizing tanks, piping and processing equipment in: Dairies, Dairy Farms, Cheese Plants, Ice Cream Plants, Restaurants, Beverage Plants, and Food Processing Plants.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with labeling. Remove gross food particles and soll before the application of Acidisol by a pre-flush or pre-scrape or when necessary by a pre-soak treatment.

Thoroughly wash the surface or objects with a detergent or compatible cleaner followed by a potable water rinse prior to the application of the sanitizing solution.

For sanitizing by immersion: use 1 oz. Acidisol for every 2 gallons water. Contact time 2 minutes unless advised differently by your Health Department.

Dairy and Food Plant Equipment - Sanitize at least 5 minutes before products are run.

TO SANITIZE BY FLOW METHOD: Add 1 oz. Acidisol to 2 gallons water. Pump through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under pressure for at least 2 minutes to insure contact with all internal surface, drain the solution and air dry.

CIP: Rinse with warm water. Add 1 oz. Acidisol to 2 gallons water. Pump'solution through system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valve and hold under pressure for at least 10 minutes to insure contact with all internal surface. Drain the solution and air dry.

CIP OF FLOW METHOD OF SANITIZING: Add 1 oz. Acidisol to 2 gallons water. Pump through the system until full flow is obtained at all extremities. The system is completely filled with the sanitizer and all air is removed from the system. Close drain valve and hold under pressure for at least 2 minutes for the flow method and 10 minutes for CIP. Drain the solution and air dry.

**IMMERSION:** Use 1 oz. Acidisol for every 2 gallons of water. Contact time 2 minutes unless advised differently by your Health Departmentdry.

Acidisol fulfills the criteria of Appendix "F" of Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the Health Service when tested by the Chambers Method.

