



ITEM
330

Chlorine
BOOSTER

**An effective method of restoring the clarity
and sparkle to your swimming pool water.**

- * To clear up "Green" looking pool water
- * For use as a shock treatment

Active Ingredient: SODIUM DICHLORO - TRIAZINETRIONE - 100%

DANGER

Read antidote statement
and other precautions
on back panel.

Keep Out of Reach of Children

Rinse empty container thoroughly and discard it.

Net Wt. 1 Lb.

EPA 9157-18

SUN SWIMMING POOL CHEMICALS

Division of SUN CLEANSER CO.
LIVONIA, MICH.

DIRECTIONS

1. Test for pH and adjust to 7.2 to 7.6.
2. Add 1 lb. of Sun Chlorine Booster to each 10,000 gallons of water. Do not dump into pool at one place. Use a clean dry scoop made of metal, ceramic, plastic or glass.
3. Add directly to the pool water by sprinkling on the surface of the water, preferably at the deep end of the pool.
4. If possible keep filter running for 24 hours after adding Sun Chlorine Booster. Clean filter if necessary.
5. Test for Chlorine the next day and resume swimming when the Chlorine reading is below 1.5.
6. Resume adding to your daily dose of Chlorine so as to maintain a Chlorine Residual of 0.6 to 1.0 PPM.

DANGER

Danger. Corrosive. Causes eye damage. May cause burns on moist or broken skin. Do not get in eyes, on skin or on clothing. Dust is irritating to nose and throat. Avoid breathing dust. Harmful if swallowed. Strong oxidizing agent. Contact with organic material may cause fire or explosive decomposition. Moisture contamination or fire may liberate irritating and hazardous chlorine-gases. Do not mix with other chemicals. Keep container tightly closed when not in use. Keep away from open flames, heaters or other heat sources. Protect from moisture contamination.

ANTIDOTE:

EXTERNAL-In case of contact with skin or eyes, flush with water for 15 minutes. For eyes, get medical attention. Clean contaminated clothing before reuse.

INTERNAL-Drink milk or mucilage, followed by cooking oil. Call a physician.

MADE AND PRINTED IN U.S.A.

A

UND
FUN
FOR
ED

SANITIZER

DIRECTIONS

1. Establishment of Chlorine Residual of 0.6 to 1.0 ppm—For initial chlorination of newly filled pool, add 1 quart of Sun Sanitizer for each 20,000 gallons of water. Test pool water with a chlorine test kit, and if below 0.6 to 1.0 ppm repeat dosage until a chlorine residual of 0.6 to 1.0 ppm is established.

2. Routine chlorination Dosage—For maintaining a chlorine

residual of 0.6 to 1.0, add 1 quart of Sun Sanitizer daily or as needed as determined by the use of a chlorine test kit. Sun Sanitizer may be added to the pool water by pouring directly from container or with a feeding device. Use test kit frequently to determine chlorine residual. Frequency of dosage will depend on water temperature, pool usage, sunlight and pool size.

3. Maintenance of pH—should be maintained in the 7.2—7.6 range.

DEGRADES WITH AGE. USE A TEST KIT AND INCREASE DOSAGE
AS NECESSARY TO OBTAIN REQUIRED LEVEL OF AVAILABLE CHLORINE.

WARNING

Harmful if swallowed. May produce chemical burns. Strong oxidant.

Do not get on skin or in eyes. Irritating to eyes, skin, and lungs.

ANTIDOTES

External: Flood with cold water. For eyes, get medical attention.

Internal: Drink milk or raw egg whites. Call a doctor.

CAUTION

Keep bottle in upright position. Make sure cap is on tight. Do not spill on clothes. Do not store in direct sunlight. Do not store longer than 2 months. Rinse empty container thoroughly with water before discarding.

SUN SWIMMING POOL CHEMICALS

DIVISION OF SUN CLEANSER CO.

LIVONIA, MICHIGAN