

ACCEPTED
 4/3/70
 UNDER THE FEDERAL INSECTICIDE ACT
 FUNGICIDE ACT OF 1947 REGISTERED
 FOR ECONOMIC PEST CONTROL REGISTERED
 UNDER NO. 9159-88 SUBJECT
 TO ATTACHED COMMENTS.

U.S.D.A. Reg. No. 9159-88

DIRECTIONS FOR USE

Use ^{1/3} to 3/4 gallon KAY-KOP 80 diluted with water per acre in low or high volume sprays. The use of Fungicides for disease control is based on prevention, the plant surfaces must be covered with the Fungicide to successfully prevent infection.

INFORMATION

Since weather conditions and disease incidence vary, consult your State Agricultural Extension Service for timing and initial applications. Spray Equipment - fittings, nozzles, and all metal parts in contact with spray solution should be steel, iron, aluminum or plastic. Do not use in contact with brass, bronze, copper.

CAUTION

Harmful if swallowed. Avoid contact with skin. May cause irritation to the nose and throat. In case of ingestion, call physician immediately. Toxic to fish. Keep this product out of lakes, ponds or streams. Before re-use, flush container with water and detergent. Do not contaminate water sources by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

NOTICE

The Seller assumes no responsibility for injury to persons, animals or damage to plants resulting from failure to follow directions for the use or to observe cautions on this label. The buyer accepts the use of this material on these conditions.

KAY - KOP 80

FUNGICIDE SPRAY CONCENTRATE for Leafspot Control

ACTIVE INGREDIENTS

Copper, Metallic (from Copper Ammonium Carbonate)	8%
INERT INGREDIENTS	92%
TOTAL	100%

KAISER
AGRICULTURAL
CHEMICALS

KAY-KOPTM 80
 FUNGICIDE SPRAY CONCENTRATE
 for Leafspot Control

**CAUTION: KEEP OUT OF
 THE REACH OF CHILDREN**
 See other cautions and directions on side panels.

Manufactured For **KAISER AGRICULTURAL CHEMICALS**
 P.O. BOX 246, SAVANNAH, GEORGIA

KAISER

AGRICULTURAL CHEMICALS

KAY-KOPTM 80

**FUNGICIDE SPRAY CONCENTRATE
for Leafspot Control**

**CAUTION: KEEP OUT OF
THE REACH OF CHILDREN**
See other cautions and directions on side panels.

Manufactured For **KAISER AGRICULTURAL CHEMICALS**
P.O. BOX 246, SAVANNAH, GEORGIA

FOR USE ON:

BEANS

For control of common and bacterial blight. Beginning at disease appearance, spray 4 to 5 times at 10 - 14 day intervals.

CARROTS

For control of Early and Late Blight. When plants are 6" high, 3 to 5 applications at 7 - 10 day intervals.

CELERY

For control of Early and Late Blight. On appearance of disease, 4 to 7 applications at weekly intervals.

CUCURBITS

Cantaloupes, Melons, Cucumbers, Squash for control of Powdery Mildew, Downy Mildew, Scab, Alternaria and Angular leaf spots. Apply weekly beginning after vining and continue until mid-harvest.

GRAPES

For control of Anthracnose. When first leaves appear, just before blooming, just after blooming and at 2 week intervals thereafter until 2 weeks before harvest. Use with 4 pounds hydrated lime.

PEANUTS

For control of Cercospora Leaf Spot. On appearance of disease, 4 to 8 sprays, 10 - 14 day intervals.

PEPPERS

To control Cercospora Leaf Spot, Bacterial Spot. Apply at 7 to 10 day intervals.

POTATOES (Irish)

To control Early and Late Blight. Starting on or before first appearance of the disease in the area, 4 to 6 sprays, 7 to 10 day intervals.

STRAWBERRIES

To control Leaf Spot and Scorch. Spray at 7 to 10 day intervals from time new growth starts until harvest.

SUGAR BEETS

To control Cercospora Leaf Spot. Starting on disease appearance, 3 to 6 sprays at 14 day intervals.

TOMATOES

To control Early and Late Blight and Bacterial Spot, spray in seed beds at emergence and continue on 4 to 5 day schedules. In field, begin applications immediately after transplanting or emergence and continue at 4 to 5 day intervals. In areas or seasons when disease is less severe, time between applications may be 5 to 7 days. Complete coverage is essential to good control.