

DIRECTIONS: Place tablespoon amounts of bait (¼ to ½ ounce) in small paper plate or other disposable container at 8 to 12 foot intervals in locations where mice feed, water or travel. Bait placements along walls, in corners and concealed locations where mice can feed unobserved are desirable. Baiting should continue until all signs of feeding have stopped. DO NOT PLACE BAIT IN LOCATIONS ACCESSIBLE TO CHILDREN, PETS OR DOMESTIC ANIMALS.

Maintain an uninterrupted supply of bait for at least 15 days. After feeding has ended, leftover bait may be poured back into original container for future use. Contaminated or spoiled bait should be replaced promptly.

LIPP

MOUSE BAIT

Net Wt. 2 ounces

PROFESSIONAL
PEST CONTROL
FORMULA



ACCEPTED

1-18-71

9172-3

ACTIVE INGREDIENT:
(3-(Alpha-acetyl-
furfuryl)-
4-hydroxycou-
marin) 00.025%

INERT
INGREDIENTS: 99.975%

CAUTION: Keep out of reach of children, pets or domestic animals. In case of ingestion by humans call a physician at once.

NOTE TO PHYSICIAN: Treat as for overdose of Dicumarol.

Consumer Products Division
SCOTT CHEMICAL COMPANY
1135 BASSE ROAD
SAN ANTONIO, TEXAS 73212

USDA Reg. No. 9172-3 © 1970