

20 LBS. NET WT.

FATAL

**TO CRAB GRASS
IN 2500 SQ. FT. OF LAWN**

**Pre-Emergence
CRAB GRASS KILLER**

- One application of FATAL in spring stops crabgrass.
- Easy to apply—use standard lawn seed & fertilizer spreader.
- Flows freely and evenly—is clean & dustless.
- Fertilizer can be applied the same day as FATAL, if desired.
- For use on home lawns, parks & golf courses.

ACCEPTED
1-17-73
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR USE AS A CRAB GRASS KILLER
REGISTERED UNDER NO. 9198-1

DIRECTIONS

The contents of this package should be applied uniformly on 2500 sq. ft. of lawn area. Use on established lawns of Merion bluegrass, Kentucky bluegrass, mixed bluegrass, — fescue, Bermuda, Centipede, St. Augustine or Zoysia. Do not use on Dichondra.

Missed spots or streaks may allow crabgrass growth. Uniform and complete coverage with Fatal will prevent this. For best coverage use two applications at the half rate—one perpendicular to the other (note spreader setting chart).

Pre-Emergence CRAB GRASS KILLER

ACCEPTED
1-17-73
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR EQUIPMENT WHICH IS REGISTERED
UNDER NO. 9198-1

- One application of FATAL in spring stops crabgrass.
- Easy to apply—use standard lawn seed & fertilizer spreader.
- Flows freely and evenly—is clean & dustless.
- Fertilizer can be applied the same day as FATAL, if desired.
- For use on home lawns, parks & golf courses.

DIRECTIONS

The contents of this package should be applied uniformly on 2500 sq. ft. of lawn area. Use on established lawns of Merion bluegrass, Kentucky bluegrass, mixed bluegrass, —fescue, Bermuda, Centipede, St. Augustine or Zoysia. Do not use on Dichondra.

Missed spots or streaks may allow crabgrass growth. Uniform and complete coverage with Fatal will prevent this. For best coverage use two applications at the half rate—one perpendicular to the other (note spreader setting chart).

Application can be made in the fall, winter or spring when ground is not frozen and before crabgrass seeds germinate.

MAKE & MODEL	Approx. Setting for Double Application (TWICE OVER TREATMENT)	Approx. Setting for Single Application
ASGROW	6¾	8¼
CYCLONE	5¼	6¾
LAWN BEAUTY	6¾	8¼
SCOTT	5½	8½
GARDEN MARK	7	8½
JACKSON	6¾	8½
SEARS	6¾	8¼

CAUTION

Do not take internally. Avoid contact with skin. Avoid contact with eyes. Avoid contamination of feed, foodstuff and fish ponds. Keep out of reach of children. Wash thoroughly after handling. Do not reuse empty bag. Destroy bag by burying with waste. Do not burn.

Notice: If not used as directed, seller makes no warranty, expressed or implied, concerning the use of this product. User then assumes all risk of loss or damage arising from use, handling and the effects thereof.

MFG. BY *Free-Flow Fertilizer Co., Maumee, Ohio.* 43537

FOR PRE-EMERGENCE TREATMENT

ACTIVE INGREDIENTS Dimethyl Ester of Tetrachloroterephthalic Acid* 2.87%

INERT INGREDIENTS (RELATIVE TO HERBICIDAL CONTENT) 97.13%

Dacthal® Reg. TM of Diamond Shamrock Corporation **TOTAL.....100.00%**

*Use as preemergence herbicide patented under U.S. Patent No. 2,923,634.

CAUTION: KEEP OUT OF REACH OF CHILDREN—SEE REVERSE SIDE FOR ADDITIONAL CAUTIONS

Free-Flow Fertilizer Co., Maumee, Ohio.

43537

ACCEPTED

1-17-73

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. 9198-1