

Wickes



**WEED-N-FEED
24-6-6 plus 2**

**High Nitrogen Fertilizer
Plus Silvex & 2-4-D
Controls Broadleaf Weeds**
HERBICIDES

ACTIVE INGREDIENTS
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 0.69%
Isocetyl Ester of Silvex 2-(2,4,5-Trichlorophenoxy) Propionic Acid 0.66%

INERT INGREDIENTS 98.65%
100.00%


EQUIVALENT TO:
2,4-Dichlorophenoxyacetic Acid 0.57%
2-(2,4,5-Trichlorophenoxy) Propionic Acid 0.46%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See reverse side for additional cautions

EPA REG. NO. 9198-15

Dist. By MICHIGAN BEAN COMPANY / SAGINAW, MICHIGAN 48605

NET WT. 40 LBS.



**Wickes
BIG
ACRE
WEED-N-FEED
24-6-6
PLUS 2**

ACCEPTED

5-17-73

9198-15

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N).....	24%
Urea Nitrogen.....	18%
Ammoniacal Nitrogen.....	2%
Water Insoluble Nitrogen.....	4%
Available Phosphoric Acid (P ₂ O ₅).....	6%
Soluble Potash (K ₂ O).....	6%

This product controls the following weeds: broadleaf plantain, buck-horn plantain, dandelion, ground ivy, pigweed, purslane, lambsquarter, smartweed, common chickweed, mouse-ear chickweed, honeysuckle, and pennywort.

DIRECTIONS

Apply this product to lawns.
Apply anytime during growing season when temperature is above 70°. Apply when grass is wet from dew, rain or sprinkling.
For stubborn weeds, use two or three applications four weeks apart. Do not use on desirable clovers, bent, St. Augustine or dichondra grass. Do not water, mow or use lawn for 36 hours after application. Do not use on newly seeded areas until after third mowing. Treated areas may be seeded three to four weeks later - after rains or sprinkling.
Use a clean, properly adjusted spreader at settings shown below.
Suggested setting for 40 lbs. per 10,000 sq. ft.

Jackson.....	5	Modem.....	5
Sears.....	5	Scott.....	5
Cyclone.....	4½	Greenfield.....	6½

Do not reuse container. Destroy by burying in a safe place. Do not burn.

CAUTION:

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Do not contaminate feed, foodstuff or ponds. Do not store near seed, fertilizer, insecticides or fungicides. Keep out of reach of children. Wash thoroughly after using. Avoid drift of dust to nearby susceptible plants. Avoid breathing dust. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation or domestic purposes.

This product is toxic to fish. Keep out of any body of water. Destroy empty bags by burying. Do not reuse container. Do not burn. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

NOTICE

IF NOT USED AS DIRECTED, SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, CONCERNING THE USE OF THIS PRODUCT. USER THEN ASSUMES ALL RISK OF LOSS OR DAMAGE ARISING FROM USE, HANDLING AND THE EFFECTS THEREOF.

Dist. By MICHIGAN BEAN COMPANY / SAGINAW, MICHIGAN 48605

Free Trade Fertilizer Co. P.O. Box 111, Ann Arbor, Mich 48106

WEED-N-FEED

24-6-6 plus 2

High Nitrogen Fertilizer Plus Silvex & 2-4-D Controls Broadleaf Weeds HERBICIDES

ACCEPTED
MAY 17 1973
UNDER
FUNCTIONAL
FOR ECONOMIC
ED UNDER NO. 9198-15
TO ATTACHED C.
SIDE
ACT
SER.
SUBJECT

ACTIVE INGREDIENTS

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid	0.69%
Isooctyl Ester of Silvex 2-(2,4,5-Trichlorophenoxy) Propionic Acid.....	0.66%
INERT INGREDIENTS.....	98.65%
	100.00%

EQUIVALENT TO:

2,4-Dichlorophenoxyacetic Acid	0.57%
2-(2,4,5-Trichlorophenoxy) Propionic Acid	0.46%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See reverse side for additional cautions

EPA REG. NO. 9198-15

Dist. By MICHIGAN BEAN COMPANY / SAGINAW, MICHIGAN 48605

NET WT. 40 LBS.