NET WT. 40 L35.

WEEDSFEED

13-5-9

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)			. 18%
		2.0%	
		5.0%	
		11.0%	
),)	. 5%
		· · · · · · · · · · · · · · · · · · ·	
Boron (B)		Manganese (Mn)	.175%
Copper (Cu)	175%	Zinc (Zn)	.175%
Iron (Fe)		Molybdenum (Mo)0	0175%

WEED CONTROL

2,4-D for control of Broadleaf Plantain, Buckhorn Plantain, Bandelion, Purslane, Pigweed and Smartweed.

Silvex for control of Ground Ivy, Common Chickweed, Mouse-ear Chickweed, White Clover and Black Medic.

DIRECTIONS FOR APPLICATIONS

Apply to 10,000 square feet of lawn area.

Apply when temperature is above 60°.

Apply when lawn is wet from rain, dew or watering.

On newly seeded areas, wait until grass has been cut two or three times.

Treated areas may be seeded three or four weeks later.

Do not use on desirable clovers, bent, St. Augustine, or Dichondra Grass.

Apply when weeds are actively growing.

SPREADER SETTINGS

	cott
--	------

These settings are approximate. Age and condition of spreader may cause variations in settings.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid breathing dust. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate feed, foodstuffs or ponds. Do not store near seed, fertilizer, insecticides or fungicides. Drift of dust or vapors may cause injury to nearby susceptible plants.

Container Disposal: Do not reuse container, Destroy by burying with waste materials. Do not burn.

Fish and Wildlife Caution: This product is toxic to fish. Keep out of lakes, streams, or pends. Do not contaminate water by cleaning of equipment, or disposal of wastes and apply this product only as specified on this label. EPA REG. NO. 9198-25

P.O. BOX 119, MAUMEE, OHIO 43537