


Net Weight
40 lb.

Covers
8,000 sq. ft.

TURF CARE
FOR
PROFESSIONAL LAWN MAINTENANCE
20-4-10
with
0.92% Benefin



GUARANTEED MINIMUM FERTILIZER ANALYSIS

Total Nitrogen (N)	20%
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	10%

ACTIVE INGREDIENTS

N-butyl-n-ethyl-alpha, alpha, alpha-trifluoro-2,6 - Dinitro -p- Toluidine	0.92%
---	-------

INERT INGREDIENTS	99.08%
Total	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feeds.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal wastes. The manufacturer makes no warranties, express or implied, concerning this product or it's use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed. Destroy bags when empty. Do not reuse.

PRECAUTIONS: Apply Turf Care to the turf prior to the germination of weed grasses, since Turf Care will not control established weeds. Turf Care may thin established annual bluegrass (Poa annua) turfs and fine-leaved fescues at rates above 215 pounds per acre (2 lbs. active ingredient). Turf Care should not be applied in the spring to turfgrass planted the previous fall. Apply the product evenly over the turf area avoiding skips or missed spots. Follow application rates carefully do not overlap or over apply. Over application can weaken turf grass during stress periods allowing fungus or insects to cause damage. In bermudagrass that has been overseeded with winter grasses, a spring application of Turf Care may thin the overseeded grasses. Because of varying cultural practices Turf Care should not be used on putting greens. Turf Care should not be used on dichondra or bentgrass turf or applied to newly sprigged areas of bermudagrass, St. Augustinegrass or centipedegrass until these turfs are well established.

DIRECTIONS FOR USE OF TURF CARE: Turf Care is a pre-emergence herbicide with added growth promoting fertilizer ingredients which is applied to established turf for the control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass (silver crabgrass or crow's foot), watergrass (barnyardgrass) and yellow and green foxtail. Turf Care controls these annual weeds by killing their seeds as they germinate. Apply Turf Care before these annual weeds begin to germinate since it does not control established weeds.

When Turf Care is applied as directed, the following established turfgrasses are tolerant; perennial bluegrasses, perennial ryegrass, centipedegrass, fescue, zoysia-grass, bermudagrass, St. Augustinegrass and bahiagrass. Turf Care will not harm nearby established ornamental trees and shrubs.

Apply Turf Care to turf before the germination of annual grass weed. General times of application are discussed below for each type of weed. Precise time of application for a specific area will vary because of climate and individual cultural practices. Consult agricultural experimental station or extension service turf grass specialists for additional information regarding specific weed germination times in your area.

Any cultural practices that disturb the soil, such as aerification or verticutting should be done prior to applying Turf Care.

1. Annual bluegrass (Poa annua) control: In areas of heavy infestation of annual bluegrass (Poa annua) it's elimination will temporarily result in thin turf areas. Proper fertilization with Turf Care and irrigation will encourage the existing turfgrasses to fill in these thin areas formerly occupied by annual bluegrass (Poa annua).

2. Northern (cool season) Turf Grasses: For pre-emergence control of Poa annua apply 40 pounds to 8,000 sq. ft. (2 pounds active ingredient) of Benefin per acre in late summer to early fall (Mid-July to mid-September); or in early spring prior to annual blue grass (Poa annua) seed germination.

3. Crabgrass, goosegrass, watergrass and foxtail control: Northern (cool season) Turf Grasses: For pre-emergence control apply 30 pounds to 8,000 sq. ft. (1.5 pounds active ingredient of Benefin per acre) in the spring 2 to 4 weeks prior to expected annual weed grass seed germination. In the southern half of Ohio, Illinois, Indiana and all points south apply 40 pounds to 8,000 sq. ft. (2 pounds active ingredient of Benefin per acre) in the spring 2 to 4 weeks prior to expected annual weed grass seed germination.

RESEEDING: Delay reseeding at least 6 weeks after application of Turf Care.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: THIS PRODUCT IS DESIGNED AND PACKAGED EXCLUSIVELY FOR SALE AND APPLICATION BY PROFESSIONAL TURF MAINTENANCE PERSONNEL.

E.P.A. Reg No. 9198-30
E.P.A. Est. No. 9198-OH-1

MANUFACTURED BY:

FREE FLOW FERTILIZER
P. O. Box 119
Maumee, Ohio 43537

Net Weight
40 lb.

Covers
8,000 sq. ft.

TURF CARE
FOR
PROFESSIONAL LAWN MAINTENANCE
20-4-10
with
0.92% Benefin

ACCEPTED

11/12/1977

GUARANTEED MINIMUM FERTILIZER ANALYSIS

Total Nitrogen (N)	20%
Available Phosphoric Acid (P_2O_5)	4%
Soluble Potash (K_2O)	10%

ACTIVE INGREDIENTS

N-butyl-n-ethyl-alpha, alpha, alpha-trifluoro- 2,6 - Dinitro -p- Toluidine	0.92%
---	-------

INERT INGREDIENTS	99.08%
Total	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feeds.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal wastes. The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed. Destroy bags when empty. Do not reuse.

PRECAUTIONS: Apply Turf Care to the turf prior to the germination of weed grasses, since Turf Care will not control established weeds. Turf Care may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 215 pounds per acre (2 lbs. active ingredient). Turf Care should not be applied in the spring to turfgrass planted the previous fall. Apply the product evenly over the turf area avoiding skips or missed spots. Follow application rates carefully do not overlap or over apply. Over application can weaken turf grass during stress periods allowing fungus or insects to cause damage. In bermudagrass that has been overseeded with winter grasses, a spring application of Turf Care may thin the overseeded grasses. Because of varying cultural practices Turf Care should not be used on putting greens. Turf Care should not be used on dichondra or bentgrass turf or applied to newly srrigged areas of bermudagrass, St. Augustinegrass or centipedegrass until these turfs are well established.

DIRECTIONS FOR USE OF TURF CARE: Turf Care is a pre-emergence herbicide with added growth promoting fertilizer ingredients which is applied to established turf for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goose grass (silver crabgrass or crowfoot), watergrass (barnyardgrass) and yellow and green foxtail. Turf Care controls these annual weeds by killing their seeds as they germinate. Apply Turf Care before these annual weeds begin to germinate since it does not control established weeds.

When Turf Care is applied as directed, the following established turfgrasses are tolerant; perennial bluegrasses, perennial ryegrass, centipedegrass, fescue, zoysia-grass, bermudagrass, St. Augustinegrass and bahiagrass. Turf Care will not harm nearby established ornamental trees and shrubs.

Apply Turf Care to turf before the germination of annual grass weed. General times of application are discussed below for each type of weed. Precise time of application for a specific area will vary because of climate and individual cultural practices. Consult agricultural experimental station or extension service turf grass specialists for additional information regarding specific weed germination times in your area.

Any cultural practices that disturb the soil, such as aerification or verticutting should be done prior to applying Turf Care.

1. Annual bluegrass (Poa annua control: In areas of heavy infestation of annual bluegrass (Poa annua) its elimination will temporarily result in thin turf areas. Proper fertilization with Turf Care and irrigation will encourage the existing turfgrasses to fill in these thin areas formerly occupied by annual bluegrass (Poa annua).

2. Northern (cool season) Turf Grasses: For pre-emergence control of Poa annua apply 40 pounds to 8,000 sq. ft. (2 pounds active ingredient) of Benefin per acre in late summer to early fall (Mid-July to mid-September); or in early spring prior to annual blue grass (Poa annua) seed germination.

3. Crabgrass, goosegrass, watergrass and foxtail control:
Northern (cool season) Turf Grasses: For pre-emergence control apply 30 pounds to 8,000 sq. ft. (1.5 pounds active ingredient of Benefin per acre) in the spring 2 to 4 weeks prior to expected annual weed grass seed germination. In the southern half of Ohio, Illinois, Indiana and all points south apply 40 pounds to 8,000 sq. ft. (2 pounds active ingredient of Benefin per acre) in the spring 2 to 4 weeks prior to expected annual weed grass seed germination.

RESEEDING: Delay reseeding at least 6 weeks after application of Turf Care.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: THIS PRODUCT IS DESIGNED AND PACKAGED EXCLUSIVELY FOR SALE AND APPLICATION BY PROFESSIONAL TURF MAINTENANCE PERSONNEL.

E.P.A. Reg No. 9198-30
E.P.A. Est. No. 9198-OH-1

MANUFACTURED BY:

FREE FLOW FERTILIZER
P. O. Box 119
Maumee, Ohio 43537