

Net Weight 40 Lbs. (18.2 kg.)
Treats 14,000 sq. Ft.

Tee Time™ Fertilizer 30-3-5 with Dursban®

GUARANTEED ANALYSIS

Total Nitrogen (N) 30%
1 2% Ammoniacal Nitrogen
22 8% Urea Nitrogen
6 0% Coated Slow Release Urea Nitrogen
Available Phosphorus Acid (P₂O₅) 3%
Soluble Potash (K₂O) 5%
Sulfur (S) 4.5%
Primary Plant Foods Derived from Urea, Murates of Potash, Ammoniacal Phosphates, and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea.
Potential Acidity 1104 pounds Calcium Carbonate equivalent per ton.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS May Be Harmful If Swallowed

CANTINE: Do not take internally—Avoid contact with eyes, skin, and clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Avoid breathing dust. Keep children and pets off treated areas until the material is washed into the soil and grass is dry. Do not treat vegetable gardens or other plants used for food purposes.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
IF ON SKIN: In case of contact, wash contaminated clothing and immediately wash skin with soap and water.
IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

NOTE TO PHYSICIAN: Chelerythrin is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply to direct treatment or residues on plants are highly susceptible. Do not apply directly to water. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Application: Apply at the rate of 2.0 pounds per 1,000 square feet or the equivalent of 1.13 pounds actual chelerythrin per acre. At this rate the product will control chinch bugs and weevils (brown moths) and brown dog ticks, chiggers, cutworms, and earwigs. Water down after application.
To control certain white grubs (larvae of European chaffer and Japanese beetle) use 5 pounds of this product per 1,000 sq. ft. Apply the granules uniformly over the area treated followed immediately by thoroughly watering to wash the insecticide into the soil. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for greater facts to treat in your area.

PRODUCT APPLICATION RATES		LBS. OF NUTRIENT/50 FT.			
Lbs. Material/Sq. Ft.	Lbs. Material/Acre	N	P	K	S
2.0	121.9	64	60	14	13

SUGGESTED SPREADER SETTINGS

Spreader	Speed	Coverage Width	Settings	
			Full Rate	Half Rate
Cyclone	3 MPH	6 ft.	4 1/4"	3 1/4"
Scots	3 MPH	6 ft.	1 1/2"	1 1/4"
Lady	5 MPH	30 ft.	4"	3"

Important: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

Store in a dry place. Do not contaminate water, food, or feed by storage or disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Seller assumes the risk of any contrary use. Seller makes no other warranty or implied warranty including the warranty of fitness for a particular purpose, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material in which a claim is made. Seller: This product is designed and packaged exclusively for sale and application by professional turf maintenance personnel.
Dursban® is a registered trademark of E.I. du Pont de Nemours & Co.

AT73004

Manufactured by
**the professional's
partner™**

**The
Andersons**

Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537

Reg # 9198-32

PM-12

194

(FRONT PANEL)

TURF CARE FOR LAWN MAINTENANCE

38-0-0

WITH DURSBAN(R) BRAND INSECTICIDE

ACTIVE INGREDIENT:

Chlorpyrifos [O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl)
phosphorothioate] 0.92%

INERT INGREDIENTS 99.08%

TOTAL 100.00%

Dursban(R) Trademark of DowElanco

KEEP OUT OF REACH OF CHILDREN


CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS


NET WEIGHT __ LBS.

TREATS _____ SQ.FT.

EPA Reg. No. 9198-32



EPA Est. No. 9198-OH-1



(Back Panel)

NET WEIGHT __ LBS.

TREATS _____ SQ.FT.

TURF CARE FOR LAWN MAINTENANCE

38-0-0

WITH DURSBAN(R) BRAND INSECTICIDE

GUARANTEED ANALYSIS

Total Nitrogen (N)	38%
Available Phosphoric Acid (P_2O_5)	0%
Soluble Potash (K_2O)	0%

Primary Plant Foods Derived From: Urea, Muriate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes. If in eyes, flush with plenty of water. Get prompt medical attention if irritation persists. Do not take internally. Wash thoroughly after using. Wash contaminated clothing before reuse. Do not treat vegetable gardens or other plants for food purposes.


STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.



ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Bees exposed to direct treatment or residues on plants are highly susceptible. Do not apply directly to water. Keep out of lakes, streams and ponds. Do not contaminate water by disposing of equipment washings. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SURFACE INSECT CONTROL

Apply at the rate of 2.5-2.8 lbs. per 1000 square feet, or the equivalent of 1.00-1.13 lbs. actual active ingredient per acre. At this rate, the product will control chinch bugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, grasshoppers, armyworms, centipedes, clover mites, fiery skipper, fleas, greenbugs, June beetles, leafhoppers, millipede mites, mosquitoes, sowbugs and ticks.

Treat when insects first appear, re-treat as necessary. For best results, product should be watered in after application.

GRUB CONTROL

Apply at the rate of 5-10 lbs. per 1000 square feet, or the equivalent of 2-4 lbs. actual active ingredient per acre. At this rate, the product will control certain white grubs (larvae of European chaffer and Japanese beetle).

Treat when grubs are young and actively feeding near the soil surface. Consult your local agricultural extension service for proper time to treat your area. For best results, product should be thoroughly watered in to wash the insecticide into the zone of insect infestation.

To control Hyperodes (turfgrass) weevils, uniformly apply 5 pounds of granules per 1,000 sq. ft. as directed above. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialists for best time to treat in your area.

SUGGESTED SPREADER SETTINGS

(available at final printing)

Important: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

