


Constantine's

**WEED KILLER
AND FERTILIZER**




**WEEDS AND FEEDS
5000 SQ. FT.**

RICHFIELD

**WEED KILLER
AND FERTILIZER**

Constantine's



HARDER

20 lbs. net weight Covers 5,000 sq. ft.

Constantine's WEED KILLER AND FERTILIZER

18-4-6

GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
4.00% Water Insoluble Nitrogen	
Available Phosphoric Acid (P_2O_5)	4%
Soluble Potash (K_2O)	6%

**MANUFACTURED FOR
CONSTANTINE'S
NURSERY & GARDEN, INC.
RICHFIELD, OHIO 44286**

DOUBLE ACTING KILLS LAWN WEEDS—FEEDS GRASS

WEED and FEED does two jobs at one time. Kills weeds and feeds the turf in the same application. No more back breaking digging of dandelion or plantain when WEED and FEED is used. The high quality of turf food gives outstanding results in a thicker, greener turf. Stays green too for weeks longer. WEED and FEED can be applied any time the weeds are in a rapid state of growth. DO NOT APPLY WHEN TEMPERATURES ARE BELOW 60° (except for chickweed) OR WHEN MOISTURE IS LACKING IN THE SOIL.

DIRECTIONS

ESTABLISHED LAWNS — It is most important to apply evenly. Contents will cover 5,000 sq. ft. For more effective results lawn may be watered lightly before applying. Lawn seed may be sown 10-14 days after application of WEED and FEED.

MOWED — Grass should not be mowed for several days before and after application.

The following weeds are controlled by WEED and FEED. Chickweed (see below), white clover, henbit, thyme-leaved speedwell, yarrow, spotted spurge, dandelion, plantain, buckhorn, ragweed, burdock, lawn perennials, mustard, nettle, pig weed, ground ivy, sea holly and jimson weed.

CHICKWEED — Early spring control is preferred, however, late summer or early fall is also recommended to destroy weeds that may have developed during the summer or spring. Application can be made any time temperature is above freezing.

IMPORTANT — Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that the plants be syringed thoroughly with water as soon as possible. Vapors from the product may injure susceptible plants in the immediate vicinity.

Do not store near other fertilizers, seeds, insecticides, or fungicides. Certain grasses such as bent, St. Augustine, Dichondra, and carpet grasses are sensitive to herbicides and should not be treated.


CAUTION — Avoid prolonged or repeated inhalations of dust or contact with skin, eyes or clothing. Wash thoroughly after handling. To protect fish and wildlife, don't use where runoff will contaminate streams, lakes or ponds. This product is toxic to fish; keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated as this product may injure susceptible plants in the immediate vicinity. Do not reuse empty container. Burn or discard in a safe place away from water supplies. Clean spreader thoroughly after application. Do not contaminate water used for domestic or irrigation purposes.

SPREADER SETTING GUIDE


NUTRI	GREENFIELD	MODERN
3%	6	4
JACKSON	CYCLONE	SCOTT
5	4%	6

These settings are approximate. Settings vary with model, age and condition of spreader.

HARDER



WEED KILLER AND FERTILIZER



WEED KILLER AND FERTILIZER



20 lbs. net weight Covers 5,000 sq. ft.

Constantine's WEED KILLER AND FERTILIZER

18-4-6

GUARANTEED ANALYSIS

Total Nitrogen (N)	1.8
C:N Ratio - Insoluble Nitro.	1.8
Avg. Soluble Acids (P ₂ O ₅)	6.1
Soluble Potash (K ₂ O)	1.1

MANUFACTURED FOR
**CONSTANTINE'S
NURSERY & GARDEN, INC.**
RICHFIELD, OHIO 44286

DOUBLE ACTING KILLS LAWN WEEDS-FEEDS GRASS

Wildlife officials say two days of shooting this week at a "Feed the Birds" reservoir near the lake drew a total of 120 deer, including 40 whitetails, 75 mule deer and 15 elk. The results were gratifying.

DIRECTIONS

ESTABLISHED LAWNS - The most common lawns are those that have been established by seed or by cuttings. These lawns are usually more expensive than those established by sod.

MOWING - grass should be mown for several days after the blossoms. The mowing wheels are to be cleaned and dried. Then the mowing machine can be used to clean the straw in the long grass. This will help to keep the grass from becoming too tall.

CHICKWEED (*Calandrinia*) - This is a common weed, especially in lawns, and it is very difficult to eradicate.

IMPORTANT - Several hundred acres of land in the eastern part of the state were planted to corn and other cereal grains in the fall of 1937. The yield was excellent and the grain was well received by the market. It is recommended that the yields of the spring 1938 crop be limited to 10 bushels per acre. This will insure a good market for the grain.

CAUTION Avoid prolonged contact with skin. If contact occurs, wash skin with water and soap. If irritation continues, call a physician. If eyes are affected, flush eyes with water. If irritation continues, call a physician.

SPREADER SETTING GUIDE