



Tee TimeTM Fertilizer 20-4-10

with Benefin/Dursban[®]

the professional's
partnerTM



ACTIVE INGREDIENTS:
(1) butyl n-ethyl alpha, alpha, alpha trichloro 2,6-Dinitro p-Toluidine) . . . 0.92%
(O,O-Diethyl (O-(3,5,6-Trichloro-2-Pyridyl) Phosphorothioate) . . . 0.46%
INERT INGREDIENTS: . . . 98.62%
TOTAL . . . 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Weight 40 Lbs. (18.2 kg)
Treats 8,000 Sq. Ft.

Net Weight 40 Lbs. (18.2 kg) Treats 8,000 Sq. Ft. **Tee Time[®]** **Fertilizer 20-4-10** with Benefin/Dursban[®]

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
15% Ammoniacal Nitrogen	
12.5% Urea Nitrogen	
60% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	10%
Sulfur (S)	4.5%
Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, Sulfur Coated Urea, and Murates of Potash Sulfur derived from Sulfur Coated Urea	
Potential Acidity 1082 pounds Calcium Carbonate equivalent per ton	

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Avoid contact with eyes. If in eyes, flush with plenty of water. Get prompt medical attention if inhaled. Do not take internally. Wash thoroughly after using. Wash contaminated clothing before reuse. Keep children and pets off treated area until this material is washed into the soil and grass is dry. Do not feed vegetable gardens or other plants used for food purposes.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not expose to direct treatment or residues on plants are highly susceptible. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply when weather favors drift from treated area.

Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote by injection only.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Tee Time 20-4-10 with Benefin/Dursban[®] is a combination of growth promoting fertilizer ingredients, a pre-emergence herbicide, and a turf insecticide. It is designed to control a smooth and large (fine) crabgrass, goosegrass (after crabgrass or downy), bermudagrass (after crabgrass), yellow and green foxtail, annual bluegrass (Poa Annua), as well as ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, darts, grasshoppers, and sod webworms. When applied as directed, the following established turf grasses are tolerant: perennial bluegrasses, perennial ryegrass, fescue and zoysia grass. This product will not harm nearby established ornamental trees and shrubs.

Apply prior to the germination of weed grasses since it will not control established weeds. Pre-treatment of application for a specific area will vary because of climate and individual cultural practices. Consult Agricultural Experiment Station or extension service turf grass specialists for additional information regarding specific weed germination times in your area.

Because of varying cultural practices, this product should not be used on putting greens, declining, bentgrass, buff, bermudagrass, St. Augustine grass, or centipede grass. This product may damage leaves and stems at rates above 215 pounds per acre (2 lbs. active Benefin). In areas where annual bluegrass (Poa Annua) is present as a desirable species, this product should not be applied. In areas that have been over seeded with winter grasses, a spring application may harm the over seeded grasses. To prevent thinning of desirable grasses, it should not be applied in the spring to turf grasses planted the previous fall and reseeded of new areas should be delayed at least 6 weeks after application.

Because Tee Time 20-4-10 with Benefin/Dursban[®] is generally applied very early in the insect cycle, it may be necessary to give a second treatment of insecticide late in the season. Do not apply this combination product twice in the same season.

Any cultural practices that disturb the soil, such as aerification or verticuring, should be done prior to application. This material must be incorporated into the soil for best results. Check application rate of your spreader before and during application.

How to Apply: Apply at the rate of 5 pounds per 1,000 sq. ft. (50 lb. x 20 ft. = 1,000 sq. ft.) For best control, the product must be applied evenly over the turf without skips or misses. Best results are obtainable by using a two step procedure, which applies half the required amount on the first trip over the area. Spread the second half of the required amount over the same area by moving at right angles to the first application in a checkboard fashion. This procedure will minimize the chances of color streaking and erratic insecticide coverage control. Do not over or under apply. Over application can weaken turf grasses during stress periods allowing fungus or insects to cause damage.

Annual Bluegrass (Poa Annua) Control: For pre-emergence control, apply 5 pounds per 1,000 sq. ft. in late summer to early fall (mid-July to mid-September) or in early spring prior to seed germination. In areas of heavy infestation, the elimination of Poa Annua will temporarily result in thin turf areas. Proper fertilization with Tee Time and single-don will encourage starting turf grasses in fall in these thin areas.

Crabgrass, Goosegrass, Watergrass, and Foxtail Control: Apply in the spring 2 to 4 weeks prior to expected annual weed grass seed germination at 5 pounds per 1,000 sq. ft.

Ant, Armyworms, Chinch Bugs, Crickets, Brown Dog Ticks, Cutworms, Earwig, Grasshopper, Sod Webworms Control: Tee Time 20-4-10 with Benefin/Dursban[®] will control these insects present at the time of or soon after treatment when applied at the same rate for pre-emergence control of the listed weed species.

PRODUCT APPLICATION RATES		LBS OF NUTRIENT PER SQ FT				
Lbs Material	Sq Ft	Lbs Material/Acre	N	P	K	S
5		215	1	2	3	225

SUGGESTED SPREADER SETTINGS

Spreader	Speed	Coverage Width	Settings	
			Fertilizer	Inerticide
Cyclone	3 MPH	8 ft.	4 1/4 G/W	3 1/4 G/W
Scout	3 MPH	8 ft.	4 1/4 G/W	0 1/4 G/W
Lely	5 MPH	30 ft.	6 W	4 W

Important: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

Storage: Keep container closed and away from food, feedstuff or domestic water supplies. Disposal: Weeds resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Notice: This product is designed and packaged exclusively for sale and application by professional turf maintenance personnel.

Dursban[®] - trademark of Dow Chemical Company U.S. Patent No. 3,244,286

Manufactured by
**the professional's
partner[™]**

**The
Andersons**

Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537

BEST AVAILABLE COPY