

(front panel)

TEE TIME FERTILIZER

20-4-10

WITH BENEFIN/DURSBAN(R) BRAND INSECTICIDE

the professional's partner™

THE ANDERSONS

ACTIVE INGREDIENTS

(N-butyl-n-ethyl-alpha,alpha,alpha-trifluoro-2,6-Dinitro-p-Toluidine . .	0.92%
Chlorpyrifos: [O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl)	
phosphorothioate]	0.46%
INERT INGREDIENTS.	<u>98.62%</u>

TOTAL

100.00%

Dursban(R) Trademark of DowElanco

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT __ LBS.

TREATS _____ SQ.FT.

EPA Reg. No. 9198-37

EPA Est. No. 9198-OH-1

(back panel)

NET WEIGHT __ LBS.

TREATS _____ SQ.FT.

TEE TIME

FERTILIZER

20-4-10

WITH BENEFIN/DURSBAN(R) BRAND INSECTICIDE

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
Available Phosphoric Acid (P_2O_5)	4%
Soluble Potash (K_2O)	10%

Primary Plant Foods Derived From: Urea, Muriate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not take internally -- Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash contaminate clothing before reuse. Avoid breathing dust. Do not treat vegetable gardens or other plants used for food purposes.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Bees exposed to direct treatment or residues on plants are highly susceptible. Do not apply directly to water. Keep out of lakes, streams and ponds. Do not contaminate water by disposing of equipment washings. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Tee Time 20-4-10 with Benefin/Dursban(R) Brand Insecticide is a combination of growth promoting fertilizer ingredients, a preemergence herbicide and a turf insecticide. It is designed to control smooth and large (hairy) crabgrass, goosegrass (silver crabgrass or crowfoot), barnyard grass (watergrass), low and green foxtail, annual bluegrass (Poa Annua), as well as ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers, and sod webworms. When applied as directed, the following established turf grasses are tolerant: perennial bluegrasses, perennial ryegrass, fescue and zoysia grass. This product will not harm nearby established ornamental trees and shrubs.

Apply prior to the germination of weed grasses since it will not control established weeds. Precise time of application for a specific area will vary because of climate and individual cultural practices. Consult Agricultural Experiment Station or extension service turf grass specialists for additional information regarding specific weed germination times in your area.

Because of varying cultural practices, this product should not be used on putting greens, dichondra, bentgrass, turf, bermudagrass, St. Augustine grass, or centipede grass. This product may thin fine-leaved fescues at rates above 215 pounds per acre (2 lbs. active Benefin). In areas where annual bluegrass (Poa Annua) is present as a desirable species, this product should not be applied. In areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. To prevent thinning of desirable grasses, it should not be applied in the spring to turf grasses planted the previous fall and reseeded of the new areas should be delayed at least 6 weeks after application.

Because Tee Time 20-4-10 with Benefin/Dursban(R) Brand Insecticide is generally applied very early in the insect cycle, it may be necessary to give a second treatment of insecticide late in the season. Do not apply this combination product twice in the same season.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application. This material must be adequately watered into the soil for best results. Check application rate of your spreader before and during application.

How To Apply: Apply at the rate of 5 pounds per 1,000 sq.ft. (50 ft. x 20 ft. = 1,000 sq.ft.). For best control, this product must be applied evenly over the turf without skips or misses. Best results are obtainable by using a two-step procedure, which applies half the required amount on the first trip over the area. Spread the second half of the required amount over the same area by moving at right angles to the first application in a checkerboard fashion. This procedure will minimize the chances of color streaking and erratic insect/crabgrass control. Do not over or under apply. Over application can weaken turf grasses during stress periods allowing fungus or insects to cause damage.

Annual Bluegrass (Poa Annua) Control: For preemergence control, apply 5 pounds per 1,000 sq.ft. in late summer to early fall (mid-July to mid-September) or in early spring prior to seed germination. In areas of heavy infestation, the elimination of Poa Annua will temporarily result in thin turf areas. Proper fertilization with Tee Time and irrigation will encourage existing turf grasses to fill in these thin areas.

Crabgrass, Goosegrass, Watergrass, and Foxtail Control: Apply in the spring 2 to 4 weeks prior to expected annual weed grass seed germination at 5 pounds per 1,000 sq.ft.


Ant, Armyworms, Chinch Bugs, Crickets, Brown Dog Ticks, Cutworm, Earwig, Grasshopper, Sod Webworm Control: Tee Time 20-4-10 with Benefin/Dursban(R) Brand Insecticide will control these insects present at the time of or soon after treatment when applied at the same rate for preemergent control of the listed weed species.

PRODUCT APPLICATION RATES		LBS. OF NUTRIENT/SQ.FT.			
LBS.MATL/1000 SQ.FT.	LBS. MATL/ACRE	N	P	K	S
5	218	1	2	5	225

SUGGESTED SPREADER SETTINGS

(available at final printing)

Important: These settings are only approximate. Ag and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.



STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact you State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonable fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By:

the professional's partner™

THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537