(front panel)

TURF CARE

FOR

SOUTHERN LAWNS

0.47% DURSBAN(R) BRAND INSECTICIDE

	ve Ingredient : .orpyrifos [0,0-Di	ieth	ıyl	0.	– C	3,5	5,6	5-t	ir i	i ct	nla	ع د	o~:	2 ~ -g	p y r	-ic	dy]											
	phosphorothicate:		•			•													•	•	•	•	•	-	•	Ø	. 477	•
I s ert	: Ingredients: .			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		-	•	•	•	•	99	<u>, 5</u> 3	•
										Total						1 188	.00.00%											
	_																											

Dursban(R) Trademark of DowElanco

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT __ LBS.

TREATS ____ SQ. FT.

EPA Reg. No. 9198-43

EPA Est. 9198 UM

(back panel)

NET WEIGHT __ LBS.

TREATS ____ SQ. FT.

TURF CAPE

FOR

SOUTHERN LAWNS

Ø.47% DURSBAN(R) BRAND INSECTICIDE

			GUARANTEED ANALYSIS																
Potal Ni	truger	r (142								•		•		-	•	•		-	J81.
Primary	flant	Facd	Sour	ces:	Unes	and	Sul	. Fur	Coa	tea	Uje-	ea.							

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

TION: Avoid contact with eyes. If in eyes, flush with plenty of water. Get prompt medical attention if irritation pensists. Do not take internally. Wash thoroughly after using. Wash contaminated clothing before reuse. Do not treat vegetable gardens or other plants used for food purposes.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink two glasses of water and induce vomiting by touching path of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Freat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Bees exposed to direct treatment or residues on plants are highly susceptible. Do not apply directly to water. Keep out of lakes, streams and ponds. Do not contaminate water by disposing of equipment washings. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SURFACE INSECT CONTROL

Apply at the rate of 5.0-5.5 lbs. per 1000 square feet, or the equivalent of 1.00-1.13 lbs. actual active indredient per acre. At this rate, the product will control chinch bugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, carwigs, grasshoppers, armyworms, centipedes, clover mites, frery skipper, fleas, greenbugs, June bettles, leafhoppers, millipedes, mites, mosquitoes, sowbugs and ticks.

Treat when insects first appear, re-treat as necessary. For best results, product should be watered in after application.

GRUB CONTROL

Apply of the male of 18-10 lbs. per 1000 square feet, or the euclivatent of 2-1 lbs. actual active ingredient per acre. At this rate, the product will control centum white grubs (larvae of European chaffer and Japanese beetle).

Treat when grubs are young and actively feeding near the soil surface. Consult your local agricultural extension service for proper time to treat in your area. For best results, product should be thoroughly watered in to wash the insecticide into the zone of insect infestation.

To control Hyperodes (turfgrass) weevils, uniformly apply 10 pounds of granules per 1,000 sq. ft. as directed above. Treat suspected problem armass in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialists for best time to treat in your area.

SUGGESTED SPREADER SETTINGS (available at final printing)

Important: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess posticide, spray mixture, or rinsate, is a violation of Federal Lawn and may taminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact you State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay our of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably rit for the purposes set forth in the Directions for Use. EXCEPT AS SU WARRANTED. THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OF IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

THE ANDERSONS LAWN FERTILIZER DIVISION P.D. BOX 119 MAUMEE, DHIO 43537

:9198~43R 04/03/90