


The contents of this bag  
treat 5000 sq. ft.

TO OPEN:  
Pull Kordon Tape



# ANDERSONS WEED AND FEED XX-XX-XX

Contains 2, 4-D and MCPP

## HERBICIDE

### ACTIVE INGREDIENTS:


Dimethylamine Salts of 2, 4-Dichlorophenoxyacetic Acid	0.01%
Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propanoic acid	0.01%

### EQUIVALENT TO:

2, 4-Dichlorophenoxyacetic Acid	0.84%
2-(2-methyl-4-chlorophenoxy) propanoic acid	0.84%

### INERT INGREDIENTS.....

TOTAL	97.93%
	100.00%



KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautions

Net Wt. 20 lbs.  
(9.1 Kg.)

EPA Reg. No. 9198-46

EPA Est. No. 9198-0H-1

ANDERSONS  
WEED AND FEED  
XX-XX-XX

Contains 2, 4 D and MCPP

GUARANTEED ANALYSIS

Total Nitrogen (N) .

X 4% Ammonium Nitrogen

X 36 Urea Nitrogen

X 16 Coated Slow Release Urea Nitrogen

Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) .

X 4%

Soluble Potash (K<sub>2</sub>O) .

X 4%

Sulfur (S)

X 4%

Primary Plant Foods Derived from Ammoniated Phosphate, Urea, Murate of Potash and Buttar Coated Urea. Buttar derived from Sulfur Coated Urea. Potential Acidity 100 lb. of Calcium Carbonate Equivalent per ton.

Andersons Weed & Feed kills undesirable weeds and feeds your lawn to help it grow thick and green.

WEED CONTROL

2, 4 D for control of Broadleaf Plantain, Buckhorn Plantain, Dandelion, Purslane, Pigweed and Smartweed

MCPP for control of Common Chickweed, Mouse ear Chickweed, Red Clover, White Clover, Ground Ivy, Stichwort and Knotweed

For use on Turf except Desirable Clovers, Bent, St. Augustine, Dichondra and Carpet Grass

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Avoid contact with clothing, skin or eyes. Avoid breathing dust. Wash thoroughly after handling. Avoid contamination of food or foodstuffs. Do Not Hand Spread

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

WHEN TO APPLY

Use anytime weeds are actively growing and temperature is above 60°. Control is slower if weather is cool. Application in Spring and Fall provide most effective kill. Apply when lawn is wet from rain, dew or sprinkling. On newly seeded areas, wait until grass has been cut two or three times. Delay planting new grass for at least 3 weeks as 2, 4 D may temporarily inhibit seed germination or plant growth. Do not apply when turf is under stress from heat, moisture or drought. Do not over or under apply.

HOW TO APPLY

Apply evenly over entire lawn area at the rate of 1 1/4 lb per 1000 sq ft. Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Age and condition of spreader will affect application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. Fill spreader on driveway or sidewalk to eliminate chance of spillage on grass. Do not hand spread.

Rain, sprinkling or mowing within 24 hours may reduce effectiveness. Wait at least two weeks before making a second application.

IMPORTANT

Fill of dust or vapor may cause injury to nearby susceptible plants such as vegetables, flowers, fruit trees and ornamentals. Avoid contact with flower beds or shrubs. If accidentally deposited on flowers or shrubs hose off immediately. Do not use on crops that will be used for food or forage purposes. Do not graze animals on treated areas. Do not contaminate water used for irrigation or domestic purposes.

SUGGESTED SPREADER SETTINGS

Gro Wall Drop	4 1/2	Gro Wall Broadcast	1 1/2 OZ
Cyclone R1-84	4 5	Lely Type W	5 1/2
Scotts Drop PE?	4 1/2	Scotts Broadcast R21M	1 1/2 OZ
Scotts Professional Broadcast R7X	1 1/2 F		

NOTE: Above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Calculate the size of area to be treated and do not over or under apply. Close spreader when filling, stopping or turning. DO NOT HAND SPREAD.

STORAGE & DISPOSAL

Do not store near feed, foodstuffs, seeds, fertilizer, insecticides or fungicides. Store in a dry place. Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag. Discard bag in trash.

WARRANTY STATEMENT

"Seller warrants that this material conforms in its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty including the warranty of fitness for a particular purpose, and no agent or seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made."

Manufactured by

FAT FLOW FERTILIZER  
P.O. BOX 119, MAUMEE, OHIO 43537