

The contents of this bag
treats 5000 sq. ft.

TO OPEN:
Pull Narrow Tape



ANDERSONS WEED KILLER

Contains 2, 4-D and MCPP

HERBICIDE

9000-47

Use only on lawns, lawns, and lawns.
Do not use on lawns, lawns, and lawns.
Do not use on lawns, lawns, and lawns.
Do not use on lawns, lawns, and lawns.

0.44%
0.84%

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
See Back Panel for Additional Precautions

Net Wt. 15 lbs.

Net Wt. 15 lbs.
(6.8 Kg.)

ANDERSONS WEED KILLER

Contains 2, 4-D and MCPP

WEED CONTROL

2, 4-D for control of Broadleaf Plantain, Buckhorn Plantain, Dandelion, Purslane, Pigweed and Smartweed

MCPP for control of common Chickweed, Mouse-ear Chickweed, Red Clover, White Clover, Ground Ivy, Stitchwort, and Knotweed

For Use on Turfs except Desirable Clovers, Bent, St. Augustine, Dichondra and Carpet Grass

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals.

CAUTION

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Avoid contamination of feed or foodstuffs. **Do Not Hand Spread**

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHEN TO APPLY:

Use anytime weeds are actively growing and temperature is above 60°. Control is slower if weather is cool. Applications in Spring and Fall provide most effective kill. Apply when lawn is wet from rain, dew or sprinkling. On newly seeded areas, wait until grass has been cut two or three times. Delay planting new grass for at least 3 weeks as 2, 4-D may temporarily inhibit seed germination or plant growth. **Do not apply** when turf is under stress from heat, moisture, or drought.

HOW TO APPLY

Apply evenly over entire lawn area at the rate of 3 lbs. per 1000 sq. ft. Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Age and condition of spreader will affect application. Adjust spreader if necessary. Be sure to shut off spreader when turning or stopping to prevent accidental over application and burning of grass. **Do not hand spread.**

Rain, sprinkling, or mowing within 24 hours may reduce effectiveness. Wait at least 2 weeks before making a second application.

IMPORTANT:

Drift of dust or vapors may cause injury to nearby susceptible plants such as vegetables, flowers, fruit trees and ornamentals. Avoid contact with flower beds or shrubs. If accidentally deposited on flowers or shrubs, hose off immediately. **Do not use on crops that will be used for food or forage purposes.** Do not graze animals on treated areas. Do not contaminate water used for irrigation or domestic purposes.

SUGGESTED SPREADER SETTINGS

Scott's B-1	Scott's G-1
Ornitho 100	Scott's #15
Ornitho 100	Sears

STORAGE AND DISPOSAL

Do not store near feed, foodstuffs, seeds, fertilizer, insecticides or fungicides. **Store in dry place.** Do not reuse container. Wrap container and put in trash collection.

WARRANTY STATEMENT

Seller warrants that this material conforms to the formula description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty, including the warranty of fitness for a particular purpose, and the agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Manufactured by FREE FLOW FERTILIZER P.O. Box 119, Maumee, Ohio 41537

EST. NO. 1108 (04)