
The contents of this bag treats up to 6,000 sq. ft.

For Immediate Seeding of Bluegrass, Perennial Ryegrass and Fescue.



ACCEPTED

OCT 8 - 1986

Under the Federal Insections Fungicide, and Rodestinide Act, as amended, for the peciale tered under

THE ANDERSONS PRE-EMERGENT **CRABGRASS KILLER PLUS FERTILIZER**

XX-XX-XX

Contains 3.5% Siduron HERBICIDE

TIVE INGREDIENT	
Siduron (1-(2-Methylcyclohexyl) 3-phenylurea)	•
ERT INGREDIENTS 96 54	•
77AL 100 04	٠
GUARANTEED FERTILIZER ANALYSIS	
lal Ndrogen (N)	4
XX% Ammonacal Nitrogen	
#Heble Phosphoric Acid (P ₂ O ₆)	٨
Huble Potesh XX	À
mary Plant Foods Derived From	
many Plant Foods Derived From	

Potential Acidity xxx pounds Calcium Carbonate Per Ton

KEEP OUT OF THE REACH OF CHILDREN **CAUTION** See Back Panel for Additional Precautions

> Net Wt. 20 lbs. (9.1 Kg.)

EPA Reg No 9198 50

EPA Est. Nos. 9198 OH 1. 8198 NJ 1/0, Superscript used is first letter of run chide on end of container

THE ANDERSONS PRE-EMERGENT CRABGRASS KILLER PLUS FERTILIZER

XX-XX-XX

ADVANTAGES

- Keeps out Crabgress, Foxtell and Bernyard Grass by killing the sprouting seed
 May be used on freshly seeded Bluegress, Perennial Ryagrass, Feacus and certain other gras
 Easy to apply with standard seed and fertilizer apreaders

KEEP OUT OF REACH OF CHILDREN CUATION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ed. Avoid contact with skin, eyes and clothing. Wash thoroughtly after using. Avoid breathing dust. Do not co water used for trigetion or domestic purposes. Do not contaminate feed, loodstuffs or ponds. Do not store near seed, other fertilizers, insecticides or fungicides. Do not apply to vegetable crops.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of takes, streams and ponds. Do not conteminate water by cleaning of equipment or disposal of wastes. Drift of dust may cause injury to nearby susceptible plants.

DIRECTIONS FOR USE

It is a violation of Faderal Law to use this product in a menner inconsistent with its labeling

This product will control weedgrasses such as Crabgrasa (smooth and helry). Fostall and Barnyerd Grass in newly seeded or established plantings of a Bluegrass, Fescue, Perennial Ryegrass, Fledtop, Smoothbrome, Orchardgrass, Zoysla and the following strains of Bentgrass: Penncross, Seaside, Highland, Astoria, Miniella, C-1, C-19. This product is not for use on other Bentgrass strains, Bermuda Grass, vegetable or Brown new forms. or flower gardens.

WHEN TO APPLY

Apply prior to germination of weed grasses in the spring. Pracise time of application for a specific area will vary due to climate and individual cultural practices. Generally application during February or March will give best results, although control can be expected by applications—up to the time that Forsythia blooms fall. If in doubt, check with your Agricultural Extension Service. For newly seeded areas, apply se the feel

HOW TO APPLY

For best results, this product must be applied with no skips or misses. Determine the correct amount for your lawn area. For freshly seeded grass apply at the rate of 3.3 pounds per 1,000 equare feet. For established turl apply at 6.6 pounds per 1,000 equare feet. The best results are obtained by using the two-step procedure at the recommended half rate settings. Apply half the required amount in one direction and the other half at right angles to the first application. This procedure will minimize the chances of color streaking and erratic crabgress control. Use the suggested apreader settings as a guide, being sure to check the application rate at the start of each application. At least 1/2 inch of water must be provided by rainfall or watering within 3 days of treatment in order to carry the chemical into the root cone of germinating needs.

SUGGESTED SPREADER SETTINGS

SPREADER	Established Lawns 6.6 lbs:/1,000 sq. ft.		New Lawne 2.3 lbs./1,000 sq. ft.	
	Normal Plate	Half Rate	Normal Rate	Half Rate
Central	81/2	7	7	•
Central Broadcast	1 5	4	1	314
Cyclone	5%	414	414	4
Greenview	81/4	7	7	•
Ortho Drop	71/0	•		5
Ortho Broadcard	j 3	1	1 1	Not Recommended
Scotts Drop	1 •	6%	644	•
Scotts Broadcast	l a	F	F	} 0
Sears Drop	l i	•		7
Seers Broadcast	l š	4	4	3%

NOTE: Above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. This bag should cover approximately 3,000 square feet (30" x 100") on new fewns, Calculate the size of the area to be treated and do not over or under apply. Close apreader when fitting stopping or turning. DO NOT HANO SPREAD,

STORAGE AND DISPOSAL

Store in a dry place. Securely wrap original conteiner in several layers of newspaper and discard in trash. Do not reuse bag. Discard

"Seller warrants that this material conforms to its chemical description and is reasonably it for the purposes stated on the label when used In accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty including the warranty of liness for a particular purpose, and arrand to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made."