An Section 1997

FEB 1 1995

Mr. Daniel Paradiso, Jr.
The Andersons Lawn Fertilizer Division
DBA Free Flow Fertilizer
P.C. Box 119
Maumee, On 43537

bear Mr. Paradiso:

Subject: Easy Weeder Flower and Garden Weed Preventer EPA Registration No. 9198-60 Application Dates Haron 25, 1994, Request To Market as Either Homeowner or Commercial, Compliance with PR Rotice 93-3, MPS Exclusionary Statements and Format Revisions

The proposed labeling submitted with your application dated March 25, 1994 has been reviewed and found acceptable under the rederal Insecticise, Eungicide and Rodenticise Act (FIFRA) as amended, provided that:

e submit one (1) copy of the final printed label before releasing the product for shipment under the subject labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section o(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copies of the labeling are enclosed for your records.

Sincerely yours,

Joanne I. Hiller Product Manager (23) Fungicice-Herbicide Branch Registration Division (7505C)

Enclosure

E.milson:Diskette ABC17:02-01-95

BEST AVAILABLE COPY

) -1 h

9198-60H Page 1 of 5

(FRONT PANEL)

HOMEOWNER USE ONLY

ACCEPTED

with COMMENTS
for Comments

BASY WEEDER

FEB | 1995

FLOWER AND GARDEN WEED PREVENTER

To be the Lotter of Concess, Line wide, and there is a many time for the posterior registered order EDA Re. No.

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT LBS.

TREATS __ SQ. FT.

EPA Reg. No. 9198-60

EPA Est. 9198 OH-1

•••

9198-60H Page 2 of 5

(BACK PANEL)

HOMEOWNER USE ONLY

NET WEIGHT __ LBS.

TREATS SQ. FT.

EASY WEEDER

FLOWER AND GARDEN WEED PREVENTER

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to tish. Do not apply to water. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by disposing of equipment washings.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

HOMEOWNER USE ONLY

EASY WEEDER PREVENTS THE FOLLOWING WEEDS:

Annual Grassy Weeds: Annual Bluegrass, Barnyard Grass (Watergrass), Brachiaria, Bromegrass, Cheat, Crabgrass, Foxtail, Goosegrass, Johnsongrass (from seed). Junglerice, Sandbur, Stinkgrass, Sprangletop, Texas Panicum, Wildcane.

Annual Broadleaf Weeds: Carelessweed, Carpetweed, Chickweed, Florida Purslane (Pusley), Goosefoot, Knotweed, Kocchia, Lambsquarter, Pigweed (Spiney and Redroot), Purslane, Russian Thistle, Stinging Nettle.

USE EASY WEEDER AS DIRECTED FOR THE FOLLOWING:

Vegetables from Seed: cantaloupes, cucumbers, carrots, okra, southern or cow peas, field peas, black-eyed peas, lima beans, snap beans.

Vegetables before Transplanting: (2-3" Tall) broccoli, brussel sprouts, cabbage, cauliflower, peppers, tomatoes.

Vegetables after Transplanting: (2-3" Tall) Irish potatoes

Established Flowers: (2-3" Tall) Achillea, African Daisy, Ageratum, Arcototis, Aster (Perennial), Balsam, Blackeyed Susan, Calendula, California Poppy, Calliopsis, Carnation, Centaurea, Chrysanthemum, Coreopsis, Cornflower, Cosmos, Dahlia, Dianthus, Dimorphetheca, Dusty Miller, Floss Flower, Forget-Me-Not, Four O'Clock, Gaillardia, Gladiolus (Corms at least 1"diameter), Hollyhock, Impatiens, Ixora (Coccinea), Lobelia, Lupines, Marigold, Morning Glory, Nasturtium, Nicotania, Periwinkle, Petunia, Phlox, Portulaca, Rudbeckia, Rose, Salvia, Scabiosa, Shasta Daisy, Snapdragon, Stocks, Snow On The Mountain, Sunflower, Sweet Alyssum, Sweet Pea, Sweet William, Vinca, Yarrow, Zinnia.

Groundcovers: Aaronsbeard, Asparagus Fern, Bellflower (Adriatic), Bellflower (Poscharsky), Bigblue Lilyturf, Ceanothus, Coreopsis, Cotoneaster, Coyote Brush, Crown Vetch, Daisy Trailing African, Gazania, Germander, Largeleaf Ice Plant, Ivy (Algerian), Ivy (English), Lily-of-the-Nile, Marigold, Myoporum, Plumbago (Dwarf), Rockrose, Rosemary, Rupturewort, Snow-in-Summer, Speedwell, St. Johnswort, Stonecrop (Sedum), Strawberry (Beach), Thrift, Verbena, Wirevine (Creeping), Yarrow (Woolly), Zoysiagrass

Shrubs and Trees: Use around most common varieties for season-long woed prevention.

9198-60H Page 4 of 5

(BACK PANEL)

HOMEOWNER USE ONLY

When to Apply: Early application is best for easy weed control all season long. Or, apply any time around shrubs, trees and established plants. Before sowing listed vegetables by seed, apply product no earlier than 10 weeks prior to planting.


Use spreader in garden beds only. Do not use Easy Weeder Flower and Garden Weed Preventer on lawns. Never use when planting seed except for the vegetables listed. Do not use with flower seeds.

How to Apply: Since Easy Weeder works by preventing weed seed germination, existing weeds must be removed before applying. Avoid over-application to prevent carryover problems with sensitive plants the following season. Do not use in enclosed areas.


Established Plants, at lease 2-3" Tall: Flowers, listed vegetables, ornamentals and trees: sprinkle on soil surface (wearing gloves or with a spreader) to match Example A (this is 1 oz. per 10 sq. ft.)

Before Planting: Prior to transplanting 2-3" listed flowers, or, less than 10 weeks prior to planting vegetable seed; apply in medium soil to match Example B and mix in (this equals 1 oz. per 60 sq.ft.) Mix into soil with rake or roto-tiller 1-3" deep. Place transplant. Do not apply after transplanting, except on Irish potatoes. (See chart below for light [sandy] or heavy [clay] soil.)

Example A



Example B



Application Rates for Mixing into Soil:

Soil Texture Rates*	Light (Sandy)	Medium	Heavy (Clay)
Use 1 ounce per	80 sq. ft.	60 sq. ft.	40 sq. ft.
Cr 1 cup per	1,280 sq. ft.	960 sq. ft.	640 sq. ft.

^{*1} cup (tightly packed) equals % pound.

HOMEOW. TER USE ONLY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: Free Flow Fertilizer

P. O. Box 119 Maumee, OH 43537

Date Printed: 01/24/94

Date Approved: Supersedes: Label Change:

9198-60C Page 1 of 5

(FRONT PANEL)

COMMERCIAL USE ONLY

APTEPLE.

EASY WEEDER

FLOWER AND GARDEN WEED PREVENTER

EEB 1 20%

ACTIVE INGREDIENT:	9	198	-60.
Trifluralin:alpha, alpha, alpha-trifluoro-1, 6-dinitro-N, N-dipropyl-o-toluidine	. ,		1.47%
INERT INGREDIENTS			98.53%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT __ LBS.

TREATS ____ SQ. FT.

EPA Reg. No. 9198-60

EPN Ect. 9198-CH-1

: . .

COMMERCIAL USE ONLY

NET WEIGHT LBS.

TREATS ____ SQ. FT.

EASY WEEDER

FLOWER AND GARDEN WEED PREVENTER

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by disposing of equipment washings.

DIRECTIONS FOR USE

It is a violation ϵ Federal Law to use this product in a manner incons stent with its labeling.

Not for use on plants being grown for sale or other commercial use, or, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamertal gardens or parks, or on golf courses or lawns.

COMMERCIAL USE ONLY

EASY WEEDER PREVENTS THE FOLLOWING WEEDS:

Annual Grassy Weeds: Annual Bluegrass, Barnyard Grass (Watergrass), Brachiaria, Bromegrass, Cheat, Crabgrass, Foxtail, Goosegrass, Johnsongrass (from seed). Junglerice, Sandbur, Stinkgrass, Sprangletop, Texas Panicum, Wildcane.

Annual Broadleaf Weeds: Carelessweed, Carpetweed, Chickweed, Florida Purslane (Pusley), Goosefoot, Knotweed, Koccnia, Lambsquarter, Pigweed (Spiney and Redroot), Purslane, Russian Thistle, Stinging Nettle.

USE EASY WEEDER AS DIRECTED FOR THE FOLLOWING:

Vegetables from Seed: cantaloupes, cucumbers, carrots, okra, southern or cow peas, field peas, black-eyed peas, lima beans, snap beans.

Vegetables before Transplanting: (2-3" Tall) broccoli, brussel sprouts, cabbage, cauliflower, peppers, tomatoes.

Vegetables after Transplanting: (2-3" Tall) Irish potatoes

Established Flowers: (2-3" Tall) Achillea, African Daisy, Ageratum, Arcototis, Aster (Perennial), Balsam, Blackeyed Susan, Calendula, California Poppy, Calliopsis, Carnation, Centaurea, Chrysanthemum, Coreopsis, Cornflower, Cosmos, Dahlia, Dianthus, Dimorphetheca, Dusty Miller, Floss Flower, Forget-Me-Not, Four O'Clock, Gaillardia, Gladiolus (Corms at least 1"diameter), Hollyhock, Impatiens, Ixora (Coccinea), Lobelia, Lupines, Marigold, Morning Glory, Nasturtium, Nicotania, Periwinkle, Petunia, Phlox, Portulaca, Rudbeckia, Rose, Salvia, Scabiosa, Shasta Daisy, Snapdragon, Stocks, Snow On The Mountain, Sunflower, Sweet Alyssum, Sweet Pea, Sweet William, Vinca, Yarrow, Zinnia.

Groundcovers: Aaronsbeard, Asparagus Fern, Bellflower (Adriatic), Bellflower (Poscharsky), Bigblue Lilyturf, Ceanothus, Coreopsis, Cotoneaster, Coyote Brush, Crown Vetch, Daisy Trailing African, Gazania, Germander, Largeleaf Ice Plant, Ivy (Algerian), Ivy (English), Lily-of-the-Nile, Marigold, Myoporum, Plumbago (Dwarf), Rockrose, Rosemary, Rupturewort, Snow-in-Summer, Speedwell, St. Johnswort, Stonecrop (Sedum), Strawberry (Beach), Thrift, Verbena, Wirevine (Creeping), Yarrow (Woolly), Zoysiagrass

Shrubs and Trees: Use around most common varieties for season-long weed prevention.

9198-60C Page 4 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

When to Apply: Early application is best for easy weed control all season long. Or, apply any time around shrubs, trees and established plants. Before sowing listed vegetables by seed, apply product no earlier than 10 weeks prior to planting.

Use spreader in garden beds only. Do not use Easy Weeder Flower and Garden Weed Preventer on lawns. Never use when planting seed except for the vegetables listed. Do not use with flower seeds.

How to Apply: Since Easy Weeder works by preventing weed seed germination, existing weeds must be removed before applying. Avoid over-application to prevent carryover problems with sensitive plants the following season. Do not use in enclosed areas.

Established Plants, at lease 2-3" Tall: Flowers, listed vegetables, ornamentals and trees: sprinkle on soil surface (wearing gloves or with a spreader) to match Example A (this is 1 oz. per 10 sq. ft.)

Before Planting: Prior to transplanting 2-3" listed flowers, or, less than 10 weeks prior to planting vegetable seed; apply in medium soil to match Example B and mix in (this equals 1 oz. per 60 sq.ft.) Mix into soil with rake or roto-tiller 1-3" deep. Place transplant. Do not apply after transplanting, except on Irish potatoes.

(See chart below for light [sandy] or heavy [clay] soil.)

Example A

Example B

* *

Application Rates for Mixing into Soil:

Soil Texture Rates*	Light (Sandy)	Medium	Heavy (Clay)
Use 1 ounce per	80 sq. ft.	60 sq. ft.	40 sq. ft.
Or 1 cup per	1,280 sq. ft.	960 sq. ft.	640 sq. ft _. .

^{*1} cup (tightly packed) equals % pound.

9198-60C Page 5 of 5

(BACK PANEL)

COMMERCIAL USE ONLY

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OF IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: Free Flow Fertilizer P. O. Box 119

Maumee, OH 43537

Date Printed: 01/24/94

Date Approved: Supersedes: Label Change: