

Net Wt. 10 Lbs. (4.5 Kg.)

Treats 5,000 Sq. Ft.

TURF CARE Lawn and Garden Insecticide

5% DIAZINON

ACTIVE INGREDIENTS
5.0% DIAZINON
INERT INGREDIENTS
95.0%

BY WEIGHT
5.0%
95.0%

ACC
JAN 28 1981
U.S. Patent Office
EPA Reg. No. 9199-62

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

If swallowed - May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply respiration if indicated.

If on skin - Remove or contaminate clothing and wash affected areas with soap and water.

If in eyes - Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

Toxic to Fish - In the presence of signs of photosensitivity, fish present atropine anticholinergic 2-PAM is a so antidote and may be administered in conjunction with it.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are active by visiting the crop, flower crop, or weeds blooming in the treated area. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

This is a general label for the use of this product. Use a more specific label with the following:

Turf Care Lawn and Garden Insecticide controls the following lawn insects on grass and Dichondra lawns:
Ants, Armyworms, Bermuda Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Hyperodes Weevils (New York only), Lawnmoths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs, Springtails, (Columbian).

Do not hand spread. Apply Turf Care Lawn and Garden Insecticide to your entire lawn in a 5 foot by 3 foot application around or near the foundation to control these insects which may enter a house at a rate of 2 lbs. per 1,000 sq. ft. of area (an area 20 by 50 or 32 oz. per 100 sq. ft. of area) to 1/4 inch. For heavy textured lawns such as St. Augustine use 3 lbs. per 1,000 sq. ft. (or 4.8 oz. per 100 sq. ft.). See back panel for recommended spreader settings. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

Turf Care Lawn and Garden Insecticide also controls Billbugs and White Grubs of Japanese Beetle, European Chaffer, Southern Chaffer and Atractodes spretulus (Dung Beetle). Apply at 2.5 lbs. per 1,000 sq. ft. For Billbugs apply when activity is first noticed in lawn. For Atractodes and Japanese Beetle damage, apply when damage is first noticed. For Grub control, including Dung Beetle, treat any time between late July and early October. For Dung Beetle Grubs, apply in spring or early summer when damage is first noticed. For Japanese Beetle water grass with 1/4 inch of water after application. Repeat treatment in late summer or early fall if a second generation is in evidence. For Hyperodes Weevils, apply at a rate of 2 lbs. per 1,000 sq. ft. in mid April to mid May, watering thoroughly after each application. See back panel for recommended spreader settings.

Turf Care Lawn and Garden Insecticide can be used for localized control of ants and Digger Wasps. Apply 1/2 teaspoon over and around each nest. For a digger wasp nest, wash ground nest opening. Apply to wasp nests a hard dust when activity is low. For ants, apply directly to the nest. Apply Turf Care Lawn and Garden Insecticide as listed for crops below. Apply just before planting and immediately water to the specified depth.

Cutworms: Alfalfa and Beans, Corn, Beans, Cabbage, Cantaloupe, Cauliflower, Celery, Cucumber, Cucumbers, Endive, Escarole, Kale, Lettuce, Lima Beans, Mung Beans, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash.

Use 2 1/2 lbs. per 1,000 sq. ft. of area to 1/4 inch depth (2 1/2 lbs. for control of surface cutworms and 3 1/2 lbs. for control of cutworms that burrow into the soil).

Mole Crickets: Beans, Corn, Cabbage, Cauliflower, Celery, Cucumber, Cucumbers, Kale, Lettuce, Mustard, Peppers, Radishes, Swiss Chard, Tomatoes, Turnips.

Use 4 1/2 lbs. per 1,000 sq. ft. of area to 1/4 inch depth.

Wireworms: Beans, Cabbage, Cantaloupe, Cauliflower, Celery, Cucumber, Cucumbers, Kale, Lettuce, Lima Beans, Mung Beans, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Swiss Potatoes, Tomatoes, Turnips, Watermelons, Winter Squash.

Use 2 1/2 lbs. per 1,000 sq. ft. of area to 1/4 inch depth.

STORAGE AND DISPOSAL

Keep this product in its original container until it is used. Do not reuse empty containers. Do not contaminate water by cleaning of equipment or disposal of wastes.

WARRANTY STATEMENT

See label for full details. This product is warranted to control the insects listed on the label when used as directed. If the product does not control the insects listed on the label, the user may return the product to the manufacturer for a refund of the purchase price. This warranty is void if the product is used in a manner not specified on the label. The manufacturer is not responsible for damage to property or crops caused by the use of this product in a manner not specified on the label.



Manufactured by The Andersons Lawn Fertilizer Division • P.O. Box 119 Maumee, Ohio 43537

EPA Reg. No. 9199-62
EPA Est. No. 77-111

SUGGESTED SPREADER SETTINGS


For control of:

Ants, Armyworms, Bermuda Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Hyperodes Weevil (New York only), Lawnmoths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs, Springtails (Collembola).

MODEL	TURF TYPE	
	FINE TO MEDIUM TEXTURE 2 LB./1,000 SQ. FT.	HEAVY (ST. AUGUSTINE) 3 LB./1,000 SQ. FT.
Central	7-1/2	8
Cyclone	3-1/4	3-3/4
Greenview	7-1/2	8
Ortho 3000	7-1/2	8-1/2
Ortho 4000	3/4	1-1/4
Scotts Drop	4	5
Scotts Broadcast	A-1/2	B-1/2
Sears	9	10

For control of Billbugs and White Grubs:

MODEL	2.5 LB./1,000 SQ. FT.
Central	7-3/4
Cyclone	3-1/2
Greenview	7-3/4
Ortho 3000	8
Ortho 4000	1
Scotts Drop	4-1/2
Scotts Broadcast	B
Sears	9-1/2



The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of terrain. Calculate the size of the area to be treated and do not over-or under-apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader settings, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning. Do not hand spread.

Important: See front panel for complete use directions.