



9198-65

02/02/1982

1/2



ACCEPTED  
FEB - 2 1982

9198-65

# Lofts

## Crabgrass Preventer

### With Tupersan

**A PRE-EMERGENCE HERBICIDE FOR THE CONTROL OF COMMON CRABGRASS IN ESTABLISHED AND NEWLY SEEDING LAWNS**

Treats 2,000 Sq. Ft. (Established Lawns)  
Treats 4,000 Sq. Ft. (Newly Seeded Lawns)



<b>ACTIVE INGREDIENT</b>	
Siduron [1-(2-Methylcyclohexyl)-3-Phenylurea]	4.70%
<b>INERT INGREDIENTS</b>	95.30%
<b>TOTAL</b>	<b>100.00%</b>

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

**Net Weight 10 LBS. (4.5 kg)**

LFTCP10

12-81

Net Weight 10 LBS. (4.5 kg)



2/a

# Crabgrass Preventer

With \*Tupersan®

### GUARANTEED ANALYSIS

ACTIVE INGREDIENT	4.70%
Siduron (1-(2-Methylcyclohexyl)-3-Phenylurea)	4.70%
INERT INGREDIENTS	95.30%
TOTAL	100.00%

\*Tupersan® Reg. T.M. of E.I. DuPont de Nemours Co., Inc. for Siduron weed killer.

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate feed, foodstuff, or ponds. Do not store near seed, other fertilizers, insecticides, or fungicides. Do not apply to vegetable crops. **DO NOT HAND SPREAD.**

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift of dust may cause injury to nearby susceptible plants.

### GENERAL INFORMATION

LOFTS CRABGRASS PREVENTER is formulated to prevent the growth of both smooth and hairy crabgrass in either a newly seeded or well established lawn. It will also aid in controlling certain other annual weed grasses such as Foxtail and Barnyard grass. LOFTS CRABGRASS PREVENTER will not damage most of the commonly used lawn grasses even if applied at time of seeding.

#### IMPORTANT:

Use only on bluegrass, fescue, reedtop, smooth brome, perennial ryegrass, orchard grass, and the following strains of bentgrass: Penncross, Seaside, Astoria, Nirmella, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (Poa annua), clover, or most broadleaf weeds.

### DIRECTIONS FOR USE

Apply to the lawn during the spring of the year prior to the germination of crabgrass. Best results are obtained when application is made approximately 2-3 weeks before crabgrass appears. It is, therefore, suggested that application be avoided during February or early March.

#### NEWLY SEED LAWNS

1. Apply contents evenly over 4,000 square feet immediately after seeding (2½ lbs. per 1,000 sq. ft.). For best results repeat applications in 4 weeks.
2. Provide sufficient moisture to insure germination of new grass seed.
3. An application of Lofts Turf Food is recommended after new turf is growing well.

#### ESTABLISHED, OVERSEEDED OR PATCHED LAWNS

1. Apply contents evenly over 2,000 square feet before crabgrass germinates (5 lbs. per 1,000 sq. ft.).
2. For best results, the lawn should be watered thoroughly within 3 days after application.
3. Before application, lawn should be raked to remove extraneous material and insure uniform application. In order to obtain expected results, CRABGRASS PREVENTER should be evenly applied with no skips or misses.

NOTE—DO NOT RAKE LAWN or DISTURB SOIL SURFACE after application. Application of Lofts Turf Food is recommended in the Summer and Fall.

### SUGGESTED SPREADER SETTINGS

	2000 Sq. Ft.	4000 Sq. Ft.
Central	8½-10½	8½-9
Central Broadcast	8½-8½	4½-5
Cyclone	6-6½	5½-5½
Greenview	9½-10½	8½-9
Ortho 3000	11½-12½	9½-10½
Ortho 4000	7-8½	3-4½
Scotts Drop	12½-13½	9-10
Scotts Broadcast	L-M½	H½-J
Sears	12-13	10-11
Sears Broadcast	5½-5½	4½-5

These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

Be sure to calibrate your spreader with each application.

### STORAGE AND DISPOSAL

Do not store near feed, foodstuff, seeds, fertilizers, insecticides, or fungicides. Store in dry place. Do not reuse container. Wrap container and put in trash collection.



Manufactured For  
**LOFTS SEED, INC.**  
CHIMNEY ROCK ROAD