

1/9

(FRONT PANEL)

HOMEOWNER USE ONLY

THE ANDERSONS

CRABGRASS PREVENTER

WITH 4.7% TUPERSAN

A PRE-EMERGENCE HERBICIDE FOR THE CONTROL OF COMMON CRABGRASS IN ESTABLISHED AND NEWLY SEEDED LAWNS.

**ACTIVE INGREDIENT:**

Siduron (1-(2-Methylcyclohexyl)-3-Phenylurea) ..... 4.70%

INERT INGREDIENTS ..... 95.30%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 10 LBS (4.5 KG)

TREATS 2,000 SQ. FT. (ESTABLISHED LAWNS)

TREATS 4,000 SQ. FT. (NEWLY SEEDED LAWNS)

EPA Reg. No. 9198-65

**ACCEPTED**  
MAY 26 1994  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 9198-65

EPA-Reg. 9198-OH-1

(BACK PANEL)

HOMEOWNER USE ONLY

Net Weight 10 Lbs. (4.5 kg)

THE ANDERSONS

CRABGRASS PREVENTER  
with 4.7% Tupersan

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate feed, foodstuff, or ponds. Do not store near seed, other fertilizers, insecticides, or fungicides. Do not apply to vegetable crops. DO NOT HAND SPREAD.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to water. Do not contaminate water when disposing of equipment washwaters. Drift of dust may cause injury to nearby susceptible plants.

GENERAL INFORMATION

This product is formulated to prevent the growth of both smooth and hairy crabgrass in either a newly seeded or well established lawn. It will also aid in controlling certain other annual weed grasses such as Foxtail and Barnyard grass. This product will not damage most of the commonly used lawn grasses even if applied at time of seeding.

IMPORTANT:

Use only on bluegrass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross, Seaside, Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (Poa annua), clover, or most broadleaf weeds.

(BACK PANEL)

HOMEOWNER USE ONLY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

Apply to the lawn during the spring of the year prior to the germination of crabgrass. Best results are obtained when application is made approximately 2-3 weeks before crabgrass appears. It is, therefore, suggested that application be avoided during February or early March.

NEWLY SEED LAWNS

- 1. Apply contents evenly over 4,000 square feet immediately after seeding (2 1/2 lbs. per 1,000 sq ft.). For best results repeat applications in 4 weeks.
- 2. Provide sufficient moisture to insure germination of new grass seed.


ESTABLISHED, OVERSEEDED OR PATCHED LAWNS

- 1. Apply contents evenly over 2,000 square feet immediately after seeding (5 lbs. per 1,000 sq. ft.).
- 2. For best results, the lawn should be watered thoroughly within 3 days after application.
- 3. Before application, lawn should be raked to remove extraneous material and insure uniform application. In order to obtain expected results, this product should be evenly applied with no skips or misses.

NOTE-DO NOT RAKE LAWN or DISTURB SOIL SURFACE after application. Application of a turf food is recommended in the Summer and Fall.

SUGGESTED SPREADER SETTINGS

(AVAILABLE FOR FINAL PRINTING)



(BACK PANEL)

HOMEOWNER USE ONLY

NOTE: These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.


WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By: The Andersons  
Lawn Fertilizer Division  
P.O. Box 119  
Maumee, OH 43537

Date Printed: 01/24/94  
Date Approved:  
Supersedes:  
Label Change:



5/9

(FRONT PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS

CRABGRASS PREVENTER

WITH 4.7% Tupersan

A PRE-EMERGENCE HERBICIDE FOR THE CONTROL OF COMMON CRABGRASS IN ESTABLISHED AND NEWLY SEEDED LAWNS.

ACTIVE INGREDIENT:

Siduron (1-(2-Methylcyclohexyl)-3-Phenylurea)	.....	4.70%
INERT INGREDIENTS	.....	95.30%
	TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 10 LBS (4.5 KG)

TREATS 2,000 SQ. FT. (ESTABLISHED LAWNS)  
TREATS 4,000 SQ. FT. (NEWLY SEEDED LAWNS)

EPA Reg. No. 9198-65

**ACCEPTED**  
MAY 26 1994  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 9198-65

EPA Est. 9198-OH-1

619

(BACK PANEL)

COMMERCIAL USE ONLY

Net Weight 10 Lbs. (4.5 kg)

THE ANDERSONS

CRABGRASS PREVENTER  
with 4.7% Tupersan

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS


Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate feed, foodstuff, or ponds. Do not store near seed, other fertilizers, insecticides, or fungicides. Do not apply to vegetable crops. DO NOT HAND SPREAD.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift of dust may cause injury to nearby susceptible plants.

GENERAL INFORMATION

THE ANDERSONS CRABGRASS PREVENTER is formulated to prevent the growth of both smooth and hairy crabgrass in either a newly seeded or well established lawn. It will also aid in controlling certain other annual weed grasses such as Foxtail and Barnyard grass. This product will not damage most of the commonly used lawn grasses even if applied at time of seeding.



(BACK PANEL)

COMMERCIAL USE ONLY

IMPORTANT:

Use only on bluegrass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross. Seaside, Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (Poa annua), clover, or most broadleaf weeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

Apply to the lawn during the spring of the year prior to the germination of crabgrass. Best results are obtained when application is made approximately 2-3 weeks before crabgrass appears. It is, therefore, suggested that application be avoided during February or early March.

NEWLY SEED LAWNS

1. Apply contents evenly over 4,000 square feet immediately after seeding (2 1/2 lbs. per 1,000 sq ft.). For best results repeat applications in 4 weeks.
2. Provide sufficient moisture to insure germination of new grass seed.

ESTABLISHED, OVERSEEDED OR PATCHED LAWNS

1. Apply contents evenly over 2,000 square feet immediately after seeding (5 lbs. per 1,000 sq. ft.).
2. For best results, the lawn should be watered thoroughly within 3 days after application.
3. Before application, lawn should be raked to remove extraneous material and insure uniform application. In order to obtain expected results, product should be evenly applied with no skips or misses.

NOTE-DO NOT RAKE LAWN or DISTURB SOIL SURFACE after application. Application of a turf food is recommended in the Summer and Fall.

819

(BACK PANEL)

COMMERCIAL USE ONLY

**SUGGESTED SPREADER SETTINGS**

(AVAILABLE FOR FINAL PRINTING)

**NOTE:** These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

Be sure to calibrate your spreader with each application.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.



9/9 ..

(BACK PANEL)

COMMERCIAL USE ONLY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by: The Andersons  
Lawn Fertilizer Division  
P.O. Box 119  
Maumee, OH 43537

Date Printed: 01/25/94  
Date Approved:  
Supersedes:  
Label Change:

