


9198-66

02/04/1982

1/2



CRABGRASS



ACCEPTED
FEB - 4 1982

LOFTS

LAWN FOOD PLUS
CRABGRASS
PREVENTER
25-3-3

2 IN 1



NATURAL ORGANIC CARRIER

HERBICIDE


ACTIVE INGREDIENTS:

N-butyl-n-ethyl-alpha, alpha, alpha-trifluoro-2, 6-Dinitro-p-Toluidine..... 1.2%

INERT INGREDIENTS 98.72%

TOTAL 100.00%

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
See Back Panel for Additional Precautions



NET WT. 18 LBS. (8.2 kg)
TREATS 5,000 SQ. FT.

OVER 55 YEARS OF QUALITY PRODUCTS

EPA Reg. No. 0190-60

LFTCP18
EPA Est. No. 0190 CH 1

At work -
back

6
5-10
1-10



NET WT. 18 LBS. (8.2 kg)

2/2



2 IN 1 LAWN FOOD PLUS CRABGRASS PREVENTER 25-3-3

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
23.8% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	3%
Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, and Murates of Potash.	
Potential Acidity 620 lbs. Calcium Carbonate Equivalent per ton.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs, or feed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by direct application, cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass, goosegrass, and other annual weed species in established:

BLUEGRASS (perennial)	FESCUE	BERMUDAGRASS	ST. AUGUSTINEGRASS
RYEGRASS (perennial)	BAHIAGRASS	CENTPEDEGRASS	ZOYAGRASS

GENERAL INFORMATION

LOFTS' 2 IN 1 LAWN FOOD PLUS CRABGRASS PREVENTER has the following advantages:

- 1. CONVENIENT & ECONOMICAL**—Loft's 2 In 1 is on a corn cob carrier, can be used effectively in a rotary type spreader as well as a drop type spreader.
- 2. GOOD SAFETY MARGIN**—Applied as directed, will not injure perennial Bluegrass, perennial Ryegrass, Fescue, Bermudagrass, Zoysiagrass, St. Augustinegrass, Centpede and Bahiagrass. Do not apply to new lawns started the previous fall.
- 3. ESTABLISHES A WEED CONTROL ZONE**—Loft's 2 In 1 establishes a weed control zone by killing certain annual weed seeds as they germinate.
- 4. WEEDS CONTROLLED**—Application prior to weed germination will prevent smooth and hairy crabgrass, annual bluegrass (goosegrass), goosegrass (silver crabgrass or crowfoot), watergrass (barnyardgrass), and yellow and green foxtail.
- 5. FEEDS YOUR LAWN**—Loft's 2 In 1 feeds your lawn a balanced diet of nitrogen, phosphorus, and potash.

WHEN TO APPLY

FOR CRABGRASS CONTROL: Apply in the spring before crabgrass germinates. Re-seeding should be delayed for 30 to 120 days after application. To insure satisfactory re-seeding results after using a pre-emergence chemical, the amount of seed should be increased.

FOR POA ANNUEA (ANNUAL BLUEGRASS) CONTROL: Apply late to early fall (mid-July to mid-September) prior to annual bluegrass first seed germination. Chemicals that effectively control Poa annua seeds may thin established Poa annua and fine-leaved fescues. Re-seeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY

Apply LOFTS' 2 IN 1 LAWN FOOD PLUS CRABGRASS PREVENTER evenly over the turf area without skips or misses. A even uniform application can be made by spreading half the required amount of this product over the area, and then repeating the spreading half at right angles or cross-wise. Both drop-type and rotary-type spreaders may be used to apply.

Listed below are suggested spreader settings. These settings are only approximates. Differences may occur due to amount of spreading material at which applied, and roughness of area treated. This bag should cover approximately 5000 sq. ft. (Brooming, 10 ft. or 100 ft. = 5000 sq. ft.) Calculate the size of area to be treated and do not over or under apply. Check spreader when filling, stopping, or turning. DO NOT RANG SPREAD.

SUGGESTED SPREADER SETTINGS

Model	Recommended Two-Step Settings (lb./row)	One Step Setting (lb./row) (recommended)
Control	7-7%	6-6%
Control Broadcast	4-4%	4-4%
Cyclone	4-4-4%	5-5%
Greenline	7-7%	6-6%
Orbit 2000	7-4-6%	5-4-5%
Orbit 4000	1-1%	2
Scotts Drop	4-4-5%	5-4-5%
Scotts Broadcast	D-E	F-G
Scotts	5-5	10-11
Scotts Broadcast	4-4%	4-4-4%

STORAGE AND DISPOSAL

Storage: Keep container closed and away from food, feedstuff, or domestic water supplies. Disposal: Do not reuse empty container. Wrap container and put in trash collection.

*Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty including the warranty of fitness for a particular purpose, and the extent of seller's liability is limited to the actual purchase price of the material as shown on the invoice. No agent shall collect a fee for any breach of warranty except the purchase price of the material as shown on the invoice.



Manufactured For
LOFTS SEED, INC.
CHIMNEY ROCK ROAD
BOUND BROOK, NEW JERSEY 08805