

9198-66

08/01/1988

1/5

(FRONT PANEL)


LOFTS

"2 in 1"

LAWN FOOD PLUS

CRABGRASS PREVENTER

25-3-3



ACTIVE INGREDIENT:

N-Butyl-n-ethyl-alpha,alpha,alpha-trifluoro-2,6-Dinitro-p-Toluidine.... 1.28%

INERT INGREDIENTS	98.72%
TOTAL	100.00%

KEEP OUT OF THE REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Weight 18 LBS. (8.2 kg)

TREATS 5,000 SQ. FT.

EPA Reg. No. 9198-66

EPA Est. 9198-OH-1



(Back Panel)

Net Weight 18 LBS. (8.2 kg)

LOFTS

"2 in 1"

LAWN FOOD

PLUS CRABGRASS PREVENTER

25-3-3

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
23.8% Urea Nitrogen	
Available Phosphoric Acid (P2O6)	3%
Soluble Potash (K2O)	3%
Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, and Muriate of Potash.	
Potential Acidity 620 lbs. Calcium Carbonate Equivalent per ton.	

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs, or feed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation Federal Law to use this product in a manner inconsistent with its labeling.

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass (poa annua), goosegrass, and other annual weed grasses in established:

- | | | | |
|------------------------|--------------|------------------|---------------------|
| (UEGRASS (perennial)✓ | FESCUE ✓ | BERMUDAGRASS | ST. AUGUSTINEGRASS✓ |
| RYEGRASS (perennial)✓ | BAHIAGRASS ✓ | CENTIFEDEGRASS ✓ | ZOYSIAGRASS ✓ |

GENERAL INFORMATION


LOFTS 2 IN 1 LAWN FOOD PLUS CRABGRASS PREVENTER has the following advantages:

1. CONVENIENT & ECONOMICAL - Loft's "2 in 1" is on a corn cob carrier, can be used effectively in a rotary-type spreader as well as a drop-type spreader.
2. GOOD SAFETY MARGIN - Applied as directed, will not injure perennial Bluegrass, perennial Ryegrass, Fescue, Bermudagrass, Zoysiagrass, St. Augustinegrass, Centipede and Bahiagrass. Do not apply to new lawns started the previous fall.
- (ESTABLISHES A WEED CONTROL ZONE - Loft's "2 in 1" establishes a weed control zone by killing certain annual weed seeds as they germinate.
4. WEEDS CONTROLLED - Application prior to weed germination will prevent smooth and hairy crabgrass, annual bluegrass (poa annua), goosegrass (silver crabgrass or crowfoot), watergrass (barnyardgrass) and yellow and green foxtail.
5. FEEDS YOUR LAWN - Loft's "2 in 1" feeds your lawn with nitrogen, phosphorous, and potash.

WHEN TO APPLY

FOR CRABGRASS CONTROL: Apply in the spring before crabgrass germinates. Reseeding should be delayed for 90 to 120 days after application. To insure satisfactory reseeding results after using a pre-emergence chemical, the amount of seed should be increased.

FOR POA ANNUA (ANNUAL BLUEGRASS) CONTROL: Apply late summer to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Reseeding should be delayed 12 to 18 weeks after application.



HOW TO APPLY

Apply LOFT'S "2 in 1" LAWN FOOD PLUS CRABGRASS PREVENTER evenly over the turf area without skips or misses. A more uniform application can be made by spreading half the required amount of this product over the area, and then applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used to apply.

Stated below are suggested spreader settings. These settings are only approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. This bag should cover approximately 5,000 sq.ft. (example, 50 ft. x 100 ft. = 5000 sq.ft.). Calculate the size of area to be treated and do not over or under apply. Close spreader when filling, stopping or turning. DO NOT HAND SPREAD.

SUGGESTED SPREADER SETTINGS

(AVAILABLE FOR FINAL PRINTING)

These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

