



1682
1723
9198-75

Tee TimeTM Fertilizer 20-4-10 with RONSTAR[®]

with comment
FEB 15 1985
9198-75

the professional's partnerTM

The Andersons

ACTIVE INGREDIENT
Oxadiazin (2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-6,7,8-triazolo[4,5-d]pyrimidin-5-ylidene-3-oxo-1,2,4-triazole-5-carboxamide N-hydroxy-
INERT INGREDIENTS

1.30%
98.62%

KEEP OUT OF REACH OF CHILDREN
WARNING

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN wash with soap and water.
IF IN EYES flush with plenty of water; get
medical attention.
See Back Panel for Additional Precautionary
Statements.

For sale to, use and storage by professional turf maintenance personnel
only. Not for use on home lawns. For use on ornamental turf.

Net Weight 50 Lbs. (22.75 kg)
Treats Up to 17,000 Sq. Ft.

EPA Reg. No. 9198-75

EPA Est. No. 9198-04.1

Net Weight 50 Lbs. (22.75 kg)
Treats Up To 17,000 Sq. Ft.

2 of 2

Tee Time™ Fertilizer 20-4-10 with RONSTAR®

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
15% Ammoniacal Nitrogen	
12.5% Urea Nitrogen	
6.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	10%
Sulfur (S)	4.5%

Primary Plant Foods Derived from: Ammoniacal Phosphates, Urea, Murate of Potash and Sulfur Coated Urea
Sulfur derived from Sulfur Coated Urea
Potential Acidity 653 pounds Calcium Carbonate Equivalent per ton

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes eye and skin irritation. Avoid contact with skin, eyes, or clothing. Avoid contamination of feed and feed-stuffs. Keep opened containers away from children and animals. Wash clothing and equipment after handling this product.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with labeling.

STORAGE AND DISPOSAL

STORAGE:

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or fuels. In the event of spill, sweep up remaining dust and use or dispose of according to directions.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION: Tee Time 20-4-10 with RONSTAR is a selective pre-emergent herbicide for control of many annual grasses and broadleaf weeds in ornamental turf on a wide variety of soils. Ronstar is a selective post-emergent herbicide for control of many annual grasses and broadleaf weeds in ornamental turf on a wide variety of soils. The product controls weeds by killing the young seedlings before they become established. Ronstar is a selective post-emergent herbicide for control of many annual grasses and broadleaf weeds in ornamental turf on a wide variety of soils. The product controls weeds by killing the young seedlings before they become established.

Do not apply to wet turf as deterioration may occur.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use. **ORNAMENTAL TURF INCLUDING GOLF COURSE FAIRWAYS, TEES, AND PARKS:** Established perennial Bluegrass, Bermudagrass, Perennial Ryegrass, Tall Fescue, Zoysa and St. Augustinegrass turf.

For most ornamental use, rates of Tee Time 20-4-10 with Ronstar should be based on pre-emergent weed population, time of application and length of growing season. A well-maintained turf with the weed infestation will require less herbicide for effective control than will a thin turf with low fertility. Maximum maintenance and heavy traffic. Ornamental areas with larger growing seasons require higher rates. When natural or practical require application of Tee Time 20-4-10 with Ronstar in advance of weed seed germination, rates should be adjusted upward to compensate for normal breakdown of the active ingredient.

Rate turf and time of season. Mowing a necessary as grasses reach top surface. Measure a 10' x 10' area to be treated and uniformly apply required amount using a fertilizer spreader. Cross treating with one half required amount in a 90° direction will give best results. If an a 90° direction spray application, water turf thoroughly to increase effectiveness. When using rates below 100 pounds of product per acre, thoroughly watering assures more even distribution of the herbicide and "washes" grass in coverage.

AMOUNT TO USE: Crabgrass, Goosegrass, Green and Yellow Foxtail and Sandbar Control. Use the chart below to calculate the most economical rate for use in your turf maintenance program.

SUGGESTED APPLICATION RATES FOR GRASSY WEED CONTROL

Anticipated Grassy Weed Population*	Lib. A	Lib. B	Lib. C	Anticipated Days of Control
High (over 1000 seeds per sq. ft.)	1.75 to 2.00	1.5	2.0	110
Medium	1.5 to 1.75	1.25	1.5	110
Low (under 1000 seeds per sq. ft.)	1.25 to 1.5	1.0	1.25	110

* Estimate weed count on 10' x 10' area or other convenient area. Count only weeds which are visible and growing. Do not count dead or dormant weeds. Do not count weeds which are not growing. Do not count weeds which are not growing.

Annual Grasses (Pre-emergent) and Bermudagrass Control: For pre-emergent control, apply 200 pounds per acre (10 pounds per 1000 square feet) in early spring prior to weed seed germination.

Broadleaf weeds such as Shells, Puffy Spurge, Yorkrass, Spotted, Common Purslane, Florida Purslane, Common Groundsweeper, Carpetgrass, Pigeon, Spiny Tuff. For pre-emergent control of these weeds, apply 200 pounds per acre (10 pounds per 1000 square feet) in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 1.75 pounds per acre (1.25 pounds per 1000 square feet) of Tee Time 20-4-10 with Ronstar. This is normally sufficient for control of St. Augustinegrass and St. Augustinegrass.

Red Fescue and Zoysa are not species of this product. The use of Tee Time 20-4-10 with Ronstar on turf containing Red Fescue or Zoysa should be limited to situations where they are a undesirable grasses.

Do not apply to Bermudagrass, Carpetgrass or newly seeded areas.

Because of long residual activity of Tee Time 20-4-10 with Ronstar, seeding and weeding areas should be delayed until two months after treatment.

Used carefully with flowers and shrubs. This product should not be used on putting greens because of the varying culture practices used.

SUGGESTED SPREADER SETTINGS

Spreader	Speed	Coverage (sq. ft.)	Settings				
			3 lbs. M	4 lbs. M	5 lbs. M	6 lbs. M	7 lbs. M
Compro	3 mph	6 ft.	2%	4	6%	6%	6%
Booms	3 mph	6 ft.	5%	6	6	6%	6%
Loys	3 mph	30 ft.	5	5%	6	6%	7

IMPORTANT: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each treatment.

NOTICE: This product is designed and labeled exclusively for use and application by professional turf maintenance personnel. The manufacturer makes no warranty, express or implied, concerning the product or its use when offered beyond the description on the label. All statements made concerning the product apply only when used as directed.

This product conforms to the chemical description on the label panel and is recommended for use on turf only when used in accordance with directions. Other turf use directions follow directions carefully. Timing and method of application, weather and conditions. Measure with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material and is not responsible for damage to property.

Manufactured by
**the professional's
partner™**

Wear protective clothing including long sleeve shirt and gloves when handling. Wash thoroughly after handling.

Do not use for use

new but efficient weeder
Higher rate ok
354-659



Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537