

9198-75

06/30/1988

1/5

Best Available Copy

TEE TIME
FERTILIZER
20-4-10
with RONSTAR

ACCEPTED
JUN 30 1988
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 9198-75

ACTIVE INGREDIENT:

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-
isopropoxyphenyl)-1,3,4-oxadiazolin-5-one] 1.38%

INERT INGREDIENTS 98.62%
TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN wash with soap and water

IF IN EYES flush with plenty of water; get medical attention

See Back Panel for Additional Precautionary Statements

Net Weight 50 Lbs. (22.75 kg)

Treats Up to 17,000 Sq. Ft.

EPA Reg. No. 9198-75

EPA Est. 9198-OH-1

Net Weight 50 Lbs. (22.75 kg)

Treats Up To 17,000 Sq. Ft.

TEE TIME

FERTILIZER 20-4-10

with RONSTAR

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
1.5% Ammoniacal Nitrogen	
12.5% Urea Nitrogen	
6.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5)	4%
Soluble Potash (K2O)	10%
Sulfur (S)	4.5%

Primary Plant Foods Derived From: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea.

Potential Acidity 653 pounds Calcium Carbonate Equivalent per ton.

PRECAUTIONARY STATEMENTS

WARNING

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Avoid contamination of feed and foodstuffs. Keep opened containers away from children and animals. Wear protective clothing including long-sleeved shirt and gloves when handling. Wash thoroughly after handling. Wash clothing and equipment after handling this product.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not contaminate water by disposing equipment washings. Do not apply directly to water.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: Tee Time 20-4-10 with FONSTAR is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf. This product controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Do not apply to wet turf as discoloration may occur.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

ORNAMENTAL TURF INCLUDING GOLF COURSE FAIRWAYS, TEES AND PARKS Established perennial Bluegrass, Bermudagrass, Perennial Ryegrass, Tall Fescue, Zoysia and St. Augustinegrass turf:

For most economical use, rates of Tee Time 20-4-10 with Fonstar should be based on anticipated weed population, time of application and length of growing season. A well-maintained turf with low weed infestations will require less herbicide for effective control than will a thin turf with low fertility, minimum maintenance and heavy traffic. Geographic areas with longer growing seasons require higher rates. When cultural practices require application of Tee Time 20-4-10 with Fonstar well in advance of weed seed germination, rates should be adjusted upward to compensate for normal breakdown of the active ingredient.

Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results.

If rain is not expected shortly after application, water turf thoroughly to increase effectiveness. When using rates below 180 pounds of product per acre, thoroughly watering assures more even distribution of the herbicide and avoids gaps in coverage.

AMOUNT TO USE: Crabgrass, Goosegrass, Green and Yellow Foxtail and Sandbur
Control: Use the chart below to calculate the most economical rate for use in your turf maintenance program.

SUGGESTED APPLICATION RATES FOR GRASSY WEED CONTROL

Anticipated Grassy Weed Population*	Lbs/A	Lbs/M	Lbs a.i./A	Anticipated Days of Control	
High (thin turf, low fertility, etc.)	175-290	4-6.7	2.4-4	110	200
Medium	153-290	3.5-6.7	2.1-4	110	250
Low (dense turf, history of good fertility and maintenance)	130-290	3-6.7	1.8-4	110	270

In seasons when heavy rains or other environmental factors would prolong weed seed germination, a supplemental application of Tee Time 20-4-10 with Ronstar may be made six to eight weeks after initial treatment. Total product applied should not exceed 290 Lbs./A (4 Lbs. active/A).

Annual Bluegrass (Poa annua) and Barnyardgrass Control: For preemergence control, apply 220 pounds per acre (3 pounds active ingredient) or 5 pounds per 1000 square feet in early spring prior to weed seed germination.

Broadleaf weeds such as Oxalis, Petty Spurge, Veronica Speedwell, Common Purslane, Florida Pursley, Common Groundsel, Carpetgrass, Pigweed, Stinging Nettle: For preemergent control of these weeds, apply 220 pounds per acre (3 pounds per active ingredient) or 5 pounds per 1000 square feet in spring prior to weed seed germination.

St. Augustinegrass turf, do not exceed the rate of 175 pounds per acre (2.4 pounds active ingredient). Tee Time 20-4-10 with Ronstar may cause temporary discoloration of Bermudagrass and St. Augustinegrass. This is normally outgrown within 2-3 weeks.

Red Fescue and Bentgrass are not tolerant of this product. The use of Tee Time 20-4-10 with Ronstar on turf containing Red Fescue or Bentgrass should be limited to situations where they are undesirable grasses.

Do not apply to Dichondra, Centipedegrass or newly seeded areas.

Because of long residual activity of Tee Time 20-4-10 with Ronstar, seeding into treated areas should be delayed until four months after treatment.

Avoid contact with flowers and shrubs. This product should not be used on putting greens because the varying cultural practices used.

Apply with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

SUGGESTED SPREADER SETTINGS

(available at label printing)

IMPORTANT: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds. In the event of spill, sweep up minimizing dust and use or dispose of according to directions. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

MANUFACTURER'S WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured by:

THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537